

**PINK BISMUTH- bismuth subsalicylate liquid**  
**Ger-Care Pharmaceuticals, Corp**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**GC PEPTO**

**Active ingredient (in each 15 mL Tablespoon = TBSP)**

Bismuth subsalicylate 262 mg

**Purposes**

upset stomach reliever and antidiarrheal

**Uses**

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea

**Warnings**

**Reye's syndrome:**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use** if you have

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool

**Ask a doctor or pharmacist before use if you are** taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout

- arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if**

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- shake well before using
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- for accurate dosing, use dose cup

**adults and children 12 years and over**

- take 2 TBSP every 1/2 hour to 1 hour as needed
- do not exceed 16 TBSP in 24 hours
- use until diarrhea stops but not more than 2 days

**children under 12 years:**ask a doctor

**Other information**

- **each TBSP contains:** sodium 6 mg. salicylate 130mg
- low sodium
- sugar free
- protect from freezing
- avoid excessive heat (over 104F or 40C)
- store at room temperature
- **TAMPER-EVIDENT: Do not use if imprinted neckband is missing or broken.**
- for health information visit [www.info-xtra.com](http://www.info-xtra.com) for health information

**Inactive ingredients**

benzoic acid, D and C red no. 22, D and C red no. 28, flavor, magnesium aluminum silicate, methylcellulose, purified water, saccharin sodium, salicylic acid, simethicone emulsion, sodium salicylate, sorbic acid, xanthan gum

**package Label**

GERICARE

NDC 57896-392-16

pink bismuth

regular strength

bismuth subsalicylate 262mg/ 15mL

upset stomach reliever/antidiarrheal

soothing relief for 5 symptoms

heartburn  
 indigestion  
 upset stomach  
 nausea  
 diarrhea

compare to the active ingredient of Pepto-Bismol  
 16 FL OZ (1 PT) (473 mL)

SEALED WITH PRINTED NECKBAND FOR YOUR PROTECTION

**Drug Facts** (continued)

**Directions**

- shake well before using
- for accurate dosing, use dose cup
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- use until diarrhea stops but not more than 2 days
- adults and children 12 years and over: take 2 TBSP every 1/2 to 1 hour as needed
- do not exceed 16 TBSP in 24 hours
- children under 12 years: ask a doctor

**Other Information**

- each tablespoon contains: sodium 6 mg
- salicylate 130 mg
- low sodium
- sugar free
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)
- visit [www.info-xtra.com](http://www.info-xtra.com) for health information.
- **TAMPER-EVIDENT:** Do not use if imprinted neckband is missing or broken.

**Inactive ingredients** benzoic acid, flavor, magnesium aluminum silicate, methylcellulose, purified water, red 22, red 28, saccharin sodium, salicylic acid, simethicone emulsion, sodium salicylate, sorbic acid, xanthan gum.

Distributed by: Gericare Pharmaceuticals Corp.  
 1650 63<sup>rd</sup> Street, Brooklyn, NY 11204

\*Geri Care is not affiliated with the owner of the registered trademark Pepto-Bismol®.



NDC 57896-392-16

**GERI CARE**

**PINK REGULAR STRENGTH BISMUTH**

**BISMUTH SUBSALICYLATE 262 mg/15 mL  
 UPSET STOMACH RELIEVER/ANTIDIARRHEAL**

**Soothing Relief for 5 Symptoms**

- Heartburn
- Indigestion
- Upset Stomach
- Nausea
- Diarrhea

*\*Compare to the Active Ingredient of Pepto-Bismol®*

16 FL OZ (1 PT) (473 mL)

**Drug Facts**

**Active ingredient** (In each 15 mL Tablespoon = TBSP)  
 Bismuth subsalicylate 262 mg ..... Upset stomach reliever and antidiarrheal

**Purposes**

**Uses** relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use** if you have

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for**

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if**

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

310-02111-0 REV1113

## PINK BISMUTH

bismuth subsalicylate liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:57896-392
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ, BISMUTH CATION - UNII:ZS9CD1I8YE)	BISMUTH SUBSALICYLATE	262 mg in 15 mL

### Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC)	
METHYLCELLULOSE (100 CPS) (UNII: 4GFU244C4J)	
WATER (UNII: 059QF0K00R)	

SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SALICYLIC ACID (UNII: O414PZ4LPZ)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SODIUM SALICYLATE (UNII: WIQ1H85SYP)	
SORBIC ACID (UNII: X045WJ989B)	
XANTHAN GUM (UNII: TTV12P4NEE)	
METHYL SALICYLATE (UNII: LAV5U5022Y)	

**Product Characteristics**

<b>Color</b>	pink	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	MINT	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57896-392-08	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/01/2012	11/01/2019
2	NDC:57896-392-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/01/2012	09/01/2019

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part335	01/01/2000	11/01/2019

**Labeler** - Geri-Care Pharmaceuticals, Corp (611196254)

**Registrant** - GCP Laboratories (965480861)

**Establishment**

Name	Address	ID/FEI	Business Operations
GCP LABORATORIES		965480861	manufacture(57896-392) , pack(57896-392)