

**VICKS NYQUIL COUGH DM PLUS CONGESTION- dextromethorphan hydrobromide, doxylamine succinate and phenylephrine hydrochloride liquid**  
**The Procter & Gamble Manufacturing Company**

-----  
**VICKS® NyQuil™**  
**COUGH DM + CONGESTION**

***Drug Facts***

**Active ingredients (in each 15 mL)**

Dextromethorphan HBr 10 mg

Doxylamine succinate 6.25 mg

Phenylephrine HCl 5 mg

**Purpose**

Cough suppressant

Antihistamine

Nasal decongestant

**Uses**

temporarily relieves cold symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

**Warnings**

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

**Ask a doctor before use if you have**

- heart disease
- high blood pressure

- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are** taking sedatives or tranquilizers.

### **When using this product**

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- symptoms do not improve within 7 days or occur with a fever
- cough persists for more than 7 days, comes back or occurs with a fever, rash, or persistent headache.

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

- take only as directed
- only use dose cup provided
- do not exceed 4 doses per 24 hrs

---

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	not use unless directed by a doctor
children under 4 yrs	do not use

---

### **Other information**

- **each 15 mL dose cup contains:** sodium 44 mg

- store at no greater than 25°C

## Inactive ingredients

citric acid, FD&C Blue No. 1, FD&C Red No. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

## Questions?

**1-800-362-1683**

TAMPER EVIDENT: Do not use if printed shrinkband is broken or missing.

**DIST. BY PROCTER & GAMBLE,  
CINCINNATI OH 45202**

## PRINCIPAL DISPLAY PANEL - 236 ml Bottle Label

**VICKS®**

**NyQuil™**

**COUGH DM + CONGESTION**

*Dextromethorphan HBr, Phenylephrine HCl, Doxylamine Succinate*

- **Cough**
- **Nasal Congestion**
- **Sneezing, Runny Nose**

**8 FL OZ (236 ml)**

*Nighttime Relief*

**TAMPER EVIDENT:**  
Do not use if printed shrinkband is broken or missing.  
← PEEL BACK FOR DRUG FACTS

**Drug Facts**  
**Active ingredients (in each 15 mL)** Purpose  
Dextromethorphan HBr 10 mg Cough suppressant  
Doxylamine succinate 6.25 mg Antihistamine  
Phenylephrine HCl 5 mg Nasal decongestant

**Uses** temporarily relieves common cold symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

**PARENTS:**  
Learn about how to use this medicine safely.  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

DIST. BY PROCTER & GAMBLE,  
CINCINNATI OH 45202  
Patents: [www.pg.com/patents](http://www.pg.com/patents)  
[www.vicks.com](http://www.vicks.com)

**P&G**  
[www.pg.com](http://www.pg.com)  
90889547

Discard from Recycle Bin  
PLASTIC BOTTLE

2390004120 2

**Drug Facts (continued)**  
**Warnings**  
**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers**

**When using this product**

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

**Drug Facts (continued)**

- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- symptoms do not improve within 7 days or occur with a fever
- cough persists for more than 7 days, comes back or occurs with a fever, rash, or persistent headache

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	do not use unless directed by a doctor
children under 4 yrs	do not use

**Other information**

- each 15 mL contains: sodium 44 mg
- store at no greater than 25°C

**Inactive ingredients** citric acid, FD&C Blue No. 1, FD&C Red No. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

**Questions? 1-800-362-1683**

# VICKS NYQUIL COUGH DM PLUS CONGESTION

dextromethorphan hydrobromide, doxylamine succinate and phenylephrine hydrochloride liquid

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69423-999
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg in 15 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>SORBITOL</b> (UNII: 506T60A25R)	

## Product Characteristics

<b>Color</b>	red	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-999-08	236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/08/2020	
2	NDC:69423-	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a	07/08/2020	

999-12

Combination Product

07/08/2020

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M012	07/08/2020	

**Labeler** - The Procter & Gamble Manufacturing Company (004238200)**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
R L FINE CHEM		871410195	api manufacture(69423-999)

Revised: 10/2023

The Procter &amp; Gamble Manufacturing Company