

PHYSICIANS CARE MOTION SICKNESS- meclizine hcl tablet
Acme United Corporation

PhysiciansCare Motion Sickness

Drug Facts

Active ingredient (in each tablet)

Meclizine HCl 25 mg

Purpose

Antiemetic

Uses

for the prevention and treatment of the nausea, vomiting, or dizziness associated with motion sickness.

Warnings

Do not use for children under 12 years of age unless directed by a doctor.

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- glaucoma
- a breathing problems such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product

- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center

(1-800-222-1222) right away.

Directions

- to prevent motion sickness, take the first dose $\frac{1}{2}$ hour to 1 hour before starting activity
- to treat motion sickness, take at first signs of symptoms

Adults and children: (12 years and older) 1 to 2 tablets once daily, or as directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Other information

- tamper-evident sealed packets
- do not use any opened or torn packets
- store at 77°F (25°C); excursions permitted between 59- 86° F (15-30°C)
- protect from heat and humidity
- use by expiration date on packet

Inactive ingredients

corn starch, D&C yellow # 10 aluminum lake, lactose anhydrous, magnesium stearate, silicon dioxide

Questions or comments? 1-800-835-2263

<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; border-radius: 15px; padding: 5px; font-weight: bold;">Physicians <i>CARE.</i></div> <div style="font-weight: bold;">91274</div> </div> <p style="text-align: center; font-size: 1.2em; margin-top: 20px;">Motion Sickness</p> <p style="text-align: center; font-size: 1.1em;">Meclizine HCl 25 mg, Antiemetic 6 Packets (2 tablets each)</p> <div style="display: flex; justify-content: space-between; align-items: flex-end; margin-top: 20px;"> <div style="font-size: 0.8em;"> <p>Manufactured for: © 2021 Acme United Corporation 1 Waterview Dr, Shelton, CT 06484 www.FirstAidOnly.com 807597-revA</p> </div> <div style="text-align: center;">  <p>0 92265 91274 7</p> </div> </div>	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td colspan="2">Drug Facts</td> </tr> <tr> <td style="width: 70%;">Active ingredient (in each tablet)</td> <td style="text-align: right;">Purpose</td> </tr> <tr> <td>Meclizine Hydrochloride 25 mg</td> <td style="text-align: right;">Antiemetic</td> </tr> <tr> <td colspan="2">Uses for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness.</td> </tr> <tr> <td colspan="2">Warnings</td> </tr> <tr> <td colspan="2">Do not use for children under 12 years of age unless directed by a doctor.</td> </tr> <tr> <td colspan="2">Ask a doctor before use if you have</td> </tr> <tr> <td colspan="2"> <ul style="list-style-type: none"> ■ difficulty in urination due to enlargement of the prostate gland ■ glaucoma ■ a breathing problem such as emphysema or chronic bronchitis </td> </tr> <tr> <td colspan="2">Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.</td> </tr> <tr> <td colspan="2">When using this product</td> </tr> <tr> <td colspan="2"> <ul style="list-style-type: none"> ■ drowsiness may occur ■ avoid alcoholic beverages ■ alcohol, sedatives and tranquilizers may increase drowsiness ■ use caution when driving a motor vehicle or operating machinery </td> </tr> <tr> <td colspan="2">if pregnant or breastfeeding, ask a health professional before use.</td> </tr> <tr> <td colspan="2">Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1.800.222.1222) right away.</td> </tr> <tr> <td colspan="2">Directions</td> </tr> <tr> <td colspan="2"> <ul style="list-style-type: none"> ■ to prevent motion sickness, take the first dose ½ hour to 1 hour before starting activity ■ to treat motion sickness, take at first signs of symptoms </td> </tr> <tr> <td style="width: 50%;">Adults and children: (12 years and older)</td> <td style="width: 50%;">1 to 2 tablets once daily, or as directed by a doctor.</td> </tr> <tr> <td>Children under 12 years</td> <td>Do not give to children under 12 years of age.</td> </tr> <tr> <td colspan="2">Other information</td> </tr> <tr> <td colspan="2"> <ul style="list-style-type: none"> ■ tamper-evident sealed packets ■ do not use any opened or torn packets ■ store at 77°F (25°C); excursions permitted between 59°-86°F (15°-30°C) ■ protect from heat and humidity ■ use by expiration date on packet </td> </tr> <tr> <td colspan="2">Inactive ingredients</td> </tr> <tr> <td colspan="2">corn starch, D&C yellow #10 aluminum lake, lactose anhydrous, magnesium stearate, silicon dioxide</td> </tr> <tr> <td colspan="2">Questions? 800.835.2263</td> </tr> </table>	Drug Facts		Active ingredient (in each tablet)	Purpose	Meclizine Hydrochloride 25 mg	Antiemetic	Uses for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness.		Warnings		Do not use for children under 12 years of age unless directed by a doctor.		Ask a doctor before use if you have		<ul style="list-style-type: none"> ■ difficulty in urination due to enlargement of the prostate gland ■ glaucoma ■ a breathing problem such as emphysema or chronic bronchitis 		Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.		When using this product		<ul style="list-style-type: none"> ■ drowsiness may occur ■ avoid alcoholic beverages ■ alcohol, sedatives and tranquilizers may increase drowsiness ■ use caution when driving a motor vehicle or operating machinery 		if pregnant or breastfeeding , ask a health professional before use.		Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1.800.222.1222) right away.		Directions		<ul style="list-style-type: none"> ■ to prevent motion sickness, take the first dose ½ hour to 1 hour before starting activity ■ to treat motion sickness, take at first signs of symptoms 		Adults and children: (12 years and older)	1 to 2 tablets once daily, or as directed by a doctor.	Children under 12 years	Do not give to children under 12 years of age.	Other information		<ul style="list-style-type: none"> ■ tamper-evident sealed packets ■ do not use any opened or torn packets ■ store at 77°F (25°C); excursions permitted between 59°-86°F (15°-30°C) ■ protect from heat and humidity ■ use by expiration date on packet 		Inactive ingredients		corn starch, D&C yellow #10 aluminum lake, lactose anhydrous, magnesium stearate, silicon dioxide		Questions? 800.835.2263	
Drug Facts																																													
Active ingredient (in each tablet)	Purpose																																												
Meclizine Hydrochloride 25 mg	Antiemetic																																												
Uses for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness.																																													
Warnings																																													
Do not use for children under 12 years of age unless directed by a doctor.																																													
Ask a doctor before use if you have																																													
<ul style="list-style-type: none"> ■ difficulty in urination due to enlargement of the prostate gland ■ glaucoma ■ a breathing problem such as emphysema or chronic bronchitis 																																													
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.																																													
When using this product																																													
<ul style="list-style-type: none"> ■ drowsiness may occur ■ avoid alcoholic beverages ■ alcohol, sedatives and tranquilizers may increase drowsiness ■ use caution when driving a motor vehicle or operating machinery 																																													
if pregnant or breastfeeding , ask a health professional before use.																																													
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1.800.222.1222) right away.																																													
Directions																																													
<ul style="list-style-type: none"> ■ to prevent motion sickness, take the first dose ½ hour to 1 hour before starting activity ■ to treat motion sickness, take at first signs of symptoms 																																													
Adults and children: (12 years and older)	1 to 2 tablets once daily, or as directed by a doctor.																																												
Children under 12 years	Do not give to children under 12 years of age.																																												
Other information																																													
<ul style="list-style-type: none"> ■ tamper-evident sealed packets ■ do not use any opened or torn packets ■ store at 77°F (25°C); excursions permitted between 59°-86°F (15°-30°C) ■ protect from heat and humidity ■ use by expiration date on packet 																																													
Inactive ingredients																																													
corn starch, D&C yellow #10 aluminum lake, lactose anhydrous, magnesium stearate, silicon dioxide																																													
Questions? 800.835.2263																																													

label image

PHYSICIANS CARE MOTION SICKNESS			
meclizine hcl tablet			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0924-1008(NDC:47682-481)
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	25 mg	

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	44;403
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0924-1008-00	2 in 1 PACKET; Type 0: Not a Combination Product	06/18/2021	
2	NDC:0924-1008-02	6 in 1 BAG	06/18/2021	
2		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	06/18/2021	

Labeler - Acme United Corporation (001180207)

Establishment

Name	Address	ID/FEI	Business Operations
Acme United Corporation		045924339	relabel(0924-1008) , repack(0924-1008)

Establishment

Name	Address	ID/FEI	Business Operations
Acme United Corporation		080119599	relabel(0924-1008) , repack(0924-1008)