CAREALL ACETAMINOPHEN- acetaminophen tablet New World Imports, Inc

Drug Facts

Active Ingredient

Acetaminophen 325mg

Purpose

Pain reliever / Fever reducer

Keep out of Reach of Children

Keep out of reach of children. Overdose warning: In case of overdose, contact a doctor or Poison Control Center immediately. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Uses

- Temporary relief of minor aches and pains associated with: headache, muscular aches, backache, minor pain from arthritis, the common cold, toothache, premenstrual and menstrual cramps
- Temporarily reduces fever

Liver warning: This product contains acetaminophen. The manxium daily dose of this product is 10 tablets (3,250 mg) in 24 hours fro adults or 5 tablets (1,625 mg) in 24 hours for children.

Severe liver damage may occur if:

- Adult takes more than 4,000mg of acetaminphen in 24 hours
- Child takes more than 5 doses in 24 hours
- Taken with other drugs containing acetaminophen
- Adult has 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters or rash. If a skin reaction occurs, stop use and seek medical help right away.

Do not use:

- with any other drug containing acetaminophen (prescription or nonprescription).
- If you are allergic to acetaminophen.

Ask a doctor before use if user has: liver disease

Ask a doctor or pharmacist before use if:

• the user is taking the blood thinnig drug warfarin.

• is a child with pain of arthritis

Stop use and ask a doctor if:

- Adults pain gets worse or lasts for more than 10 days.
- Childs pain gets worse or last fr more than 5 days.
- Fever gets worse or lasts for more than 3 days.
- New symptoms occur.
- Redness or swelling is present.

If pregnant or breast-feeding, ask a health professional before use.

Overdose warning: taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical action is critical for adults as well as for children, even if you do not notice any signs or symptoms.

Adults and children 12 years of age and older: Take 2 tablets every 4 to 6 hours while symptoms last. Do not take more than 10 tablets in 24 hours, unless directed by a doctor. do not take for more than 10 days unless directed by a doctor.

Children 6-11 years of age: Take 1 tablet every 4 to 6 hours while symptoms last. Do not take more than 5 tablets in 24 hours. Do not use more than 5 days unless directed by a doctor.

Children under 6 years of age: Ask a doctor

Corn starch, povidone, pregelatinized starch, sodium starch glycolate, stearic acid

0

ij



Acetaminophen

Regular Strength

► Pain Reliever

100

Fever Reducer

*Compare to the active ingredient of Regular Strength Tylenol®



DO NOT USE IF IMPRINTED SAFETY SEA UNDER CAP IS BROKEN OR MISSING

Active ingredient (in each tablet) Orug Facts

Purpose

For the temporarily relief of minor aches and pains Uses

■ backache ■ minor pain from arthritis ■ the common cold premenstrual and menstrual cramps associated with ■ headache ■ muscular aches ■ toothache

temporarily reduces fever

Children

under 12

years to Children 6

maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children. 4,000mg of acetaminophen in 24 hours ■ child takes more than ■ taken with other drugs containing acetaminophen ■ skin reddening
■ blisters
■ rash Allergy alert: Acetaminophen may cause severe skin reactions. If a skin reaction occurs, stop use and seek medical help right 3 or more alcoholic drinks every day while using this product adult takes more than 5 doses in 24 hours, which is the maximum daily amount Liver warning: This product contains acetaminophen. Severe liver damage may occur if Symptoms may include:

years and

children 12

Adults and

with any other drug containing acetaminophen prescription or non-prescription). If you are not sure whether a ■ if you are allergic to acetaminophen or any of the inactive drug contains acetaminophen, ask a doctor or pharmacist ingredients in this product Do not use away.

Drug Facts (continued on inside) This product is not distributed or manufactured by Johnson & Johnson Consumer Inc., owner of the registered trademark Regular Strength

Distributed By: CareALL Products, 160 Athens Way, Nashville, TN 37228 www.careallproducts.com

Stop use and ask a doctor if adult's pain gets worse or Inactive ingredients com starch, povidone, pregelatinized starch, sodium starch glycolate, stearic acid Other information 20°-25°C (68°-77°F) overdose wa contact a Poison Control Center (1-800-222-1222) right more than 5 days ■ fever gets worse or lasts more than 3 lasts more than 10 days child's pain gets worse or lasts Ask a doctor or pharmacist before use if ■ the user is taking the blood thinning drug warfarin ■ is a child with Drug Facts (continued) Monday through Friday 9AM - 5PM Questions/Comments for children even if you do not notice any signs or symptoms away. Quick medical attention is critical for adults as well as Overdose warning: In case of overdose, get medical help or petore use. Keep out of reach of children. pain of arthritis Ask a doctor before use if the user has liver disease ymptoms occur JIFECTIONS ■ do not take more than directed (see pregnant or breast-feeding, ask a health professional ■ redness or swelling is present ■ any new doctor ■ do not use for more than 10 days ■ take 1 tablet every 4 to 6 hours while symptoms last ■ do not take more than 10 tablets in 24 hours, unless directed by a ■ take 2 tablets every 4 to 6 hours while than 5 days unless directed by a doctor symptoms last ■ do not take more than 5 unless directed by a doctor ablets in 24 hours. do not use for more store at room temperature Call 1-888-952-0050

CAREALL ACETAMINOPHEN

acetaminophen tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51824-061
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
Acetaminophen (UNII: 36209ITL9D) (Acetaminophen - UNII:36209ITL9D)	Acetaminophen	325 mg	

Inactive Ingredients		
Ingredient Name	Strength	
Starch, Corn (UNII: O8232NY3SJ)		
Stearic Acid (UNII: 4ELV7Z65AP)		

Povidone (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A CORN (HNII: AG9R65PV6R)	

Product Characteristics			
Color	white (white)	Score	no score
Shape	ROUND (Round Circle)	Size	11mm
Flavor		Imprint Code	AZ;010
Contains			

ı	P	Packaging			
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:51824-061- 02	24 in 1 CASE	04/15/2016	
	1	NDC:51824-061- 01	100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/15/2016	

Labeler - New World Imports, Inc (075372276)

Revised: 11/2023 New World Imports, Inc