DR. DEAN- chlorhexidine gluconate liquid solution 2% solution USA Medical Supply LLC

Drug Facts

Active ingredient

Chlorhexidine gluconate solution, 2%

Purposes

Surgical hand scrub

Healthcare personnel handwash

Skin wound and general skin cleanser

Uses

- **surgical hand scrub:** significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- **healthcare personnel handwash:** handwash to help reduce bacteria that potentially can cause disease
- skin wound and general cleanser: helps reduce bacteria on the skin

Warnings

For external use only

Allergy alert

This product may cause a severe allergic reaction.

Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- if you are allergic to chlorhexidine gluconate or any other ingredients in this preparation
- as a properative skin preparation (especially on the head or face)
- in contact with meninges
- in the genital area

When using this product

- keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if
 placed or kept in the eye during surgical during surgical procedures or may cause
 deafness when instilled in the middle ear through perforated ear drums.
- if contact occurs, rinse with cold water right away
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be done except when underlying condition makes it necessary to reduce the bacterial population of the skin

Stop use and ask a doctor if

• irritation, sensitization, or allergic reaction occurs and lasts for 72 hours. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These product may cause irritation or chemical burns.
- use full strength
- do not dilute

Surgical hand scrub

- remove jewelry
- wet hands and forearms with water and apply 5 mL of the product
- wash/scrub hands and forearms for 3 minutes paying particular attention to the nails, cuticles, and interdigital spaces
- rinse thoroughly with water
- washfor an additional 3 minutes with 5 mL of the product and rinse under running water
- dry thoroughly

Healthcare personnel handwash

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 30 seconds
- rinse and dry thoroughly

Skin wound and general skin cleanser

- thoroughly rinse the area to be cleansed with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently. Rinse thoroughly with water.
- rinse again thoroughly

Other information

- store between 20-25 °C (68-77 °F)
- avoid excessive heat (above 104 °F/40 °C)
- for additional information, see Safety Data Sheet (SDS)

• if swallowed get medical help or contact a Poison Control Center immediately

Inactive ingredients cocamide DEA, cocamine oxide, fragrance, gluconic acid*, gluconolactone*, hydroxyethylcellulose, isopropyl alcohol 4%, PEG-75 lanolin, PEG-150 distearate, propylene glycol, quaternium-60, water

*contains one or more of these ingredients

☐ Questions or comments? call 1.800.581.2528, 24 hours a day, 7 days a week

Representative label and principal display panel

Dr. Dean

Chlorhexidine Gluconate

Liquid Solution

2.0%

Up to 6 Hour Protection

Antimicrobial Skin Soap

FOR EXTERNAL USE ONLY

Distributed by USA Medical Supply

Northbrook IL 60062

www.usamdsupply.com

4 FL OZ (118 mL)

NDC 81657-002-01

MADE IN THE USA

FDA APPROVED



Drug Facts

Active ingredient Chlorhexidine gluconate

Purposes

solution, 2%

Healthcare personnel handwash Surgical hand scrul

Skin wound and general skin cleanse

Uses

surgical hand scrub: significantly reduces the

number of microorganisms on the hands and forearm

skin wound and general skin cleanser: helps reduce healthcare personnel handwash: handwash to help reduce bacteria that potentially can cause disease prior to surgery or patient care

Lift Here For Drug Facts

bacteria on the skin

Drug Facts (continued)

Skin wound and general skin cleanser: thoroughly rinse the area to be cleansed with water apply the minimum amount of product necessary to

thoroughly with water cover the skin or wound area and wash gently. Rinse

Other information

rinse again thoroughly

store between 20-25°C (68-77°F) avoid excessive heat (above 104°F/40°C)

If swallowed get medical help or contact a Poison for additional information, see Safety Data Sheet (SDS Control Center immediately

Do not use

help right away.

gluconolactone*, hydroxyethyl cellulose, isopropyl alcohol 4%, PEG-75 lanolin, PEG-150 distearate, propylene cocamine oxide, fragrance, gluconic acid* nactive ingredients cocamide DEA

Questions or comments? call 1.800.581.2528, 24 hours a day, 7 days: contains one or more of these ingredients /col, quaternium-60, water

> When using this product in contact with meninges

in the genital area on the head or face)

keep out of eyes, ears, and mouth. May cause serious

and permanent eye injury if placed or kept in the eye

during surgical procedures or may cause deafness wher instilled in the middle ear through perforated eardrums.

 as a patient preoperative skin preparation (especially if you are allergic to chlorhexidine gluconate or any

other ingredient in this preparation

repeated general skin cleansing of large body areas

of the skin should not be routinely treated wounds which involve more than the superficial layers if contact occurs, rinse with cold water right away

Keep from freezing

Directions

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

or allergic reaction occurs and lasts for 72 hours. These Stop use and ask a doctor if irritation, sensitization

population of the skin

condition makes it necessary to reduce the bacterial should not be done except when an underlying

may be signs of a serious condition.

use full strength use with care in premature infants or infants under 2 chemical burns months of age. These products may cause irritation

do not dilute

Surgical hand scrub:

remove jewelry

wash/scrub hands and forearms for 3 minutes paying wet hands and forearms with water and apply 5 mL the product particular attention to the nails, cuticles, and

 dry thoroughly wash for an additional 3 minutes with 5 mL of the product and rinse under running water

rinse thoroughly with water

interdigital spaces

dispense about 5 mL of product into cupped hands rinse and dry thoroughly and wash in a vigorous manner for 30 seconds

Healthcare personnel handwash:

wet hands with water

Drug Facts (continued

Warnings

For external use only

Symptoms may include: Allergy alert his product may cause a severe allergic reaction

wheezing/difficulty breathing hives

facial swelling

shock

If an allergic reaction occurs, stop use and seek medical

Dr. Dean Chlorhexidine Gluconate **Liquid Solution**

Up to 6 Hour Protection

Antimicrobial Skin Soap

FOR EXTERNAL USE ONLY

Distributed by USA Medical Supply

Northbrook IL 60062


www.usamdsupply.com


8 FL OZ (236 mL)

NDC 81657-002-02

MADE IN THE USA

FDA APPROVED







Dr. Dean
Chlorhexidine Gluconate
Liquid Solution
2.0%

Up to 6 Hour Protection

Antimicrobial Skin Soap

FOR EXTERNAL USE ONLY

Distributed by USA Medical Supply

Northbrook IL 60062


www.usamdsupply.com

128 FL OZ (3785 mL)

NDC 81657-002-03

MADE IN THE USA

FDA APPROVED



DR. DEAN

chlorhexidine gluconate liquid solution 2% solution

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:81657-002(NDC:42939-127)
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII: R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	20 mg in 1 mL	

Inactive Ingredients		
Ingredient Name	Strength	
COCO DIETHANOLAMIDE (UNII: 92005F972D)		
COCAMINE OXIDE (UNII: QWA2IZI6FI)		
GLUCONIC ACID (UNII: R4R8J0Q44B)		
GLUCONOLACTONE (UNII: WQ29KQ9POT)		
HYDROXYETHYL CELLULOSE (3000 CPS AT 1%) (UNII: 7Q6P4JN1QT)		
ISOPROPYL ALCOHOL (UNII: ND2M416302)		
PEG-75 LANOLIN (UNII: 091790X7TB)		

PEG-150 DISTEARATE (UNII: 6F36Q0I0AC)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
QUATERNIUM-33 (UNII: XPS4174QZJ)	
WATER (UNII: 059QF0KO0R)	

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:81657- 002-01	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/22/1986		
2	NDC:81657- 002-03	3785 mL in 1 JUG; Type 0: Not a Combination Product	07/22/1986		
3	NDC:81657- 002-02	236 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/22/1986		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA019258	07/22/1986	

Labeler - USA Medical Supply LLC (117790416)

Revised: 6/2021 USA Medical Supply LLC