

**LACTATED RINGERS- lactated ringers injection, solution
MWI (VetOne)**

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Lactated Ringers

INDICATIONS:

For the correction of electrolyte depletion, metabolic acidosis and dehydration of cattle, calves, horses, sheep and swine.

DOSAGE AND ADMINISTRATION:

May be injected intravenously, subcutaneously or intraperitoneally (except in horses) using strict aseptic technique.

Cattle and Horses: 2 to 5 mL per pound of body weight, depending on size and condition of animal, repeated 1 to 3 times daily or as needed.

Swine and Sheep: 2 to 5 mL per pound of body weight depending on size and condition of animal, repeated 1 to 3 times daily or as needed.

If administered subcutaneously divide dosage into several sites of injection and massage points of injection to aid in absorption and help prevent inflammation and/or sloughing.

Distributed by: MWI
Boise, ID 83705
(888) 694-8381
www.vetone.net

Manufactured by: Nova-Tech, Inc.
Grand Island, NE 68801

Rev. 04/12

V1 501033

Net Contents: 1000 mL

RMS 92-1007

18-803-60

LOT NO.: EXP. DATE:

NDC 13985-803-60

1000 mL

MADE IN USA

Sterile - Nonpyrogenic Solution

For animal use only.

Keep out of reach of children.

CAUTION:

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

COMPOSITION:

Each 100 mL contains:

Sodium Chloride.....	600 mg
Sodium Lactate.....	310 mg
Potassium Chloride.....	30 mg
Calcium Chloride Dihydrate.....	20 mg
Water for Injection.....	q.s.

The Calcium, Potassium and Sodium contents are approximately 2.7, 4.0, and 130 mEq/liter, respectively. Total Osmolar Concentration: 269 mOsm/l per liter (calculated).

CAUTION:

Solution should be warmed to body temperature prior to administration and administered at a slow rate. This is a single dose unit. It contains no preservatives. If entire contents are not used, discard unused portion.

WARNING:

Do not administer to horses by intraperitoneal injection. Do not administer to animals with inadequate renal function. Not for use in lactic acidosis.

STORAGE:

Store between 15° - 30°C (59° - 86°F).

TAKE TIME OBSERVE LABEL DIRECTIONS

INDICATIONS: For the correction of electrolyte depletion, metabolic acidosis and dehydration of cattle, calves, horses, sheep and swine.

DOSAGE AND ADMINISTRATION: May be injected intravenously, subcutaneously or intraperitoneally (except in horses) using strict aseptic technique.

Cattle and Horses: 2 to 5 mL per pound of body weight, depending on size and condition of animal, repeated 1 to 3 times daily or as needed.

Swine and Sheep: 2 to 5 mL per pound of body weight depending on size and condition of animal, repeated 1 to 3 times daily or as needed.

If administered subcutaneously divide dosage into several sites of injection and massage points of injection to aid in absorption and help prevent inflammation and/or sloughing.

Distributed by: MWI
Boise, ID 83705
(888) 694-8381
www.vetone.net

Manufactured by: Nova-Tech, Inc.
Grand Island, NE 68801

Rev. 04/12



NDC 13985-803-60

1000 mL



Lactated Ringer's

Injection, USP

Sterile – Nonpyrogenic Solution

Keep out of reach of children.
For animal use only.

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

V1 501033

Net Contents: 1000 mL

COMPOSITION: Each 100 mL contains:

Sodium Chloride 600 mg
Sodium Lactate..... 310 mg
Potassium Chloride..... 30 mg
Calcium Chloride Dihydrate..... 20 mg
Water for Injectionq.s.

The Calcium, Potassium and Sodium contents are approximately 2.7, 4.0, and 130 mEq/liter, respectively. Total Osmolar Concentration: 269 mOsm/l per liter (calculated).

CAUTION: Solution should be warmed to body temperature prior to administration and administered at a slow rate. This is a single dose unit. It contains no preservatives. If entire contents are not used, discard unused portion.

WARNING: Do not administer to horses by intraperitoneal injection. Do not administer to animals with inadequate renal function. Not for use in lactic acidosis.

STORAGE: Store between 15°–30°C (59°–86°F).

RMS 92-1007

18-803-60



OBSERVE LABEL DIRECTIONS

LOT NO.:

EXP. DATE:

LACTATED RINGERS

lactated ringers injection, solution

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:13985-803
Route of Administration	INTRAVENOUS, SUBCUTANEOUS, INTRAPERITONEAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X) (CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	600 mg in 100 mL
SODIUM LACTATE (UNII: TU7HW0W0QT) (LACTIC ACID - UNII:33X04XA5AT)	SODIUM LACTATE	310 mg in 100 mL
POTASSIUM CHLORIDE (UNII: 660YQ98I10) (POTASSIUM CATION - UNII:295O53K152)	POTASSIUM CATION	30 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB)	CALCIUM CHLORIDE	20 mg in 100 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:13985-803-60	1000 mL in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		05/30/2012	

Labeler - MWI (VetOne) (019926120)

Revised: 4/2024

MWI (VetOne)