

**CONEX- dexbrompheniramine maleate, pseudoephedrine liquid**  
**Llorens Pharmaceutical International Division**

-----  
**Llorens-ConexLiq 802**

**Active Ingredients (in each 5 mL tsp.)**

Dexbrompheniramine Maleate, USP ..... 1 mg ..... Antihistamine

Pseudoephedrine HCL, USP ..... 30 mg ..... Nasal Decongestant

**Uses:**

- Temporarily relieves nasal congestion due to common cold, hay fever or other upper respiratory allergies
- Helps decongest sinus openings and sinus passages
- Reduces swelling of nasal passages, shrinks swollen membranes, and temporarily restores freer breathing through the nose
- Temporarily alleviates the following symptoms due to hay fever (allergic rhinitis) : runny nose, sneezing, itching of the nose or throat, itchy and watery eyes.

**Warnings:**

- **Do not use if** you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if you are taking a prescription drug that contains an MAOI, ask your doctor or pharmacist before taking this product.

**Ask a doctor before use if you are** taking sedatives or tranquilizers.

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- difficulty in urination due to the enlargement of the prostate gland.

**When using this product**

- do not exceed the recommended dosage
- excitability may occur, especially in children
- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever.

**Keep out of the reach of children.**In case of overdose, get medical help or contact a Poison Control Center right away.

**If pregnant or breast-feeding,**ask a Health Professional before use.

**Directions:**Do not exceed more than 4 doses in any 24-hour period or as directed by a doctor.

Age	Dose
Adults and children 12 years of age and over	take two teaspoonfuls (10 mL) every 4 to 6 hours
Children 6 to under 12 years of age	take one teaspoonful (5 mL) every 4 to 6 hours
Children under 6 years of age	ask a doctor

**Other Information:**

- Store at controlled room temperature 20-25 degree celcius ( 68-77 degree farenheit); excursions permitted to 15-30 degree clecius (59-86 degree farenheit) (See USP Controlled Room Temperture) Tamper evident by imprinted heat seal under cap
- Do not use if there is evidence of tampering
- **WARNING: Phenylketonuric: Contains 13.5 mg of Phenylalanine per 5 mL (one teaspoonful) dose.**

**Inactive Ingredients:** citric acid, glycerine, menthol, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate and sucralose

**Questions or Comments? 1-866-595-5598**

**Drug Facts**

**Active ingredients (in each 5 mL tsp.)** **Purpose**  
 Dexbrompheniramine Maleate, USP ..... 2 mg ..... Antihistamine  
 Pseudoephedrine HCl, USP ..... 60 mg ..... Nasal Decongestant

**Uses** ■ Temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies ■ helps decongest sinus openings and sinus passages ■ reduces swelling of nasal passages, shrinks swollen membranes, and temporarily restores freer breathing through the nose ■ temporarily alleviates the following symptoms due to hay fever (allergic rhinitis) ■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy and watery eyes

**Warnings**

■ **Do not use** if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if you are taking a prescription drug that contains an MAOI, ask your doctor or pharmacist before taking this product.

**Ask a doctor before use if you are** taking sedatives or tranquilizers.

**Ask a doctor before use if you have** ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ glaucoma ■ difficulty in urination due to the enlargement of the prostate gland.

**When using this product** ■ do not exceed the recommended dosage ■ excitability may occur, especially in children ■ drowsiness may occur ■ avoid alcoholic beverages ■ alcohol, sedatives and tranquilizers may increase drowsiness ■ use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if** ■ nervousness, dizziness or sleeplessness occur ■ symptoms do not improve within 7 days or occur with a fever.

NDC: 54859-802-16

**Gonex**<sup>®</sup>**COLD & ALLERGY**

Oral Solution

- Nasal & Sinus Congestion
- Runny Nose, Sneezing
- Itchy, Watery Eyes

- ANTIHISTAMINE
- NASAL DECONGESTANT

**Cherry**

Manufactured by:  
**LLORENS**  
 Llorens Pharmaceutical  
 International Division  
 Miami, FL 33147  
 www.llorensparm.com

16 FL OZ (473 mL)

**Drug Facts (continued)**

**If pregnant or breast-feeding**, ask a Health Professional before use. **Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center right away.

**Directions** Do not exceed more than 4 doses in any 24-hour period or as directed by a doctor.

Age	Dose
Adults and children 12 years of age and over	take one teaspoonful (5 mL) every 4 to 6 hours

**Other information** ■ Store at room temperature 15-30°C (59-86°F).  
 ■ Tamper-Evident: Do not use if foil seal over bottle opening is torn, broken, or missing. ■ Pharmacist - Preserve and dispense in tight, light-resistant container with a child resistant cap as defined in the USP.

**Inactive Ingredients:** Citric Acid, Flavor, Glycerin, Menthol, Methylparaben, Propylene Glycol, Propylparaben, Purified Water, Sodium Citrate and Sucralose.

**Questions or Comments?** 1-866-595-5598

THIS IS A BULK CONTAINER NOT INTENDED FOR RETAIL.

CODE# L-76 REV.:07/22



Lot. #

Exp. Date:

**CONEX**

dexbrompheniramine maleate, pseudoephedrine liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:54859-802
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>DEXBROMPHENIRAMINE MALEATE</b> (UNII: BPA9UT29BS) (DEXBROMPHENIRAMINE - UNII: 75T64B71RP)	DEXBROMPHENIRAMINE MALEATE	1 mg in 5 mL
<b>PSEUDOEPHEDRINE HYDROCHLORIDE</b> (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII: 7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg in 5 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MENTHOL</b> (UNII: L7T10EIP3A)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	

## Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54859-802-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2024	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/01/2007	

**Labeler** - Llorens Pharmaceutical International Division (037342305)

Revised: 12/2024

Llorens Pharmaceutical International Division