

FORMA-RAY - formaldehyde liquid
Gordon Laboratories

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

FORMA-RAY

A tinted aqueous perfumed base provides preferable vehicles for the topical application of formalin solution to relieve severe conditions of bromidrosis and hyperhidrosis. This formula is double the strength of Formadon. Formalin solution dries the skin's surface, and makes a very good drying agent after surgery or laser treatment.

2 oz sponge tip plastic bottle and 4 oz plastic bottle.

Formaldehyde (20% of USP strength) formulated in a aqueous perfumed base.

Indications: Antiperspirant for the treatment of hyperhidrosis and bromidrosis. Drying agent for verrucae. Drying agent for post surgical removal of warts or for nonsurgical laser treatment of warts. Directions: Apply to feet twice weekly or as prescribed by a Physician.

Contraindications/Warnings: Avoid frequent use. Avoid contact with eyes or mucous membranes. Do not apply to open wounds. Should signs of irritation develop, medications should be discontinued. Irritates eyes, nose and throat. Keep out of reach of children. For external use only. Harmful if swallowed. Contact your local poison control center immediately. Do not induce vomiting. If conscious, give eight ounce of (240ml) of milk, or water with activated charcoal.

Keep well closed in a cool place.

The image shows a red product label for 'FORMA-RAY 20% FORMALIN SOLUTION' by Gordon Laboratories. The label features a white wavy border at the top. The text is arranged in columns and includes a caution section on the left, the product name and strength in the center, and a directions/ingredients section on the right. At the bottom, there is a barcode, lot and expiration information, and the net contents.

NDC 10481-3015-2

FORMA-RAY
20% FORMALIN
SOLUTION

CAUTION: Irritates eyes, nose and throat. Avoid breathing vapor, use with adequate ventilation. In case of contact, flush with plenty of water. For eyes, get medical attention.

FOR EXTERNAL USE ONLY: Harmful if swallowed. Contact a local Poison Control Center immediately. **DO NOT INDUCE VOMITING.** If conscious, give eight ounces (240 ml) of milk, water or water with activated charcoal.

CAUTION: Should sign of irritation develop, medication should be discontinued.

CAUTION: Federal law prohibits dispensing without a prescription.

KEEP OUT OF REACH OF CHILDREN

POISON: Keep well closed in a cool place.

INDICATIONS: Antiperspirant for treatment of hyperhidrosis and bromidrosis. Drying agent for verrucae. Drying agent for postsurgical removal of warts or for nonsurgical laser treatment of warts.

INGREDIENTS: Formaldehyde (20% of USP concentrate) in an aqueous perfumed base.

DIRECTIONS: Apply to feet twice weekly as directed by your Physician. Avoid frequent use. Discontinue if irritation occurs.

Lot# Exp.

Gordon Laboratories

NET CONTENTS: 4 FL. OZ.
Gordon Laboratories • Upper Darby, PA 19082 USA

8 50217 00111 2

formaldehyde liquid

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:1048 1-30 15
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FORMALDEHYDE (UNII: 1HG84L3525) (FORMALDEHYDE - UNII:1HG84L3525)	FORMALDEHYDE	.2 mL in 59 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10481-3015-2	118 mL in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		07/15/2010	

Labeler - Gordon Laboratories (002333847)

Registrant - Gordon Laboratories (002333847)

Establishment

Name	Address	ID/FEI	Business Operations
Gordon Laboratories		002333847	manufacture

Revised: 10/2010

Gordon Laboratories