

BC CHERRY- aspirin and caffeine powder
Medtech Products Inc.

BC Cherry Aspirin

Drug Facts

Active ingredient

(in each powder)

Aspirin (NSAID*) 845 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Active ingredient

(in each powder)

Caffeine 65 mg

Purpose

Pain reliever aid

Uses

- temporary relieves minor aches and pains due to:
 - headache
 - backache
 - minor arthritis pain
 - a cold
 - muscle aches
 - toothache
 - menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- facial swelling

- asthma (wheezing)
- shock
- hives

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for longer time than directed

Caffeine warning: This product contains caffeine. Limit the use of caffeine-containing medications, foods, or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat.

Do not use

if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- stomach bleeding warnings applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- diabetes
- gout
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- any new symptoms appear
- ringing in the ears or a loss of hearing occurs.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **Adults and children 12 years of age and over:** open 1 pack, place powder on tongue every 6 hours, while symptoms persist. Drink a full glass of water with each dose, or stir powder into a glass of water or other liquid.
- Do not take more than 4 powders in 24 hours unless directed by a doctor.
- **Children under 12 years of age:** ask a doctor.

Other information

- store below 25°C (77°F)

Inactive ingredients

acesulfame potassium, colloidal silicon dioxide, flavor, guar gum, mannitol, microcrystalline cellulose, sucralose

Questions?

1-866-255-5197

PRINCIPAL DISPLAY PANEL

BC® cherry
FAST PAIN RELIEF
ASPIRIN (NSAID) – PAIN RELIEVER ● FEVER REDUCER
CAFFEINE – PAIN RELIEVER AID
24 POWDERS



PRINCIPAL DISPLAY PANEL

BC® cherry
 FAST PAIN RELIEF
 ASPIRIN (NSAID) - PAIN RELIEVER ● FEVER REDUCER
 CAFFEINE - PAIN RELIEVER AID
 4 POWDERS



Drug Facts (continued) 5	Drug Facts (continued) 6
<p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children.</p> <p>In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.</p>	<p>with each dose, or stir powder into a glass of water or other liquid. Do not take more than 4 packs in 24 hours unless directed by a doctor. Children under 12 years: ask a doctor.</p> <p>Other information</p> <ul style="list-style-type: none"> Store below 25°C (77°F) <p>Inactive ingredients</p> <p>acetamide potassium, colloidal silicon dioxide, flavor, guar gum, mannitol, microcrystalline cellulose, sucralose.</p>
<p>Directions</p> <ul style="list-style-type: none"> adults and children 12 years and over: open 1 pack, place powder on tongue every 6 hours, while symptoms persist. Drink a full glass of water. 	<p>Questions?</p> <p>1-866-255-5197 BCpowder.com</p>

BC CHERRY

aspirin and caffeine powder

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63029-200
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	845 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
GUAR GUM (UNII: E89I1637KE)	
MANNITOL (UNII: 3OWL53L36A)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63029-200-22	24 in 1 BOX; Type 0: Not a Combination Product	10/01/2012	
2	NDC:63029-200-04	4 in 1 BOX; Type 0: Not a Combination Product	10/01/2012	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	10/01/2012	

Labeler - Medtech Products Inc. (122715688)

Revised: 10/2025

Medtech Products Inc.