

**NAPROXEN SODIUM- naproxen sodium capsule, liquid filled
TOP CARE (Topco Associates LLC)**

Drug Facts

Active ingredient (in each capsule)

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID) *

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing

- it feels like the capsule is stuck in your throat

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause delivery problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

adults and children 12 years and older:	<ul style="list-style-type: none"> • take 1 capsule every 8 to 12 hours while symptoms last • for the first dose you may take 2 capsules within the first hour • do not exceed 2 capsules in any 8- to 12-hour period • do not exceed 3 capsules in a 24-hour period
children under 12 years:	<ul style="list-style-type: none"> • ask a doctor

Other information

- **each capsule contains:** sodium 20 mg
- store at 20-25°C (68-77°F) avoid high humidity and excessive heat above 40°C (104°F)
- protect from light
- read all directions and warnings before use. Keep carton.

Inactive ingredients

FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?

1-888-423-0139

Do not use if printed safety seal under cap is torn or missing

DISTRIBUTED BY TOPCO ASSOCIATES LLC

ELK GROVE VILLAGE, IL 60007

©TOPCO BMAA0521

QUESTIONS? 1-888-423-0139

topcare@topco.com www.topcarebrand.com

✓ QUALITY GUARANTEED

***This product is not manufactured

or distributed by the owners of

ALEVE® Liquid Gels.

Scan here for more

information or call

1-888-423-0139

20's Bottle Carton

+TopCare ®

health

NDC 36800-204-11

COMPARE TO ALEVE® LIQUID GELS ACTIVE INGREDIENT***

Naproxen

Sodium

Capsules, 220 mg

PAIN RELIEVER • FEVER REDUCER

(NSAID)

12 HOUR

Strength to last 12 hours

actual size

20

LIQUID-FILLED CAPSULES



NAPROXEN SODIUM

naproxen sodium capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-204
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
LACTIC ACID (UNII: 33X04XA5AT)	
MANNITOL (UNII: 3OWL53L36A)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	blue (Blue with white text)	Score	no score
Shape	CAPSULE	Size	25mm
Flavor		Imprint Code	NP1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-204-11	1 in 1 CARTON	07/02/2021	
1		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:36800-204-16	1 in 1 CARTON	07/02/2021	
2		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:36800-204-78	1 in 1 CARTON	07/02/2021	
3		80 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA021920	07/02/2021	

Labeler - TOP CARE (Topco Associates LLC) (006935977)

Registrant - Bionpharma Inc. (079637826)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

Patheon Softgels Inc.	002193829	manufacture(36800-204)
-----------------------	-----------	------------------------

Revised: 12/2022

TOP CARE (Topco Associates LLC)