

**SHISEIDO ULTIMATE SUN PROTECTOR- octisalate, titanium dioxide, octocrylene, and zinc oxide cream**  
**SHISEIDO AMERICAS CORPORATION**

-----  
**SHISEIDO ULTIMATE SUN PROTECTOR CREAM**

***Drug Facts***

<b><i>Active ingredients</i></b>	<b><i>Purpose</i></b>
Octisalate 5.0%	Sunscreen
Octocrylene 7.0%	Sunscreen
<i>Titanium Dioxide</i> 3.8%	Sunscreen
Zinc Oxide 19.4%	Sunscreen

**Uses**

- helps prevent sunburn
- if used as directed with other sun protection measures (see ***Directions***), decreases the risk of skin cancer and early skin aging caused by the sun

**Warnings**

**For external use only**

**Do not use on damaged or broken skin**

**When using this product** keep out of eyes. Rinse with water to remove.

**Stop use and ask a doctor** if rash occurs.

**Keep out of reach of children.** If product is swallowed, get medical help or contact a Poison Control Center right away.

**Flammable**

Do not use near heat, flame, or while smoking.

**Directions**

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- reapply:
  - after 80 minutes of swimming or sweating
  - immediately after towel drying
  - at least every two hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. – 2 p.m.

- wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor

### **Inactive Ingredients**

WATER ■ DIMETHICONE ■ SD ALCOHOL 40-B ■ ISOPROPYL MYRISTATE ■ PEG/PPG-9/2 DIMETHYL ETHER ■ METHYL METHACRYLATE CROSSPOLYMER ■ LAURYL PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE ■ GLYCERIN ■ ISOHEXADECANE ■ POLYBUTYLENE GLYCOL/PPG-9/1 COPOLYMER ■ ISODODECANE ■ TRIMETHYLSILOXYSILICATE ■ DISTEARDIMONIUM HECTORITE ■ DEXTRIN PALMITATE ■ PEG/PPG-14/7 DIMETHYL ETHER ■ SAXIFRAGA SARMENTOSA EXTRACT ■ SCUTELLARIA BAICALENSIS ROOT EXTRACT ■ CAMELLIA SINENSIS LEAF EXTRACT ■ ONONIS SPINOSA ROOT EXTRACT ■ SODIUM HYALURONATE ■ SOPHORA ANGUSTIFOLIA ROOT EXTRACT ■ ALUMINUM HYDROXIDE ■ SILICA ■ HYDROGEN DIMETHICONE ■ STEARIC ACID ■ SORBITAN SESQUIISOSTEARATE ■ CARBOXYDECYL TRISILOXANE ■ TRISODIUM EDTA ■ BUTYLENE GLYCOL ■ ALCOHOL ■ POLYMETHYLSILSESQUIOXANE ■ TALC ■ TOCOPHEROL ■ BHT ■ SODIUM METABISULFITE ■ PEG-6 ■ SYZYGIUM JAMBOS LEAF EXTRACT ■ PHENOXYETHANOL ■ FRAGRANCE ■

### **Other information**

Protect this product in this container from excessive heat and direct sun.

### **Questions or comments?**

Call toll free 1-800-906-7503

### **PRINCIPAL DISPLAY PANEL - 50 mL Tube Label**

SHISEIDO  
GINZA TOKYO

Ultimate  
Sun Protector Cream

50+

SynchroShield  
WetForce x HeatForce

BROAD SPECTRUM  
SPF 50+

For Face

WATER RESISTANT  
(80 MINUTES)

SUNSCREEN

50mL NET WT. 2 OZ.



SHISEIDO

GINZA TOKYO

Ultimate  
Sun Protector Cream

50+



SynchroShield

WetForce x HeatForce

BROAD SPECTRUM

SPF 50+

For Face

WATER RESISTANT

(80 MINUTES)

SUNSCREEN

50mL NET WT. 2 OZ.



SHISEIDO  
ULTIMATE SUN PROTECTOR CREAM  
BROAD SPECTRUM SPF 50+

Powerful sun protection with SynchroShield™. The protective veil becomes even stronger whenever it senses heat, water or perspiration\*. Enriched with skincare ingredients to help prevent the appearance of fine lines, wrinkles and uneven skin tone.

\*The HeatForce technology is activated on hot days, after the formula is applied to the skin. To activate the WetForce technology, expose the applied area to water or perspiration for 30 minutes.

**Drug Facts**

**Active ingredients**

	<i>Purpose</i>
OCTISALATE 5.0% .....	Sunscreen
OCTOCRYLENE 7.0% .....	Sunscreen
TITANIUM DIOXIDE 3.8% .....	Sunscreen
ZINC OXIDE 19.4% .....	Sunscreen

**Uses** ■ helps prevent sunburn ■ if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

**Warnings**

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable : Do not use near heat, flame, or while smoking.

**Directions** ■ apply liberally 15 minutes before sun exposure ■ reapply:  
· after 80 minutes of swimming or sweating · immediately after towel  
drying · at least every 2 hours ■ **Sun Protection Measures.** Spending time  
in the sun increases your risk of skin cancer and early skin aging. To  
decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF of  
15 or higher and other sun protection measures including: · limit time in  
the sun, especially from 10 a.m. – 2 p.m. · wear long-sleeve shirts, pants,  
hats, and sunglasses ■ children under 6 months: Ask a doctor

**Inactive ingredients** WATER · DIMETHICONE · SD ALCOHOL 40-B · ISOPROPYL MYRISTATE ·  
PEG/PPG-9/2 DIMETHYL ETHER · METHYL METHACRYLATE CROSSPOLYMER · LAURYL PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE · GLYCERIN ·  
ISOHEXADECANE · POLYBUTYLENE GLYCOL/PPG-9/1 COPOLYMER · ISODODECANE · TRIMETHYLSILOXYSILICATE · DISTEARDIMONIUM HECTORITE ·  
DEXTRIN PALMITATE · PEG/PPG-14/7 DIMETHYL ETHER · SAXIFRAGA SARMENTOSA EXTRACT · SCUTELLARIA BAICALENSIS ROOT EXTRACT ·  
CAMELLIA SINENSIS LEAF EXTRACT · ONONIS SPINOSA ROOT EXTRACT · SODIUM HYALURONATE · SOPHORA ANGUSTIFOLIA ROOT EXTRACT ·  
ALUMINUM HYDROXIDE · SILICA · HYDROGEN DIMETHICONE · STEARIC ACID · SORBITAN SESQUISOSEARATE · CARBOXYDECYL TRISILOXANE ·  
TRISODIUM EDTA · BUTYLENE GLYCOL · ALCOHOL · POLYMETHYLSILSESQUOXANE · TALC · TOCOPHEROL · BHT · SODIUM METABISULFITE · PEG-6 ·  
SYZYGIUM JAMBOS LEAF EXTRACT · PHENOXYETHANOL · FRAGRANCE ·

### Other Information

■ protect this product in this container from excessive heat and direct sun.

**Questions or Comments?** Call toll free 1-800-906-7503

Reducing the quantity of application will lower the level of sunscreen protection significantly.



Blue Conscious Project

Specially formulated by  
Shiseido Laboratories, Japan.

SHISEIDO AMERICAS CORPORATION DIST.  
NEW YORK, NY 10017  
MADE IN U.S.A.

GLO. 15676

[www.shiseido.com](http://www.shiseido.com)

<http://s1872.com/15676AA>

See our website for more  
information.



7 30852 15676 0

## SHISEIDO ULTIMATE SUN PROTECTOR

octisalate, titanium dioxide, octocrylene, and zinc oxide cream

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:58411-480
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>OCTISALATE</b> (UNII: 4X49Y0596W) (OCTISALATE - UNII:4X49Y0596W)	OCTISALATE	3.105 g in 50 mL
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.359 g in 50 mL
<b>OCTOCRYLENE</b> (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	4.347 g in 50 mL
<b>ZINC OXIDE</b> (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	12.047 g in 50 mL

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>WATER</b> (UNII: 059QF0K00R)	
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>ISOPROPYL MYRISTATE</b> (UNII: 0RE8K4LNJS)	
<b>PEG/PPG-9/2 DIMETHYL ETHER</b> (UNII: LM6Q66468X)	
<b>METHYL METHACRYLATE/GLYCOL DIMETHACRYLATE CROSSPOLYMER</b> (UNII: EG97988M5Q)	
<b>LAURYL PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE</b> (UNII: 25G622K2RA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>ISOHEXADECANE</b> (UNII: 918X1OUF1E)	
<b>ISODODECANE</b> (UNII: A8289P68Y2)	
<b>TRIMETHYLSILOXYSILICATE (M/Q 0.6-0.8)</b> (UNII: 5041RX63GN)	
<b>DISTEARDIMONIUM HECTORITE</b> (UNII: X687XDK09L)	
<b>DEXTRIN PALMITATE (CORN; 20000 MW)</b> (UNII: 89B2BSF9I3)	
<b>PEG/PPG-14/7 DIMETHYL ETHER</b> (UNII: 6DNW9T7YT2)	
<b>SAXIFRAGA STOLONIFERA LEAF</b> (UNII: O3TMV4903H)	
<b>SCUTELLARIA BAICALENSIS ROOT</b> (UNII: 7J95K7ID2S)	
<b>GREEN TEA LEAF</b> (UNII: W2ZU1RY8B0)	
<b>ONONIS SPINOSA ROOT</b> (UNII: FD2FMC53M1)	
<b>HYALURONATE SODIUM</b> (UNII: YSE9PPT4TH)	
<b>SOPHORA FLAVESCENS ROOT</b> (UNII: IYR6K8KQ5K)	
<b>ALUMINUM HYDROXIDE</b> (UNII: 5QB0T2IUN0)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>HYDROGEN DIMETHICONE (13 CST)</b> (UNII: 4QGR4P2YOI)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>SORBITAN SESQUIISOSTEARATE</b> (UNII: VU97D01BF9)	
<b>EDETATE TRISODIUM</b> (UNII: 420IP921MB)	
<b>BUTYLENE GLYCOL</b> (UNII: 3XUS85KORA)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>POLYMETHYLSILSESQUIOXANE (4.5 MICRONS)</b> (UNII: 59Z907ZB69)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>.ALPHA.-TOCOPHEROL</b> (UNII: H4N855PNZ1)	
<b>BUTYLATED HYDROXYTOLUENE</b> (UNII: 1P9D0Z171K)	
<b>SODIUM METABISULFITE</b> (UNII: 4VON5FNS3C)	

<b>POLYETHYLENE GLYCOL 300</b> (UNII: 5655G9Y8AQ)	
<b>SYZYGIUM JAMBOS LEAF</b> (UNII: 407Z4W5LFF)	
<b>PHENOXYETHANOL</b> (UNII: HIE492ZZ3T)	
<b>SODIUM POLYMETAPHOSPHATE</b> (UNII: P1BM4ZH95L)	
<b>.ALPHA.-CELLULOSE</b> (UNII: I355QGZ19A)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>FERROSOFERRIC OXIDE</b> (UNII: XM0M87F357)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-480-60	1 in 1 CARTON	01/01/2020	
1		50 mL in 1 TUBE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	01/01/2020	

**Labeler** - SHISEIDO AMERICAS CORPORATION (193691821)

### Establishment

Name	Address	ID/FEI	Business Operations
Shiseido America Inc.		782677132	analysis(58411-480) , manufacture(58411-480)

Revised: 1/2026

SHISEIDO AMERICAS CORPORATION