

**BOVATEC 150FP- lasalocid sodium powder**  
**Phibro Animal Health**

-----  
**Bovatec® 150 FP**

**Bovatec® 150 FP**

**(lasalocid sodium  
Type A medicated article)**

**For further manufacturing only**

**Active drug ingredient**

Each pound contains 150 grams (33.1%) of lasalocid (as lasalocid sodium activity) in a carrier suitable for incorporation in feed.

**Mixing directions**

This product should be further diluted in an intermediate blending step prior to mixing in the final feed.

**1. Complete Type C Medicated Feed**

- a. Intermediate blending: Mix 1 part of Bovatec 150 FP (Type A Medicated Article) with 29 parts of finely ground non-medicated feedstuffs to provide an intermediate preblend containing 5 grams of lasalocid sodium per pound.
- b. Using the 5 g/lb intermediate preblend, the following table would apply to the manufacture of complete Type C medicated feeds:

<b>Intended lasalocid sodium concentration in the complete Type C medicated feed</b>		<b>Add the following amounts of 5 grams per pound intermediate preblend per ton of complete Type C medicated feed mixed:</b>
<b>grams/ton</b>	<b>mg/lb</b>	
10	5	2
15	7.5	3
30*	15	6

\* Maximum approved concentration for complete Type C medicated feeds for growing beef steers and heifers fed in confinement for slaughter and sheep maintained in confinement.

- c. Using the 5 g/lb intermediate preblend, the following table would apply to the manufacture of other complete Type C medicated feeds and supplements:

<b>Desired daily</b>	<b>Add the following amounts</b>	<b>Lasalocid sodium will be present in the Type C</b>
----------------------	----------------------------------	---

Daily feeding rate	Desired daily lasalocid sodium intake	of 5 grams per pound intermediate preblend per ton of Type C medicated feed or supplement mixed:	present in the Type C medicated feed or supplement at the following concentration:
1	60	24	120
	200	80	400
	300	120	600
2	60	12	60
	200	40	200
	300	60	300
3	60	8	40
	200	26.7	133.5
	300	40	200

## 2. Dry Type C Medicated Feed Supplement

- a. Intermediate blending: Mix 1 part Bovatec 150 FP Type A Medicated Article with 6.5 parts of finely ground non-medicated feedstuffs to provide an intermediate preblend containing 20 grams of lasalocid per pound. For example, mix one 50 lb bag of Bovatec 150 FP Type A Medicated Article with 325 lbs of non-medicated feedstuffs.
- b. Using the 20 g/lb intermediate preblend, the following table would apply to the manufacture of dry Type C medicated supplements:

Intended lasalocid sodium concentration in the Type C medicated feed supplement	Add the following amounts of 20 grams per pound intermediate preblend per ton of Type C medicated feed supplement mixed:	
		grams/ton
400	200	20
600	300	30
1000	500	50
1440	720	72
1600	800	80

- c. Thoroughly mix Type C medicated feed supplement with sufficient non-medicated feedstuffs in accordance with the feeding rates listed in the use directions below.

## 3. Type C Liquid Medicated Feed Supplement Intended for Addition to Dry Feeds

- a. Type C liquid medicated feed supplements with suspending agent(s) should be in a pH range of 4-8 and maintain physical stability for up to three months with a viscosity not less than 300 cps.
- b. Conventional liquid supplements should be in a pH range of 4-8.

For liquid feeds stored in recirculating tank systems: Recirculate immediately prior to use for no less than 10 minutes, moving not less than 1 percent of the tank contents per minute from the bottom of the tank to the top. Recirculate daily as described even when not used.

For liquid feeds stored in mechanical, air, or other agitation-type tank systems: Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily as described even when not used.

## Feeding directions

Species	Dose	Indications for use
<b>Growing beef steers and heifers fed in confinement for slaughter</b>	10-30 grams lasalocid sodium per ton of total ration (90% dry matter)	For improved feed efficiency: Feed as the sole ration to provide 100 to 360 mg lasalocid sodium per head per day.
	25-30 grams lasalocid sodium per ton of total ration (90% dry matter)	For increased rate of weight gain and improved feed efficiency: Feed as the sole ration to provide 250 to 360 mg lasalocid sodium per head per day.
<b>Growing beef steers and heifers on pasture (stocker, feeder, and slaughter) and replacement beef and dairy heifers</b>	60-300 mg lasalocid sodium per head/day	For increased rate of weight gain: Feed in at least 1 pound of Type C medicated feed. Intakes of lasalocid sodium greater than 200 mg per head per day have not been shown to be more effective than 200 mg per head per day.
	60-300 mg lasalocid sodium per head/day	For increased rate of weight gain: Feed on a free-choice basis. Intakes of lasalocid sodium greater than 200 mg per head per day have not been shown to be more effective than 200 mg per head per day. Type C free-choice medicated feed must be manufactured under a medicated feed mill license utilizing an FDA approved formulation.  NOTE: Coccidiosis may occur when young pasture cattle are co-mingled with adult cattle passing coccidial oocysts.
<b>Cattle up to 800 lbs</b>	1 mg lasalocid sodium	For control of coccidiosis caused by <i>Eimeria bovis</i> and <i>E. zuernii</i> . Hand feed to provide up to 360 mg lasalocid sodium per head per day.

	per 2.2 lbs body weight/day	
<b>Beef calves less than 2 months of age and dairy calves</b>	1 mg lasalocid sodium per 2.2 lbs body weight/day	For control of coccidiosis caused by <i>Eimeria bovis</i> and <i>E. zuernii</i> . Thoroughly mix milk replacer powder with warm water and hand feed immediately.
<b>Sheep maintained in confinement</b>	20-30 grams lasalocid sodium per ton of total ration (90% dry matter)	For prevention of coccidiosis caused by <i>Eimeria ovina</i> , <i>E. crandallis</i> , <i>E. ovinoidalis</i> ( <i>E. ninakohlyakimovae</i> ), <i>E. parva</i> , and <i>E. intricata</i> : Feed as the sole ration to provide 15 to 70 mg lasalocid sodium per head per day depending on the body weight.

## WITHDRAWAL PERIODS AND RESIDUE WARNINGS

**No withdrawal period is required when used according to labeling. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.**

### User Safety Warnings

**Not for use in humans. Keep out of reach of children. When mixing and handling lasalocid sodium Type A medicated articles, use protective clothing, impervious gloves and a dust mask. Avoid contact with eyes. Operators should wash hands thoroughly with soap and water after handling.**

### Animal Safety Warnings

**Feeding undiluted or mixing errors may result in an excess lasalocid concentration which could be fatal to cattle and sheep. Do not allow horses or other equines access to Type A articles or Type B feeds containing lasalocid as ingestion may be fatal. Safety of lasalocid for use in unapproved species has not been established.**

**Questions/Comments?** For technical assistance or to obtain a Safety Data Sheet(s), contact Phibro Animal Health Corporation at 888-403-0074 or [www.pahc.com](http://www.pahc.com). To report side

effects, contact Phibro Animal Health Corporation at 1-888-475-7355. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

Phibro, Phibro logo and Bovatec are trademarks owned by or licensed to Phibro Animal Health Corporation or its affiliates.

***Distributed by:***

**Phibro Animal Health Corporation**

**Teaneck, NJ 07666**

**Net Wt 50 LB (22.68 kg)**

**176-9035-04**

**TAKE TIME**



**OBSERVE LABEL  
DIRECTIONS**

**Take Time**

**Storage Conditions**

**Store below 25°C (77°F), excursions permitted to 37°C (99°F).  
Restricted drug (California) - Use only as directed. Not for human use.  
Approved by FDA under NADA # 096-298**

**PRINCIPAL DISPLAY PANEL**



## Product Characteristics

<b>Color</b>	white (white to tan)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66104-1457-1	22.68 kg in 1 BAG		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA096298	01/01/2009	

**Labeler** - Phibro Animal Health (006989008)

Revised: 4/2026

Phibro Animal Health