

# **DYE FREE PAIN AND FEVER CHILDRENS- acetaminophen suspension**

## **Walgreens**

-----

### **Drug Facts**

#### **Active ingredient (in each 5 mL)**

**Acetaminophen 160 mg**

#### **Purpose**

**Pain reliever/fever reducer**

#### **Uses**

- temporarily reduces fever
- temporarily relieves minor aches and pains due to:
  - the common cold
  - flu
  - headache
  - sore throat
  - toothache

#### **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### **Do not use**

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this

product

**Ask a doctor before use if your child has**

liver disease.

**Ask a doctor or pharmacist before use if your child is**

taking the blood thinning drug warfarin.

**When using this product,**

**do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- **this product does not contain directions or complete warnings for adult use**
- **do not take more than directed (see overdose warning)**
- **shake well before using**
- mL = milliliter
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by a doctor
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

<b>Weight (lb)</b>	<b>Age (year)</b>	<b>Dose (mL)*</b>
under 24	under 2	ask a doctor
24-35	2-3	5 mL
36-47	4-5	7.5 mL
48-59	6-8	10 mL
60-71	9-10	12.5 mL

72-95	11	15 mL
-------	----	-------

\*or as directed by a doctor

### **Other information**

- **each 5 mL contains:**sodium 2 mg
- store between 20-25°C (68-77°F)
- do not refrigerate
- see bottom panel for lot number and expiration date

### **Inactive ingredients**

anhydrous citric acid, butylparaben, carboxymethylcellulose sodium, flavor, glycerin, high fructose corn syrup, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

### **Questions or comments?**

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

### **Principal Display Panel**

Compare to the active ingredient in Children's Tylenol® Oral Suspension††

Children's

#### **Dye-Free**

#### **Pain & Fever**

ACETAMINOPHEN 160 mg PER 5 mL

PAIN RELIEVER /FEVER REDUCER

ORAL SUSPENSION

Aspirin Free

Alcohol Free

Ibuprofen Free

AGES 2 - 11 YEARS

FL OZ (mL)

GRAPE FLAVOR

DOSAGE CUP PROVIDED

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

††This product is not manufactured or distributed by McNeil Consumer Healthcare,

distributor of Children's Tylenol® Oral Suspension.

DISTRIBUTED BY: WALGREEN CO.

DEERFIELD, IL 60015

walgreens.com

**Product Label**

**Drug Facts (continued)**

- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 doses in 24 hours
- do not give more than 5 days unless directed by a doctor
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

Weight (lb)	Age (year)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-86	11 years	15 mL

\*or as directed by a doctor

**Other information**

- each 5 mL contains sodium 2 mg
- store between 20-25°C (68-77°F) do not refrigerate
- see bottom panel for lot number and expiration date

**Inactive ingredients** butylparaben, carbonylmethylcellulose sodium, citric acid, flavors, glycerin, high fructose corn syrup, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

**Questions or comments?**  
Call 1-877-753-3838 Monday-Friday 9AM-5PM EST

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING**

**DO dosage cup provided**

DISTRIBUTED BY: WALGREEN CO.  
DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com ©2021 Walgreen Co.

NDC 0363-0285-04

Compare to the active ingredient in Children's Tylenol® Oral Suspension\*

# Children's Dye-Free Pain & Fever

**ACETAMINOPHEN 160 mg PER 5 mL PAIN RELIEVER / FEVER REDUCER ORAL SUSPENSION**

Aspirin Free  
Alcohol Free  
Ibuprofen Free

AGES 2-11 YEARS

4 FL OZ (118 mL) Grape flavor

NDC 0363-0285-04

Compare to the active ingredient in Children's Tylenol® Oral Suspension\*

# Children's Dye-Free Pain & Fever

**ACETAMINOPHEN 160 mg PER 5 mL PAIN RELIEVER / FEVER REDUCER ORAL SUSPENSION**

Aspirin Free  
Alcohol Free  
Ibuprofen Free

AGES 2-11 YEARS

4 FL OZ (118 mL) Grape flavor

**Drug Facts**

**Active ingredient (in each 5 mL)**  
Acetaminophen 160 mg. Pain reliever/fever reducer

**Uses**

- temporarily reduces fever
- temporarily relieves minor aches and pains due to:
  - the common cold
  - flu
  - headache
  - sore throat
  - toothache

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away. **Severe throat warning:** If sore throat is severe, persists for more than 3 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if your child has liver disease.**

**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dose (see Overdose warning).**

**Stop use and ask a doctor if:**

- redness or swelling is present
- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-232-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not take more than directed (see Overdose warning)
- shake well before using
- mL = milliliter

\*Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.  
\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Oral Suspension.

W3ORG0222-F REV-0624

ITEM 461454 W00000-0000-0

PLD 1254D FZ009149 PRODUCT OF CHINA

Lot No:  
Exp. Date:

**WALGREENS Children's Dye-Free Pain and Fever Grape Flavor**

**DYE FREE PAIN AND FEVER CHILDRENS**

acetaminophen suspension

**Product Information**

**Product Type**

HUMAN OTC DRUG

**Item Code (Source)**

NDC:0363-0285

<b>Route of Administration</b>	ORAL
--------------------------------	------

<b>Active Ingredient/Active Moiety</b>		
<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

<b>Inactive Ingredients</b>	
<b>Ingredient Name</b>	<b>Strength</b>
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>BUTYLPARABEN</b> (UNII: 3QPI1U3FV8)	
<b>CARBOXYMETHYLCELLULOSE SODIUM</b> (UNII: K679OBS311)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>HIGH FRUCTOSE CORN SYRUP</b> (UNII: XY6UN3QB6S)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

<b>Product Characteristics</b>			
<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	GRAPE	<b>Imprint Code</b>	
<b>Contains</b>			

<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:0363-0285-04	1 in 1 BOX	03/31/2015	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

<b>Marketing Information</b>			
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M013	03/31/2015	

**Labeler** - Walgreens (008965063)

