

**EASY CARE FIRST AID KIT - SPORTS AND TRAVEL - benzalkonium chloride, benzocaine, sd alcohol, bacitracin zinc, neomycin sulfate, polymyxin b, ibuprofen, acetaminophen  
Tender Corp dba Adventure Medical Kits**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Easy Care First Aid Kit Sports and Travel**

**Active Ingredient**

Active Ingredient:

Benzalkonium Chloride 0.40%

**Purpose**

Antiseptic

**Use**

For Professional and Hospital use. Helps prevent infection. Antiseptic cleansing of face, hands and body without soap and water.

**Warnings**

Warning: For external use only.

Keep out of reach of children: If swallowed, get medical help or contact a Poison Control Center right away.

Stop use if unusual redness, swelling or other symptoms occur. Consult a physician immediately.

Do not use in the eyes or over large areas of the body.

**Directions**

Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.

**Inactive Ingredients**

Inactive Ingredient: Purified water

LOT/EXP:                      Made in CHINA

20130301

Antiseptic Towelette

Genuine First Aid LLC, Clearwater FL 33755

[www.GenuineFirstAid.com](http://www.GenuineFirstAid.com)

1/pouch

GENUINE FIRST AID

Active Ingredient:                      Purpose:  
Benzocaine, 6% w/v.....      Topical Anesthetic  
SD alcohol, 60% w/v.....      Antiseptic

Use: For the temporary relief of pain and itching associated with minor burns, scrapes and insect bites.

Warnings: For external use only.

Avoid contact with eyes. If this happens, rinse thoroughly with water.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Flammable - keep away from fire or flame.

Do not use: In eyes, on broken skin, deep puncture wounds. If unusual redness, swelling, irritation or other symptoms occur, consult a physician immediately.

Made in CHINA

LOT/EXP:

Insect Sting Relief Pad

Genuine First Aid LLC, Clearwater FL 33755  
[www.GenuineFirstAid.com](http://www.GenuineFirstAid.com)

1/pouch

GENUINE FIRST AID

**Active Ingredients**

Active Ingredient: .....Bacitracin Zinc 400 units  
Neomycin Sulfate 5mg ( equivalent to 3.5 mg Neomycin base)  
Polymyxin B Sulfate 5000 units

**Purpose**

Triple Antibiotic  
Uses: To help prevent infection in:  
minor cuts; scrapes; burns

**Warnings**

For external use only.  
Do not use: in eyes; over large areas of the body;  
If allergic to any of the ingredients; for more than one week unless directed by a physician.  
Stop use and consult a doctor:

if the condition persists or gets worse; a rash or other allergic reaction develops

Keep out of reach of children.

If ingested, contact a Poison

Control Center right away.

**Directions**

Directions: clean affected area; apply small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily; may be covered with a sterile bandage

Other information:

Store at room temperature.

**Inactive Ingredient**

water

Genuine Triple Antibiotic

First Aid Ointment

To Help Prevent Infection

Each Gram Contains:

Bacitracin Zinc 400 units

Neomycin Sulfate 5 mg

(equivalent to 3.5 mg

Neomycin base)

Polymyxin B Sulfate 5000 units

Net Wt. 0.5g ; (1/64 oz)

Manufactured in CHINA for

GENUINE FIRST AID.

Triple Antibiotic Ointment 10pcs

Net wt. 0.9g (1/32oz)

100

Triple Antibiotic

Active Ingredient (in each tablet)

Purpose

Acetaminophen 325 mg .....

Analgesic/antipyretic

Uses

temporary relief of minor aches and pains associated with:

common cold; headache; toothache; muscular aches; backache; arthritis; menstrual cramps; and reduction of fever

## Warnings:

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if: adult takes more than 12 tablets in 24 hours, which is the maximum daily amount; child takes more than 5 doses in 24 hours; taken with other drugs containing acetaminophen; adult has 3 or more alcoholic drinks every day while using this product

Do not use: with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist; for more than 10 days for pain unless directed by a doctor; for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if the user has liver disease

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if: symptoms do not improve; pain gets worse or lasts for more than 10 days; fever gets worse or lasts for more than 3 days; new symptoms occur; redness or swelling is present; a rare sensitivity reaction occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, contact a doctor or Poison Control Center immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Do not exceed recommended dosage

## Directions

Adults and Children 12 years of age or older	Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours.
--	--

Children 6-11 years of age	Take 1 tablet every 4 to 6 hours as needed. Do not take more than 5 tablets in 24 hours.
-------------------------------	--

Children under 6 years of age	Do not use this regular strength product. This will provide more than the recommended dose (overdose) and could cause serious health problems.
----------------------------------	---

Store at 59-86 degree F (15-30 degree C)

tamper evident sealed packets; do not use any open or torn packets

Distributed by GENUINE FIRST AID  
600 Cleveland Str Suite 400, Clearwater, FL 33755

GENUINE FIRST AID      2 Tablets

NON-ASPIRIN

Active ingredient (in each tablet)	Purpose
Ibuprofen USP (NSAID*) 200mg . . . . .	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	

Uses temporarily relieves minor aches and pains due to:

the common cold

headache

toothache

muscular aches

backache

minor pain of arthritis

menstrual cramps temporarily reduces fever

### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: shock, facial swelling, asthma (wheezing) rash, skin reddening, blisters, hives. If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinner (anticoagulant) or steroid drug, take other drugs containing NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery.

Ask a doctor before use if stomach bleeding warning applies to you; you have a history of stomach problems such as heartburn; you have a high blood pressure, heart disease, liver cirrhosis, or kidney disease; you are taking a diuretic

Ask a doctor before use if you are: taking any other drug containing NSAID (prescription or nonprescription); taking aspirin for heart attack or stroke, because Ibuprofen may decrease this benefit of aspirin; taking any other drug

When using this product:  
take with food or milk if stomach upset occurs

Stop use and ask a doctor If:  
you experience any of the following signs of stomach bleeding; feel faint; vomit blood; have bloody or

black stools; have stomach pain that does get better; pain gets worse or lasts more than 10 days; fever gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions:**

do not use more than directed; the smallest effective dose should be used; do not take longer than 10 days, unless directed by a doctor.

Adults and Children (12 years and older): Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Other information: Store at controlled room temperature; avoid excessive heat 40 degree Celsius (104 degree Fahrenheit); tamper evident sealed packets; do not use any opened or torn packets

Inactive ingredients: cellulose, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, polyethylene glycol, povidone, silica, sodium starch glycolate, stearic acid, titanium dioxide, triacetin.

Distributed by GENUINE FIRST AID

600 Cleveland Str Suite 400, Clearwater, FL 33755

IBUPROFEN 2 Tablets

IBUPROFEN 2 Tablets

Reorder TAO-001

# Genuine Triple Antibiotic

**First Aid Ointment  
To Help Prevent Infection**

**Each Gram Contains:**  
**Bacitracin Zinc 400 units**  
**Neomycin Sulfate 5 mg**  
**(equivalent to 3.5 mg**  
**Neomycin base)**  
**Polymyxin B Sulfate 5000 units**  
**Net Wt 0.5g · (1/64 oz)**



Manufactured in CHINA for  
**GENUINE FIRST AID.**

## Drug Facts

Uses: To help prevent infection  
in • minor cuts • scrapes • burns

## Warnings:

For external use only

Do not use • in eyes • over large  
areas of the body • if allergic to  
any of the ingredients • for more  
than one week unless directed  
by a physician

Stop use and consult a doctor  
• if the condition persists or gets  
worse • a rash or other allergic  
reaction develops

Keep out of reach of children.

If ingested, contact a Poison  
Control Center right away.

Directions: • clean affected area  
• apply small amount of this product  
(an amount equal to the surface  
area of the tip of a finger) on the  
area 1 to 3 times daily • may be  
covered with a sterile bandage

Other information:

Store at room temperature

LOT/EXP:

20130118



GENUINE FIRST AID. 2 Tablets

# NON-ASPIRIN

**Active ingredient (in each tablet) Purpose**  
Acetaminophen 325 mg . . . . . Analgesic/antipyretic

## Uses

temporary relief of minor aches and pains associated  
with

- common cold    ■ headache    ■ toothache
- muscular aches    ■ backache    ■ arthritis
- menstrual cramps    ■ and reduction of fever

## Warnings

**Liver warning:** This product contains acetaminophen.  
Severe liver damage may occur if ■ adult takes more than  
12 tablets in 24 hours, which is the maximum daily amount  
■ child takes more than 5 doses in 24 hours ■ taken with  
other drugs containing acetaminophen ■ adult has 3 or  
more alcoholic drinks every day while using this product

**Do not use** ■ with any other drug containing  
acetaminophen (prescription or nonprescription). If you  
are not sure whether a drug contains acetaminophen,  
ask a doctor or pharmacist ■ for more than 10 days  
for pain unless directed by a doctor ■ for more than  
3 days for fever unless directed by a doctor

**Ask a doctor before use if the user has liver disease**

**Ask a doctor or pharmacist before use if the user is  
taking the blood thinning drug warfarin** ▶

## Warnings (continued)

**Stop use and ask a doctor if** ■ symptoms do not improve ■  
pain gets worse or lasts for more than 10 days ■ fever gets  
worse or lasts for more than 3 days ■ new symptoms occur ■  
redness or swelling is present ■ a rare sensitivity reaction occurs

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.** In case of accidental overdose,  
contact a doctor or Poison Control Center immediately. Prompt  
medical attention is critical for adults as well as for children even if  
you do not notice any signs or symptoms. **Do not exceed  
recommended dosage.**

## Directions

Adults and children 12 years of age and older	Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours.
Children 6-11 years of age	Take 1 tablet every 4 to 6 hours as needed. Do not take more than 5 tablets in 24 hours.
Children under 6 years of age	Do not use this regular strength product. This will provide more than the recommended dose (overdose) and could cause serious health problems.

**Other information** ■ store at 59°-86°F (15°-30°C) ■ tamper  
evident sealed packets ■ do not use any open or torn packets

**Inactive ingredients** corn starch, hydroxypropyl  
methylcellulose, polyethylene glycol, pregelatinized starch, stearic  
acid. May contain povidone and sodium starch glycolates.

Distributed by **GENUINE FIRST AID**  
600 Cleveland Str Suite 400, Clearwater, FL 33755

REORDER AST-001

**Antiseptic  
Towelette**  
**Toallitas  
Antisepicas**



Genuine First Aid LLC, Clearwater FL 33755  
www.GenuineFirstAid.com



**GENUINE FIRST AID®**



DRUG FACTS - Antiseptic Towelette

Active Ingredient: Benzalkonium Chloride 0.40% Purpose: First Aid Antiseptic

Use: For Professional and Hospital use. Helps prevent infection. Antiseptic cleansing of face, hands and body without soap and water.

Warnings: For external use only. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. If unusual redness, swelling or other symptoms occur, consult a physician immediately.

Do not use: In the eyes, or over large areas of the body.

Directions: Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.

Inactive ingredient: Purified water

Made in CHINA

LOT/EXP:  
20130301

--- TEAR HERE ---





GENUINE FIRST AID®



ISO 9001:2010



Genuine First Aid LLC, Clearwater FL 33755  
www.GenuineFirstAid.com

**Insect Sting  
Relief Pad**

---

**Toallitas para  
Picaduras de Insectos**

REORDER ISRP-001

**DRUG FACTS - Insect Sting Relief Pad**

**Active Ingredient:** Benzocaine, 6% w/v ..... Topical Anesthetic

SD alcohol, 60% w/v ..... Antiseptic

**Use:** For the temporary relief of pain and itching associated with minor burns, scrapes and insect bites.

**Warnings:** For external use only. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. Flammable - keep away from fire or flame. Avoid contact with eyes. If this happens, rinse thoroughly with water.

**Do not use:** In eyes, on broken skin, deep puncture wounds. If unusual redness, swelling, irritation or other symptoms occur, consult a physician immediately.

--- TEAR HERE ---

Made in CHINA

# IBUPROFEN 2 Tablets

**Active ingredient (in each tablet)** Purpose  
Ibuprofen USP (NSAID)\* 200mg . . . . . Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses** temporarily relieves minor aches and pains due to:  
■ the common cold ■ headache ■ toothache ■ muscular aches  
■ backache ■ minor pain of arthritis ■ menstrual cramps  
temporarily reduces fever

**Warnings** **Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ shock ■ facial swelling ■ asthma (wheezing) ■ rash ■ skin reddening ■ blisters ■ hives If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are** ■ taking any other drug containing an NSAID (prescription or nonprescription) ▶

**Warnings (continued)**

■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

**When using this product**

■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ do not use more than directed ■ the smallest effective dose should be used ■ do not take longer than 10 days, unless directed by a doctor

**Adults and Children (12 years and older):** Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

**Children under 12 years:** Do not give to children under 12 years of age.

**Other information** ■ store at controlled room temperature ■ avoid excessive heat 40° C (104° F) ■ tamper evident sealed packets ■ do not use any opened or torn packets

**Inactive ingredients** cellulose, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, polyethylene glycol, povidone, silica, sodium starch glycolate, stearic acid, titanium dioxide, triacetin

Distributed by GENUINE FIRST AID  
600 Cleveland Str Suite 400, Clearwater, FL 33755

**EASY CARE**  
**First Aid Kit**  
SPORTS + TRAVEL

**CONTENTS:**

**Cuts & Scrapes**

Adhesive Plastic Bandage 3/4"x3"	15
Adhesive Plastic Bandage 1"x3"	5
Adhesive Knuckle Fabric Bandage	2
Adhesive Spck Bandage 7/8"x7/8" Plastic	1
Elbow/Knee Fabric Adhesive Bandage	2
Antiseptic Towelette	20
Triple Antibiotic Ointment	3
Butterfly Wound Closure	2

**Blisters, Splinters & Stings**

Insect Sting Relief Pad	4
Moleskin	1
Tweezers	1

**Bleeding, Wounds & Burns**

Non-Adherent Pad 2"x3"	2
Cotton Tipped Applicator	10
Sterile Gauze Pad 2"x2"	3
Sterile Gauze Pad 3"x3"	3
First Aid Tape 1/2"x2.5yds.	1
Medical Grade Gloves	2

**Headaches & Pain**

Ibuprofen 200mg	2
Non-Aspirin 325mg	4

**Sprains & Strains**

Elastic Bandage 2"	1
Instant Cold Compress	1
Assorted Safety Pins	3
Wooden Finger Splint	2

We reserve the right to add, replace or withdraw items as needed. We offer a 100% satisfaction guarantee.

**EASY CARE**

# First Aid Kit

SPORTS + TRAVEL

**Five Pockets Organized By Injury**

Be Prepared with Easy Care!

- 1 Cuts & Scrapes
- 2 Blisters, Splinters & Stings
- 3 Bleeding, Wounds & Burns
- 4 Headaches & Pain
- 5 Sprains & Strains

For Fast & Effective First Aid



**EASY CARE**

# First Aid Kit

SPORTS + TRAVEL

**Ideal for:**

- Sports Injuries
- Home First Aid
- Workshop Accidents
- Outdoor & Travel First Aid

Sports | Outdoor | Family | Travel

benzalkonium chloride, benzocaine, sd alcohol, bacitracin zinc, neomycin sulfate, polymyxin b, ibuprofen, acetaminophen kit

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:44224-0999
---------------------	----------------	---------------------------	----------------

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:44224-0999-5	1 in 1 KIT		

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	20 PACKAGE	16 mL
Part 2	4 PACKAGE	2.0 mL
Part 3	3 TUBE	1.5 g
Part 4	4 PACKAGE	8
Part 5	2 PACKET	4

### Part 1 of 5

#### ANTISEPTIC TOWELETTE

benzalkonium chloride swab

### Product Information

<b>Item Code (Source)</b>	NDC:52124-0001
---------------------------	----------------

<b>Route of Administration</b>	TOPICAL
--------------------------------	---------

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BENZALKONIUM CHLORIDE</b> (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.4 mL in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0001-1	0.8 mL in 1 PACKAGE		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333E	04/13/2011	

## Part 2 of 5

### INSECT STING RELIEF PAD

benzocaine,alcohol swab

## Product Information

Item Code (Source)	NDC:52124-0008
Route of Administration	TOPICAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	6 mL in 100 mL
ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)	ALCOHOL	60 mL in 100 mL

## Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0008-1	0.5 mL in 1 PACKAGE		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part348	04/13/2011	

## Part 3 of 5

### GENUINE TRIPLE ANTIBIOTIC

bacitracin zinc,neomycin sulfate,polymyxin b sulfate ointment

**Product Information**

<b>Item Code (Source)</b>	NDC:52124-0003
<b>Route of Administration</b>	TOPICAL

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN ZINC	400 [iU] in 1 g
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN SULFATE	5 mg in 1 g
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B SULFATE	5000 [iU] in 1 g

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
WATER (UNII: 059QF0K00R)	

**Packaging**

#	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:52124-0003-1	0.5 g in 1 TUBE		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph final	part333B	04/13/2011	

**Part 4 of 5****NON-ASPIRIN**

acetaminophen tablet

**Product Information**

<b>Item Code (Source)</b>	NDC:52124-0010
<b>Route of Administration</b>	ORAL

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
------------------------	-----------------

STARCH, CORN (UNII: O8232NY3SJ)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE (UNII: FZ989GH94E)	

### Product Characteristics

Color	white (WHITE)	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	AZ;234
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0010-1	2 in 1 PACKAGE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	04/13/2011	

## Part 5 of 5

### IBUPROFEN

ibuprofen tablet

### Product Information

Item Code (Source)	NDC:52124-0009
Route of Administration	ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

### Inactive Ingredients

Ingredient Name	Strength
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
LACTOSE (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYDEXTROSE (UNII: VH2XOU12IE)	

<b>POLYETHYLENE GLYCOL</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	

### Product Characteristics

<b>Color</b>	white (White)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10 mm
<b>Flavor</b>		<b>Imprint Code</b>	44;352
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0009-1	2 in 1 PACKET		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	02/17/2010	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part348	04/13/2011	

**Labeler** - Tender Corp dba Adventure Medical Kits (064437304)