

**IVERMECTIN SHEEP DRENCH- ivermectin solution**  
**Fast Sold LLC**

-----  
**Ivermectin Sheep Drench**  
**(ivermectin)**  
**0.08% Solution Parasiticide**

For the Treatment and Control of Worms and Bots of Sheep

- 20- 100 lb Doses (100 mL)

**See booklet for additional information.**

**Restricted Drug** (California) - Use Only as Directed

**For Animal Use Only • Keep Out of Reach of Children**

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

**FOR ORAL USE IN SHEEP ONLY. NOT FOR HUMAN USE.**

**PRODUCT DESCRIPTION**

Ivermectin Sheep Drench is a ready-to-use, free-flowing solution of ivermectin. It is formulated to deliver the recommended dose rate of 0.2 mg ivermectin per 1 kg body weight given orally at a volume of 3.0 mL per 26 lbs body weight.

**INDICATIONS:**

Ivermectin Sheep Drench provides treatment and control of adult and fourth-stage larvae of the following parasites: Gastrointestinal Roundworms – *Haemonchus contortus*, *Ostertagia circumcincta*, *Trichostrongylus axei*, *T. colubriformis*, *Cooperia curticei*, *Nematodirus spathiger*, *N. battus*, and *Oesophagostomum columbianum*; Lungworms – *Dictyocaulus filaria*; and all the larval stages of Nasal Bot – *Oestrus ovis*. It also provides treatment and control of adult forms only of the following Gastrointestinal Roundworms – *Haemonchus placei*, *Cooperia oncophora*, *Strongyloides papillosus*, *Oesophagostomum venulosum*, *Trichuris ovis*, and *Chabertia ovina*.

**DOSAGE AND ADMINISTRATION:**

Ivermectin Sheep Drench may be used in any standard drenching equipment or in any equipment which provides a consistent dose volume.  
Ivermectin Sheep Drench is administered orally at a dose of 3.0 mL (2.4 mg ivermectin) per 26 lbs body weight or 200 mcg ivermectin per kilogram of body weight.  
Coughing may be observed in some animals during and for several minutes following drenching.  
Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Underdosing may result in ineffective treatment, and encourage the development of parasite resistance.

## **RESIDUE WARNING:**

Do not treat sheep within 11 days of slaughter.

The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report adverse effects in users, to obtain an SDS, or for assistance call 1-5714367859.

## **PRECAUTIONS:**

Ivermectin (ivermectin) Sheep Drench has been formulated for use in sheep **only**. This product should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.

## **OTHER WARNINGS:**

Parasite resistance may develop to any dewormer, and has been reported for most classes of dewormers.

Treatment with a dewormer used in conjunction with parasite management practices appropriate to the geographic area and the animal(s) to be treated may slow the development of parasite resistance.

Fecal examinations or other diagnostic tests and parasite management history should be used to determine if the product is appropriate for the herd/flock, prior to the use of any dewormer. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of a fecal egg count reduction test or another appropriate method).

A decrease in a drug's effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. Your parasite management plan should be adjusted accordingly based on regular monitoring.

### **Keep this and all drugs out of reach of children.**

Refrain from smoking and eating when handling. Avoid contact with eyes. Immediately wash hands and any spills on the skin with plenty of soap and water following use.

**Restricted Drug** (California) - Use Only as Directed.

## **Environmental Safety:**

Studies indicate that when ivermectin comes in contact with the soil, it readily and tightly binds to the soil and becomes inactive over time. Free ivermectin may adversely affect fish and certain water-borne organisms on which they feed. Do not permit water runoff from feedlots to enter lakes, streams, or ground water. Do not contaminate water by direct application or by the improper disposal of drug containers. Spills should be contained and soaked up with absorbent towels or into loose soil. Gloves should be worn to prevent skin exposure. All the collected materials (contaminated towels and soil), as well as all empty drug containers, should be placed in an impervious film (plastic) bag and disposed of by incineration or in an approved landfill.

## **STORAGE INFORMATION:**

Store at 68° - 77° F (20° - 25° C). Excursions between 59° - 86° F (15° - 30° C) are permitted.

OZIPCO

IVERMECTIN

Liquid 0.08%

SHEEP DRENCH

NDC 86241-100-01

For the Treatment and Control of Worms and Bots of Sheep

See insert or booklet for additional information.

Use Only as Doses Section

For oral use in sheep only

Net Contents: 3.4 fl oz (100 mL)



**ozipco**  
**Ivermectin**  
 Liquid 0.08%  
 SHEEP DRENCH  
 NDC 86241-100-01

**ozipco**  
**Ivermectin**  
 Liquid 0.08%  
 SHEEP DRENCH  
 NDC 86241-100-01



For the Treatment and  
 Control of Worms  
 and Bots of Sheep

See insert or booklet for  
 Additional Information. Use  
 Only as Doses section

For oral use in sheep only

Net Wt. 3.4 fl oz (100 mL)

**INDICATIONS :**

For the treatment and control of gastrointestinal round worms, lungworms, and nasal bots in sheep.

**DOSAGE AND ADMINISTRATION :**

Administer orally at a dose of 1 mL per 50 lb (23 kg) body weight (Equivalent to 200 mcg ivermectin per kg body weight)

GTIN : 199874216478

**STORAGE INFORMATION:**

Store at 68°F-77°F (20°C-25°C). Excursions between 59°F-86°F (15°C-30°C)

**Distributed By :**

**Fast Sold Llc**

2280 AVOCADO AVE UNITE-5  
 MELBOURNE MELBOURNE, FL  
 32935-5547, United States  
 Phone number: +15714367859  
 support@ozipco.us  
 www.fastsoldllc.com

**ozipco**  
**Ivermectin**  
 Liquid 0.08%  
 SHEEP DRENCH  
 NDC 86241-100-01



For the Treatment and  
 Control of Worms  
 and Bots of Sheep

See insert or booklet for  
 Additional Information. Use  
 Only as Doses section

For oral use in sheep only

Net Wt. 3.4 fl oz (100 mL)

**WARNINGS :**

- Not for use in humans
- Keep out of reach of children
- Do not underdose
- Use only in sheep
- Consult a veterinarian for assistance in diagnosis and control of parasitism

Animals must not be slaughtered for human consumption within 11 days following treatment.

**PRECAUTIONS :**

- Use with caution in debilitated animals
- Do not use in other animal species
- Overdose may result in adverse effects

**Doses section :**

Sheep Weight (lb)	Sheep Weight (kg)	Doses (ml)
25 lb	11kg	0.5 ml
50 lb	23 kg	1.0 ml
75 lb	34 kg	1.5 ml
100 lb	45 kg	2.0 ml
125 lb	57 kg	2.5 ml
150 lb	68 kg	3.0 ml
175 lb	79 kg	3.5 ml
200 lb	91 kg	4.0 ml

Lot No :  
 Expired Date :

**ozipco**  
**Ivermectin**  
 Liquid 0.08%  
 SHEEP DRENCH  
 NDC 86241-100-01

**IVERMECTIN SHEEP DRENCH**

ivermectin solution

**Product Information**

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:86241-100
<b>Route of Administration</b>	Oral		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
IVERMECTIN (UNII: 8883YP2R6D) (IVERMECTIN - UNII:8883YP2R6D)	IVERMECTIN	0.8 mg in 1 mL

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:86241-100-01	1 in 1 BOX		
1		100 mL in 1 BOTTLE		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANADA	ANADA200327	01/30/2009	

**Labeler** - Fast Sold LLC (123898597)**Registrant** - Fast Sold LLC (123898597)**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Zhejiang Hisun Pharmaceutical Co., Ltd.		654211754	api manufacture

Revised: 4/2026

Fast Sold LLC