

ALEVE HEADACHE PAIN- naproxen sodium tablet
Bayer HealthCare LLC

Aleve ® Headache Pain

Active Ingredient

Naproxen Sodium 220mg

Purposes:

Pain reliever/fever reducer

Uses:

- temporarily relieves minor aches and pains due to:
 - headache
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - toothache
 - the common cold
- temporarily reduces fever

Warnings:

Allergy Alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do Not Use:

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a Doctor:

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat

- redness or swelling is present in the painful area
- any new symptoms appear

The pregnancy/breast-feeding warning:

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children:

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none">• take 1 tablet every 8 to 12 hours while symptoms last• for the first dose you may take 2 tablets within the first hour• do not exceed 2 tablets in any 8- to 12-hour period• do not exceed 3 tablets in a 24-hour period
Children under 12 years	<ul style="list-style-type: none">• ask a doctor

Other information:

- **each tablet contains:** sodium 20 mg
- store at 20-25 ° C (68-77 ° F). Avoid high humidity and excessive heat above 40 ° C (104 ° F).

Inactive ingredient

FD&C BLUE #2 LAKE, HYPROMELLOSE, MAGNESIUM STEARATE, MICROCRYSTALLINE CELLULOSE, POLYETHYLENE GLYCOL, POVIDONE, TALC, TITANIUM DIOXIDE

Questions & comments:

1-800-395-0689 (Mon-Fri 9AM-5PM EST)

Dist. by :

Bayer Healthcare LLC

Whippany, NJ 07981

PACKAGE LABEL PRINCIPAL DISPLAY PANEL - 24 Count Tablets

ALL DAY STRONG®

ALEVE

HEADACHE PAIN

naproxen sodium tablets, 220 mg (NSAID)

Pain reliever/fever reducer

Bayer

For temporary relief of headache pain

ACTUAL SIZE

STRENGTH TO LAST 12 HOURS

TABLETS



ALEVE HEADACHE PAIN

naproxen sodium tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0280-0041
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
TALC (UNII: 7SEV7J4R1U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	

Product Characteristics

Color	blue	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	ALEVE
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0280-0041-01	1 in 1 CARTON	06/01/2021	
1		175 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0280-0041-02	1 in 1 CARTON	06/01/2021	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0280-0041-04	1 in 1 CARTON	06/01/2021	
3		24 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:0280-0041-03	1 in 1 CARTON	06/01/2021	

4		90 in 1 BOTTLE; Type 0: Not a Combination Product	
5	NDC:0280-0041-05	1 in 1 POUCH; Type 0: Not a Combination Product	06/01/2021



Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020204	06/01/2021	

Labeler - Bayer HealthCare LLC (112117283)

Revised: 12/2025

Bayer HealthCare LLC