

M-PAP- acetaminophen liquid
Bryant Ranch Prepack

M-PAP

M-PAP

Acetaminophen Liquid

160 mg/5 mL

Pain Reliever / Fever Reducer

Sugar Free / Alcohol Free

Aspirin Free

CHERRY FLAVOR

Drug Facts

Active ingredient

(in each 5 mL teaspoonful)

Acetaminophen, USP 160 mg

Purpose

Pain reliever/fever reducer

Uses

- reduces fever
- temporarily relieves minor aches and pains due to:
 - Headache
 - muscular aches
 - backache
 - arthritis
 - the common cold
 - toothache
 - menstrual cramps
 - reduces fever

Warnings

Liver Warning: This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 8 teaspoonfuls (40 mL) in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor
- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use

if you have health issues especially liver disease.

Ask a doctor or pharmacist before use

if you are taking other drugs, including the blood thinner warfarin.

When using this product do not exceed recommended dose (see Overdose warning)

Stop use and ask a doctor if

- new symptoms occur such as rash, hives, itching or hoarseness
- redness or swelling is present
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- symptoms do not improve

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical even if you do not notice any signs or symptoms.

Directions

Do not exceed recommended dosage.

- **adults and children 12 years of age and older:** Take 3 teaspoonfuls (15 mL) every 5 hours; do not exceed 15 teaspoonfuls (75 mL) in 24 hours
- **children under 12 years of age:** Under the direct guidance of a licensed professional, doctor, or pharmacist.

Other Information

- **each 5 mL contains:** Sodium 3 mL

If dispensed, dispense in a tight, light resistant container with a child-resistant cap. Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F).

Inactive Ingredients

Anhydrous citric acid, FD&C Blue #1, FD&C Red #40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solution.

Questions?

1-877-250-3427 M-F (8 a.m. to 5 p.m. CST), or directly to Medwatch at 1-800-332-1088.

Serious side effects associated with use of this product may be reported to this number.

Distributed by:

Method Pharmaceuticals

Southlake, TX 76092

Rev. 02/2024

HOW SUPPLIED

M-PAP Acetaminophen Liquid 160 mg/5 mL

- NDC: 72162-2404-2: 473 mL in a BOTTLE
- NDC: 72162-2404-4: 118 mL in a BOTTLE

If dispensed, dispense in a tight, light resistant container with a child-resistant cap. Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F).

Repackaged/Relabeled by:

Bryant Ranch Prepack, Inc.

Burbank, CA 91504

M-PAP Acetaminophen Liquid 160 mg/5 ml



Lot
Exp
GTIN

Drug Facts	
Active ingredient (in each 5 mL teaspoonful)	Purpose
Acetaminophen, USP 160 mg	Pain reliever/fever reducer
Uses	
•reduces fever •temporarily relieves minor aches and pains due to: •Headache •muscular aches •backache •arthritis •the common cold •toothache •menstrual cramps	
Warnings	
Liver Warning: This product contains acetaminophen. Severe liver damage may occur if you take: •more than 8 teaspoonfuls (40 mL) in 24 hours, which is the maximum daily amount •with other drugs containing acetaminophen •3 or more alcoholic drinks every day while using this product. Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening •blisters •rash. If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly. Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. •for more than 10 days for pain unless directed by a doctor •for more than 3 days for fever unless directed by a doctor •if you are allergic to acetaminophen or any of the inactive ingredients in this product. Ask a doctor before use if you have health issues especially liver disease. Ask a doctor or pharmacist before use if you are taking other drugs, including the blood thinner warfarin. When using this product do not exceed recommended dose (see Overdose warning). Stop use and ask a doctor if •new symptoms occur such as rash, hives, itching or hoarseness •redness or swelling is present •pain gets worse or lasts for more than 10 days •fever gets worse or lasts for more than 3 days •symptoms do not improve. These could be signs of a serious condition.	
Other Information	
•each 5 mL contains: Sodium 3 mL •if dispensed, dispense in a tight, light resistant container with a child-resistant cap. Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (between 59°F and 86°F). Cherry Flavor.	
Directions	
Do not exceed recommended dosage. •adults and children 12 years of age and older: Take 3 teaspoonfuls (15 mL) every 5 hours; do not exceed 15 teaspoonfuls (75 mL) in 24 hours. •children under 12 years of age: Under the direct guidance of a licensed professional, doctor, or pharmacist.	
Inactive Ingredients	
Anhydrous citric acid, FD&C Blue #1, FD&C Red #40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solution.	



Package
Insert

NDC 72162-2404-4

M-PAP Acetaminophen Liquid

160 mg/5 mL



Relabeled by:
Bryant Ranch Prepack, Inc.
Burbank, CA 91504 USA

4 FL OZ (118 mL)

Manufactured by:
AptaPharma Inc.



7216224044

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical even if you do not notice any signs or symptoms. Do not use if tamper evident seal under cap is broken or missing. Scan Package Insert QR Code for additional information.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

M-PAP

acetaminophen liquid

Product Information				
Product Type		HUMAN OTC DRUG	Item Code (Source)	NDC:72162-2404(NDC:58657-524)
Route of Administration		ORAL		
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)			ACETAMINOPHEN	160 mg in 5 mL
Inactive Ingredients				
Ingredient Name				Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
WATER (UNII: 059QF0KO0R)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
SACCHARIN SODIUM (UNII: SB8ZUX40TY)				
SORBITOL (UNII: 506T60A25R)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
Product Characteristics				
Color			Score	
Shape			Size	
Flavor		CHERRY	Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72162-2404-2	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/10/2024	
2	NDC:72162-2404-4	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/10/2024	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M013	04/01/2024	

Labeler - Bryant Ranch Prepack (171714327)

Registrant - Bryant Ranch Prepack (171714327)

Establishment

Name	Address	ID/FEI	Business Operations
Bryant Ranch Prepack		171714327	REPACK(72162-2404) , RELABEL(72162-2404)

Revised: 4/2025

Bryant Ranch Prepack