

NAPROXEN SODIUM- naproxen sodium tablet, film coated
Rite Aid Corporation

Rite Aid Corporation Naproxen Sodium Drug Facts

Active ingredient (in each caplet)

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing

- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

| | |
|--|--|
| Adults and children 12 years and older | <ul style="list-style-type: none"> • take 1 caplet every 8 to 12 hours while symptoms last • for the first dose you may take 2 caplets within the first hour • do not exceed 2 caplets in any 8- to 12-hour period • do not exceed 3 caplets in a 24-hour period |
| Children under 12 years | <ul style="list-style-type: none"> • ask a doctor |

Other information

- **each caplet contains:** sodium 22 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to the active ingredient of Aleve® Caplets

FREE FRV/M | GLUTEN FREE

STRENGTH TO LAST 12 HOURS

EASY OPEN ARTHRITIS CAP

NAPROXEN SODIUM TABLETS, 220 mg

PAIN RELIEVER/FEVER REDUCER (NSAID)

ACTUAL SIZE

PACKAGE NOT CHILD-RESISTANT

270 CAPLETS ♦

♦ capsule-shaped tablets

Drug Facts (continued)

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer right before or after heart surgery

Ask a doctor before use if the stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

NDC 11822-0140-1

Compare to the active ingredient of Aleve® Caplets**

FREE FROM GLUTEN FREE

STRENGTH TO LAST 12 HOURS

EASY OPEN ARTHRITIS CAP

NAPROXEN SODIUM

TABLETS, 220 mg

PAIN RELIEVER/FEVER REDUCER (NSAID)

ACTUAL SIZE

PACKAGE NOT CHILD-RESISTANT

270 CAPLETS!
capsule-shaped tablets

Drug Facts

| | |
|--|--|
| Active ingredient (in each caplet) Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)* | Purposes Pain reliever/fever reducer |
|--|--|

*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to: minor pain of arthritis, muscular aches, backache, menstrual cramps, headache, toothache, the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older

0 11822 58937 6

PEEL BACK HERE

DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

140H8 83 F3

Drug Facts (continued)

- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin taking any other drug

When using this product take with food or milk if stomach upset occurs

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling, pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, you have difficulty swallowing, it feels like the pill is stuck in your throat, redness or swelling is present in the painful area, any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions do not take more than directed

Drug Facts (continued)

- the smallest effective dose should be used
- drink a full glass of water with each dose

| | |
|--|--|
| Adults and children 12 years and older | <ul style="list-style-type: none"> take 1 caplet every 8 to 12 hours while symptoms last for the first dose you may take 2 caplets within the first hour do not exceed 2 caplets in any 8- to 12-hour period do not exceed 3 caplets in a 24-hour period |
| Children under 12 years | ask a doctor |

Other information

- each caplet contains: sodium 22 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients: FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments? 1-800-719-9260

**This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Aleve® Caplets.

DISTRIBUTED BY: RITE AID
30 HUNTER LANE, CAMP HILL, PA 17011 | www.riteaid.com

SATISFACTION GUARANTEE: If you're not satisfied, we'll happily refund your money.

ADHESIVE AREA
NO COPY

NAPROXEN SODIUM

naproxen sodium tablet, film coated

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:11822-0140 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|--------------------------|-----------------|
| NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ) | NAPROXEN SODIUM | 220 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|-----------------|
| FD&C BLUE NO. 2 (UNII: L06K8R7DQK) | |
| HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO) | |
| MAGNESIUM STEARATE (UNII: 70097M6130) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| POVIDONE, UNSPECIFIED (UNII: FZ989GH94E) | |
| TALC (UNII: 7SEV7J4R1U) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |

Product Characteristics

| | | | |
|-----------------|-------------------|---------------------|----------|
| Color | BLUE (light blue) | Score | no score |
| Shape | OVAL | Size | 12mm |
| Flavor | | Imprint Code | L368 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:11822-0140-1 | 270 in 1 BOTTLE; Type 0: Not a Combination Product | 07/01/2016 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------------|---|-----------------------------|---------------------------|
| ANDA | ANDA074661 | 07/01/2016 | |

Labeler - Rite Aid Corporation (014578892)

