

MELOXICAM- mecoxican tablet

Preferred Pharmaceuticals Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use MELOXICAM TABLETS safely and effectively. See full prescribing information for MELOXICAM TABLETS.

MELOXICAM tablets, for oral use

Initial U.S. Approval: 2000

WARNING: RISK OF SERIOUS CARDIOVASCULAR AND GASTROINTESTINAL EVENTS

See full prescribing information for complete boxed warning.

- Nonsteroidal anti-inflammatory drugs (NSAIDs) cause an increased risk of serious cardiovascular thrombotic events, including myocardial infarction and stroke, which can be fatal. This risk may occur early in treatment and may increase with duration of use (1).
- NSAIDs are contraindicated in the setting of coronary artery bypass graft (CABG) surgery (1).
- NSAIDs cause an increased risk of serious gastrointestinal (GI) events, including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients and patients with a prior history of peptic ulcer disease and/or GI bleeding are at greater risk for serious GI events (5.2).

RECENT MAJOR CHANGES

07/2024

INDICATIONS AND USAGE

Meloxicam tablets are a non-steroidal anti-inflammatory drug indicated for (1):

- Osteoarthritis (OA) (1)
- Rheumatoid Arthritis (RA) (1)
- Juvenile Rheumatoid Arthritis (JRA) in patients who weigh ≥ 60 kg (1)

DOSAGE AND ADMINISTRATION

Use the lowest effective dosage for the shortest duration consistent with individual patient treatment goals (1).

- OA (1) and RA (1):

Starting dose: 7.5 mg once daily (2)

Dose may be increased to 15 mg once daily (2)

- RA (1):

7.5 mg once daily in children ≥ 60 kg (2)

Meloxicam Tablets are not interchangeable with approved formulations of oral mecoxican even if the total mifam strength is the same (2).

DOSEAGE FORMS AND STRENGTHS

• Meloxicam Tablets USP: 7.5 mg and 15 mg (1).

CONTRAINDICATIONS

- Known hypersensitivity to mecoxican or any components of the drug product (1).
- History of asthma, urticaria, or other allergic-type reactions after taking aspirin or other NSAIDs (1).
- In the setting of CABG surgery (1).

WARNINGS AND PRECAUTIONS

- Hematology:** Inform patients of warning signs and symptoms of hepatotoxicity. Discontinue if abnormal liver function or evidence of liver or clinical signs and symptoms of liver disease develop (1).
- Hypertension:** Patients taking some antihypertensive medications may have impaired response to these medications when taking NSAIDs (1).
- Heart Failure and Edema:** Avoid use of Meloxicam in patients with severe heart failure unless benefits are expected to outweigh the risk of edema (1).
- Renal Toxicity:** Monitor renal function in patients with renal or hepatic impairment, heart failure, or hypertension. If renal function is significantly impaired, discontinue Meloxicam unless benefits are expected to outweigh risk of worsening renal function (1).
- Anaphylactic Reactions:** Seek emergency help if an anaphylactic reaction occurs (1).
- Exacerbation of Asthma:** Monitor patients with preexisting asthma (without aspirin sensitivity) (1).
- Severe Hypersensitivity:** Inform patients of signs and symptoms of severe hypersensitivity (1).
- Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS):** Discontinue Meloxicam and evaluate clinically (1).
- Use of NSAIDs:** Between about 20 to 30 weeks in pregnancy due to the risk of oligohydramnios/fetal renal dysplasia. Avoid use of NSAIDs in women at about 30 weeks of gestation due to the risks of oligohydramnios/fetal renal dysplasia and premature closure of the fetal ductus arteriosus (1).
- Hemorrhage/Toxicity:** Monitor hemoglobin or hematocrit in patients with any signs or symptoms of anemia (1).

ADVERSE REACTIONS

- Most common ($\geq 5\%$ and greater than placebo) adverse events in adults are diarrhea, upper respiratory tract infections, dyspepsia, and influenza-like symptoms (1).
- Adverse events observed in pediatric studies were similar in nature to the adult clinical trial experience (1).

To report SUSPECTED ADVERSE REACTIONS, contact Preferred Pharmaceuticals (USA), Inc. at 866-444-3633 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. (1)

DRUG INTERACTIONS

- Drugs that interfere with Hemostasis (e.g., warfarin, aspirin, SSRIs/NBSS): Monitor patients for bleeding who are taking Meloxicam and these drugs (1).
- Concomitant use of Meloxicam and any NSAID doses of aspirin is not generally recommended (1).
- ACE Inhibitors:** Angiotensin Receptor Blockers (ARBs) or Beta-Blockers: Concomitant use with Meloxicam may increase the risk of hypotension and/or orthostatic hypotension (1).
- ACE Inhibitors and ARBs:** Concomitant use with Meloxicam in elderly, volume-depleted, or those with compromised renal function may increase the risk of hypotension and/or orthostatic hypotension (1).
- Diuretics:** Monitor for signs of worsening renal function (1).
- Diuretics and Antidiabetics:** Diuretic effect of furosemide and thiazide diuretics. Monitor patients to assure diuretic efficacy including antidiabetics effects (1).

USE IN SPECIFIC POPULATIONS

- Infertility:** NSAIDs are associated with reversible infertility. Consider withdrawal of Meloxicam in women who have difficulties conceiving (1).

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide. Revised: 10/2024

FULL PRESCRIBING INFORMATION: CONTENTS***1 INDICATIONS AND USAGE**

- Osteoarthritis (OA) (1)
- Rheumatoid Arthritis (RA) (1)
- Juvenile Rheumatoid Arthritis (JRA) Pauciarticular and Polyarticular Course (1)

2 DOSAGE AND ADMINISTRATION

- Osteoarthritis Instructions (1)
- 2 Osteoarthritis (1)
- 2 Rheumatoid Arthritis (1)
- 2 Juvenile Rheumatoid Arthritis (JRA) Pauciarticular and Polyarticular Course (1)
- 2.5 Non-Interchangeability (1)
- 2.6 Non-Interchangeability with Other Formulations of Meloxicam (1)

3 DOSAGE FORMS AND STRENGTHS**4 CONTRAINDICATIONS****5 WARNINGS AND PRECAUTIONS**

- 5.1 Cardiovascular Thrombotic Events (1)
- 5.2 Gastrointestinal Bleeding, Ulceration, and Perforation (1)
- 5.3 Hepatotoxicity (1)
- 5.4 Hypertension (1)
- 5.5 Masking of Inflammation and Edema (1)
- 5.6 Renal Toxicity and Hyperkalemia (1)
- 5.7 Anaphylactic Reactions (1)
- 5.8 Exacerbation of Asthma Related to Aspirin Sensitivity (1)
- 5.9 Skin Reactions (1)
- 5.10 Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS) (1)
- 5.11 Fetal Toxicity (1)
- 5.12 Pediatric Toxicity (1)
- 5.13 Masking of Inflammation and Fever (1)
- 5.14 Laboratory Monitoring (1)

6 ADVERSE REACTIONS

- 6.1 Clinical Trial Experience (1)
- 6.2 Post-Marketing Experience (1)

7 DRUG INTERACTIONS**8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy (1)
- 8.2 Lactation (1)
- 8.3 Females and Males of Reproductive Potential (1)
- 8.4 Pediatric Use (1)
- 8.5 Geriatric Use (1)
- 8.6 Hepatic Impairment (1)
- 8.7 Renal Impairment (1)

10 OVERDOSAGE**11 DOSAGE FORMS****12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action (1)
- 12.2 Pharmacokinetics (1)

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility (1)

14 CLINICAL STUDIES

- 14.1 Osteoarthritis and Rheumatoid Arthritis (1)
- 14.2 Juvenile Rheumatoid Arthritis (JRA) Pauciarticular and Polyarticular Course (1)

16 HOW SUPPLIED/STORAGE AND HANDLING**17 PATIENT COUNSELING INFORMATION**

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: RISK OF SERIOUS CARDIOVASCULAR AND GASTROINTESTINAL EVENTS	
Cardiovascular Thrombotic Events	
Nonsteroidal anti-inflammatory drugs (NSAIDs) cause an increased risk of serious cardiovascular thrombotic events, including myocardial infarction and stroke, which can be fatal. This risk may occur early in treatment and may increase with duration of use (see <i>Warnings and Precautions</i> 1).	
Meloxicam tablets are contraindicated in the setting of coronary artery bypass graft (CABG) surgery (see <i>Contraindications</i> 1 and <i>Warnings and Precautions</i> 1).	
Gastrointestinal Bleeding, Ulceration, and Perforation	
NSAIDs cause an increased risk of serious gastrointestinal (GI) adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients and patients with a prior history of peptic ulcer disease and/or GI bleeding are at greater risk for serious GI events (see <i>Warnings and Precautions</i> 5.2).	

1 INDICATIONS AND USAGE

- Osteoarthritis (OA) (1)

Meloxicam tablets are indicated for relief of the signs and symptoms of osteoarthritis (see *Clinical Studies* 1).**1.2 Rheumatoid Arthritis (RA)**Meloxicam tablets are indicated for relief of the signs and symptoms of rheumatoid arthritis (see *Clinical Studies* 1).**1.3 Juvenile Rheumatoid Arthritis (JRA) Pauciarticular and Polyarticular Course**Meloxicam tablets are indicated for relief of the signs and symptoms of pauciarticular or polyarticular course Juvenile Rheumatoid Arthritis in patients who weigh ≥ 60 kg (see *Dosage and Administration* 1 and *Clinical Studies* 1).

2 DOSAGE AND ADMINISTRATION

2.1 General Dosing Instructions

Carefully consider the potential benefits and risks of Meloxicam tablets, and other treatment options before deciding to use Meloxicam tablets. Use the lowest effective dosage for the shortest duration consistent with individual patient treatment goals [see Warnings and Precautions (5)].

After observing the response to initial therapy with Meloxicam tablets, adjust the dose to suit the patient. In adults, the maximum recommended daily oral dose of Meloxicam tablets is 15 mg regardless of formulation. In patients with hemodialysis, a maximum daily dosage of 7.5 mg is recommended [see Use in Specific Populations (1) and Clinical Pharmacology (1)]. Meloxicam tablets may be taken without regard to timing of meals.

2.2 Osteoarthritis

For the relief of the signs and symptoms of osteoarthritis the recommended starting and maintenance oral dose of Meloxicam tablets is 7.5 mg once daily. Some patients may receive additional benefit by increasing the dose to 15 mg once daily.

2.3 Rheumatoid Arthritis

For the relief of the signs and symptoms of rheumatoid arthritis, the recommended starting and maintenance oral dose of Meloxicam tablets is 7.5 mg once daily. Some patients may receive additional benefit by increasing the dose to 15 mg once daily.

2.4 Juvenile Rheumatoid Arthritis (JRA) Pauciarticular and Polyarticular Course

For the treatment of juvenile rheumatoid arthritis, the recommended oral dose of Meloxicam tablets is 7.5 mg once daily in children who weigh ≥ 60 kg. There was no additional benefit demonstrated by increasing the dose above 7.5 mg in clinical trials. Meloxicam tablets should not be used in children who weigh <60 kg.

2.5 Renal Impairment

The use of Meloxicam tablets in subjects with severe renal impairment is not recommended.

In patients on hemodialysis, the maximum dosage of Meloxicam tablets is 7.5 mg per day [see Clinical Pharmacology (1)].

2.6 Non-Interchangeability with Other Formulations of Meloxicam

Meloxicam tablets have not shown equivalent systemic exposure to other approved formulations of oral meloxicam. Therefore, Meloxicam tablets are not interchangeable with other formulations of oral meloxicam product even if the total daily dose is the same. Do not substitute dose strengths of Meloxicam tablets with other formulations of oral meloxicam product.

3 DOSAGE FORMS AND STRENGTHS

Meloxicam Tablets USP:

- 7.5 mg: Light yellow, round flat beveled edged, tablet with U & L debossed on one side and 7.5 debossed centrally on the other side
- 15 mg: Light yellow, capsule shaped, biconvex, tablet with U & L debossed on one side and 15 debossed centrally on the other side

4 CONTRAINDICATIONS

Meloxicam tablets are contraindicated in the following patients:

- Known hypersensitivity (e.g., anaphylactic reactions and serious skin reactions) to meloxicam or any components of the drug product [see Warnings and Precautions (5).]
- History of asthma, urticaria, or other allergic-type reactions after taking aspirin or other NSAIDs. Severe, sometimes fatal, anaphylactic reactions to NSAIDs have been reported in such patients [see Warnings and Precautions (5).]
- In the setting of coronary artery bypass graft (CABG) surgery [see Warnings and Precautions (5).]

5 WARNINGS AND PRECAUTIONS

5.1 Cardiovascular Thrombotic Events

Clinical trials of several COX-2 selective and non-selective NSAIDs of up to three years duration have shown an increased risk of serious cardiovascular (CV) thrombotic events, including myocardial infarction (MI) and stroke, which can be fatal. Based on available data, it is unclear that the risk for CV thrombotic events is similar for all NSAIDs. The risk of serious CV thrombotic events appears to increase with increasing duration of use and appears to be similar in those with and without known CV disease or risk factors for CV disease. However, patients with known CV disease or risk factors had a higher absolute incidence of excess serious CV thrombotic events, due to their increased baseline risk. Most trials have not found a consistent increase in serious CV thrombotic events began as early as the first weeks of treatment. The increase in CV thrombotic risk has been observed most consistently at higher doses.

To minimize the potential risk for an adverse CV event in NSAID-treated patients, use the lowest effective dose for the shortest possible duration. Patients and patients should remain alert for signs of such events, throughout the entire treatment course, even in the absence of previous CV symptoms. Patients should be informed about the symptoms of serious CV events and the steps to take if they occur.

There is no consistent evidence that concurrent use of aspirin mitigates the increased risk of serious CV thrombotic events in patients treated with NSAIDs. Aspirin and an NSAID, such as meloxicam, increase the risk of serious gastrointestinal (GI) events [see Warnings and Precautions (5.2)].

5.1.1 Status Post Coronary Artery Bypass Graft (CABG) Surgery

Two large, controlled clinical trials of a COX-2 selective NSAID for the treatment of pain in the days following CABG surgery found an increased incidence of myocardial infarction and stroke. NSAIDs are contraindicated in the setting of CABG [see Contraindications (4)].

5.1.2 Post-MI Patients

Observational studies conducted in the Danish National Registry have demonstrated that patients treated with NSAIDs in the post-MI period were at increased risk of recurrent CV-related death and/or cardiovascular events in the first year of treatment. In this same cohort, the incidence of death in the first year post-MI was 20 per 100 person years in NSAID-treated patients compared to 12 per 100 person years in non-NSAID exposed patients. Although the absolute rate of death declined somewhat after the first year post-MI, the increased relative risk of death in NSAID users persisted over at least the next four years of follow-up.

Avoid the use of Meloxicam in patients with a recent MI unless the benefits are expected to outweigh the risk of recurrent CV thrombotic events. If Meloxicam is used in patients with a recent MI, monitor patients for signs of cardiac ischemia.

5.2 Gastrointestinal Bleeding, Ulceration, and Perforation

NSAIDs, including meloxicam, can cause serious GI (mucosal) adverse events, including inflammation, bleeding, ulceration, and perforation of the esophagus, stomach, small intestine, or large intestine, which can be fatal. These serious adverse events can occur at any time, with or without warning symptoms, in patients treated with NSAIDs. Only one in five patients who develop a serious upper GI adverse event on NSAID therapy are symptomatic. The remaining four-fifths of patients are asymptomatic, and about 2-4% of patients treated for one year. However, even short-term NSAID therapy is not without risk.

5.2.1 Risk Factors for GI Bleeding, Ulceration, and Perforation

Patients with a prior history of peptic ulcer disease and/or GI bleeding who used NSAIDs had a greater than 10-fold increased risk for developing a GI bleed compared to patients without these risk factors. Other factors that increase the risk of GI bleeding in patients treated with NSAIDs include longer duration of NSAID therapy, concurrent use of oral corticosteroids, smoking, use of alcohol, older age, and poor general health status. Most postmarketing reports of fatal GI events occurred in elderly or debilitated patients. Additionally, patients with advanced liver disease and/or coagulopathy are at increased risk for GI bleeding.

5.2.2 Strategies to Minimize the GI Risks in NSAID-treated Patients

- Use the lowest effective dosage for the shortest possible duration.
- Avoid administration of more than one NSAID at a time.
- Avoid use of patients at higher risk unless benefits are expected to outweigh the increased risk for bleeding, such as in those with active GI bleeding, consider alternate therapies other than NSAIDs.
- Remain alert for signs and symptoms of GI ulceration and bleeding during NSAID therapy.
- If a serious GI adverse event is suspected, promptly initiate evaluation and treatment, and discontinue Meloxicam until a serious GI adverse event is ruled out.
- In the setting of concomitant use of low-dose aspirin for cardiac prophylaxis, monitor patients more closely for evidence of GI bleeding [see Drug Interactions (1)].

5.3 Hepatotoxicity

Elevations of ALT or AST (three or more times the upper limit of normal [ULN]) have been reported in approximately 1% of NSAID-treated patients in clinical trials. In addition, rare, sometimes fatal, cases of severe hepatic injury, including fulminant hepatitis, liver necrosis, and hepatic failure have been reported.

Elevations of ALT or AST (less than three times ULN) may occur in up to 15% of patients treated with NSAIDs including meloxicam.

Inform patients of the warning signs and symptoms of hepatotoxicity (e.g., nausea, fatigue, lethargy, diarrhea, pruritus, jaundice, right upper quadrant tenderness, and "flu-like" symptoms). If clinical signs and symptoms consistent with liver disease develop, or if a patient develops any new symptom while taking meloxicam, discontinue meloxicam immediately, and perform a clinical evaluation of the patient [see Use in Specific Populations (1) and Clinical Pharmacology (1)].

5.4 Hypertension

NSAIDs, including Meloxicam, can lead to new onset or worsening of preexisting hypertension, either of which may contribute to the increased incidence of CV events. Patients taking angiotensin-converting enzyme (ACE) inhibitors, thiazide diuretics, or loop diuretics may have impaired response to these therapies when taking NSAIDs [see Drug Interactions (1)].

Monitor blood pressure (BP) during the initiation of NSAID treatment and throughout the course of therapy.

5.5 Heart Failure and Edema

The Coxib and traditional NSAID Trialists' Collaboration meta-analysis of randomized controlled trials demonstrated an approximately two-fold increase in hospitalizations for heart failure in COX-2 selective-treated patients and non-selective NSAID-treated patients compared to placebo-treated patients. In a Danish National Registry study of patients with heart failure, NSAID use increased the risk of MI, hospitalization for heart failure, and death.

Additionally, fluid retention and edema have been observed in some patients treated with NSAIDs. Use of meloxicam may blunt the CV effects of several therapeutic agents used

to treat these medical conditions (e.g., diuretics, ACE inhibitors, or angiotensin receptor blockers [ARBs]) (see *Drug Interactions* 0).

Avoid the use of Meloxicam in patients with severe heart failure unless the benefits are expected to outweigh the risk of worsening heart failure. If Meloxicam is used in patients with severe heart failure, monitor patients for signs of worsening heart failure.

5.6 Renal Toxicity and Hyperkalemia

Renal Toxicity

Long-term administration of NSAIDs, including Meloxicam, has resulted in renal papillary necrosis, renal insufficiency, acute renal failure, and other renal injury.

Renal toxicity has also been seen in patients in whom renal insufficiencies have a compensatory role in the maintenance of renal perfusion. In these patients, administration of an NSAID may cause a dose-dependent reduction in prostaglandin formation and, secondarily, in renal blood flow, which may precipitate overt renal decompression. Patients with preexisting renal impairment, those with reduced renal function, dehydration, hypovolemia, heart failure, liver dysfunction, those taking diuretics and ACE inhibitors or ARBs, and the elderly. Discontinuation of NSAID therapy is usually followed by recovery to the pretreatment state.

The renal effects of Meloxicam may hasten the progression of renal dysfunction in patients with preexisting renal disease. Because some Meloxicam metabolites are excreted by the kidney, monitor patients for signs of worsening renal function.

Correct volume status in dehydrated or hypovolemic patients prior to initiating Meloxicam. Monitor renal function in patients with renal or hepatic impairment, heart failure, and in patients with hypovolemia (see *Warnings and Precautions* 0).

No information is available from controlled clinical studies regarding the use of Meloxicam in patients with advanced renal disease. Avoid the use of Meloxicam in patients with advanced renal disease unless the benefits are expected to outweigh the risk of worsening renal function. If Meloxicam is used in patients with advanced renal disease, monitor patients for signs of worsening renal function (see *Clinical Pharmacology* 0).

Hyperkalemia

Increases in serum potassium concentration, including hyperkalemia, have been reported with use of NSAIDs, even in some patients without renal impairment. In patients with preexisting renal impairment, these effects have been attributed to a hyperkrenic-hypoaldosteronism state.

5.7 Anaphylactic Reactions

Meloxicam can be associated with anaphylactic reactions in patients with and without known hypersensitivity to meloxicam and in patients with aspirin-sensitive asthma (see *Contraindications* 0) and *Warnings and Precautions* 0).

Seek emergency help if an anaphylactic reaction occurs.

5.8 Escalation of Asthma Related to Aspirin Sensitivity

A subpopulation of patients with asthma may have aspirin-sensitive asthma which may include chronic rhinosinusitis complicated by nasal polyps; severe, potentially fatal bronchospasm; and/or intolerance to aspirin and other NSAIDs. Because cross-reactivity between aspirin and other NSAIDs has been reported in such aspirin-sensitive patients, avoid the use of other NSAIDs in this form of aspirin sensitivity (see *Contraindications* 0). When Meloxicam is used in patients with preexisting asthma (without known aspirin sensitivity), monitor patients for changes in the signs and symptoms of asthma.

5.9 Serious Skin Reactions

NSAIDs, including meloxicam, can cause serious skin adverse reactions, such as exfoliative dermatitis, Steven's Johnson Syndrome (SJS), and toxic epidermal necrolysis (TEN), which can be fatal. NSAIDs can also cause fixed drug eruption (FDE). FDE may present as a more severe variant known as generalized bullous fixed drug eruption (GBFDE), which can be life-threatening. These serious events may occur without warning and may occur in patients who have never taken the drug before, and to discontinue the use of Meloxicam at the first appearance of skin rash or any other sign of hypersensitivity. Meloxicam is contraindicated in patients with previous serious skin reactions to NSAIDs (see *Contraindications* 0).

5.10 Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)

Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS) has been reported in patients taking NSAIDs, such as meloxicam. Some of these reactions have been fatal or life-threatening. DRESS typically, although not exclusively, presents with fever, rash, lymphadenopathy, and/or facial swelling. Other clinical manifestations may include hepatitis, nephritis, hematological abnormalities, myocarditis, or myositis. Symptoms of DRESS may be preceded by an acute hypersensitivity reaction and/or present. Because this disorder is variable in its presentation, other organ systems not noted here may be involved. It is important to note that early manifestations of hypersensitivity, such as fever or lymphadenopathy, may be present even though rash is not evident. If such signs or symptoms are present, discontinue meloxicam and evaluate the patient immediately.

5.11 Fetal Toxicity

Premature Closure of Fetal Ductus Arteriosus

Avoid use of NSAIDs, including meloxicam, in pregnant women at about 30 weeks gestation and later. NSAIDs, including meloxicam, increase the risk of premature closure of the fetal ductus arteriosus at approximately this gestational age.

Oligohydramnios/Neonatal Birth Impairment

Use of NSAIDs, including meloxicam, at about 20 weeks gestation or later in pregnancy may result in oligohydramnios and/or oligohydramnios, and, in some cases, neonatal renal impairment. These adverse outcomes are seen, on average, after days to weeks of treatment, although oligohydramnios has been infrequently reported as soon as 4 days after NSAID initiation. Oligohydramnios is often, but not always, reversible with treatment discontinuation. Complications of prolonged oligohydramnios, for example, include limb contractures and delayed lung maturation. In some postmarketing cases of impaired neonatal renal function, invasive procedures such as exchange transfusions were required.

If NSAID treatment is necessary between about 20 weeks and 30 weeks gestation, limit meloxicam use to the lowest effective dose and shortest duration possible. Consider ultrasound monitoring of amniotic fluid if meloxicam treatment extends beyond 48 hours. Discontinue meloxicam if oligohydramnios occurs and follow up according to clinical practice (see *Use in Specific Populations* 0).

5.12 Hematologic Toxicity

Anemia has occurred in NSAID-treated patients. This may be due to occult or gross blood loss, fluid retention, or an incompletely described effect on erythropoiesis. If a patient treated with Meloxicam has any signs or symptoms of anemia, monitor hemoglobin or hematocrit.

NSAIDs, including meloxicam, may increase the risk of bleeding events. Coarcted conditions, coagulation disorders or concurrent use of warfarin, other anticoagulants, antiplatelet agents (e.g., aspirin), serotonin reuptake inhibitors (SSRIs) and serotonin norepinephrine reuptake inhibitors (SNRIs) may increase this risk. Monitor these patients for signs of bleeding (see *Drug Interactions* 0).

5.13 Masking of Inflammation and Fever

The pharmacological activity of Meloxicam in reducing inflammation, and possibly fever, may diminish the utility of diagnostic signs in detecting infections.

5.14 Laboratory Monitoring

Because serious GI bleeding, hepatotoxicity, and renal injury can occur without warning symptoms or signs, consider monitoring patients on long-term NSAID treatment with a CBC and a chemistry profile periodically (see *Warnings and Precautions* 5.2, 7).

6 ADVERSE REACTIONS

The following adverse reactions are discussed in greater detail in other sections of the labeling:

- Cardiovascular Thrombotic Events (see *Boxed Warning* and *Warnings and Precautions* 0)
- GI Bleeding, Ulceration, and Perforation (see *Boxed Warning* and *Warnings and Precautions* 0, 2, 3)
- Hemorrhage (see *Warnings and Precautions* 0)
- Hypertension (see *Warnings and Precautions* 0)
- Heart Failure and Edema (see *Warnings and Precautions* 0)
- Rash (see *Warnings and Precautions* 0)
- Anaphylactic Reactions (see *Warnings and Precautions* 0)
- Serious Skin Reactions (see *Warnings and Precautions* 0)
- Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS) (see *Warnings and Precautions* 0)
- Fetal Toxicity (see *Warnings and Precautions* 0)
- Hematologic Toxicity (see *Warnings and Precautions* 0)

6.1 Clinical Trial Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Adults

Osteoarthritis and Rheumatoid Arthritis

Meloxicam 7.5 mg/day controlled studies include 10,722 OA patients and 1012 RA patients treated with Meloxicam 7.5 mg/day, 3505 OA patients and 1351 RA patients treated with Meloxicam 15 mg/day. Meloxicam at these doses was administered to 661 patients for at least 6 months and to 312 patients for at least one year. Approximately 10,300 patients were included in the controlled osteoarthritis trials and 2363 of these patients were treated in 10 placebo- and/or active-controlled rheumatoid arthritis trials. Gastrointestinal (GI) adverse events were the most frequently reported adverse events in all treatment groups across Meloxicam trials.

A 12-week multicenter, double-blind, randomized trial was conducted in patients with osteoarthritis of the knee or hip to compare the efficacy and safety of Meloxicam with placebo and active-controlled agents. Two 12-week multicenter, double-blind, randomized trials were conducted in patients with rheumatoid arthritis to compare the efficacy and safety of Meloxicam with placebo.

Table 1a depicts adverse events that occurred in $\geq 2\%$ of the Meloxicam treatment groups in a 12-week placebo- and active-controlled osteoarthritis trial.

Table 1b depicts adverse events that occurred in $\geq 2\%$ of the Meloxicam treatment groups in two 12-week placebo-controlled rheumatoid arthritis trials.

Table 1a Adverse Events (%) Occurring in $\geq 2\%$ of Meloxicam Patients in a 12-Week Osteoarthritis Placebo- and Active-Controlled Trial

	Placebo 7.5 mg daily	Meloxicam 7.5 mg daily	Meloxicam 15 mg daily	Diclofenac 100 mg daily
No. of Patients	157	154	156	153
Gastrointestinal	17.2	20.1	17.3	28.1
Abdominal pain	2.5	1.9	2.6	1.3
Diarrhea	2.8	2.6	2.2	0.2
Dyspepsia	4.5	4.5	4.5	6.5
Flatulence	4.5	3.2	3.2	3.9
Nausea	3.2	3.9	3.8	7.2

Body as a Whole					
Accident household	1.9	4.5	3.2	2.6	
Edema	2.5	1.9	4.5	3.3	
Fatigue	0.6	2.6	0.9	3.3	
Influenza-like symptoms	9.1	4.5	5.8	2.6	
Central and Peripheral Nervous System					
Dizziness	3.2	2.6	3.6	2.0	
Headache	10.2	7.8	8.3	5.9	
Respiratory					
Pharyngitis	1.3	0.6	3.2	1.3	
Upper respiratory tract infection	1.9	3.2	1.9	3.3	
Skin					
Rash ^a	2.5	2.6	0.6	2.0	

Table 2b Adverse Events (%) Occurring in $\geq 2\%$ of Meloxicam Patients in two 12-Week Rheumatoid Arthritis Placebo- Controlled Trials

No. of Patients	Placebo	Meloxicam 7.5 mg daily	Meloxicam 15 mg daily
Gastrointestinal Disorders	14.1	18.9	16.8
Abdominal pain NOS	0.5	2.9	2.3
Dyspeptic signs and symptoms ^b	1.3	3.5	4.0
Nausea	2.6	3.3	3.8
General Disorders and Administration Site Conditions			
Influenza-like illness ^b	2.1	2.9	2.3
Infectious and Parasitic Conditions			
Upper Respiratory tract infections-pathogen class unspecified ^b	4.1	7.0	6.5
Musculoskeletal and Connective Tissue Disorders			
Joint and muscle signs and symptoms ^b	1.9	1.5	2.3
Nervous System Disorders			
Headaches NOS ^b	6.4	6.4	5.5
Skin and Subcutaneous Tissue Disorders			
Pruritus	1.7	1.6	2.1

^a MedDRA preferred term; ^b nausea, abdominal pain NOS, influenza-like illness, headaches NOS, and rash NOS

^b MedDRA high level term (preferred term), dyspeptic signs and symptoms (dyspepsia, dysuria, aggravated, eructation, pain), influenza-like illness (influenza, influenza-like signs and symptoms), pruritus (itching), rash (erythema, rash), joint related signs and symptoms (arthralgia, arthralgia aggravated, joint crepitus, joint effusion, joint swelling)

The adverse events that occurred with Meloxicam in $\geq 2\%$ of patients treated short-term (4 to 6 weeks) and long-term (6 months) in active-controlled osteoarthritis trials are presented in Table 2.

Table 3 Adverse Events (%) Occurring in $\geq 2\%$ of Meloxicam Patients in 4 to 6 Weeks and 6 Month Active-Controlled Osteoarthritis Trials

No. of Patients	4-6 Weeks Controlled Trials		6 Month Controlled Trials	
	Meloxicam 7.5 mg daily	Meloxicam 15 mg daily	Meloxicam 7.5 mg daily	Meloxicam 15 mg daily
Gastrointestinal				
Abdominal pain	11.8	18.0	26.6	24.2
Constipation	2.7	2.3	4.7	2.9
Diarrhea	0.8	1.2	1.8	2.6
Dyspepsia	1.9	2.7	5.9	2.6
Flatulence	0.5	0.4	4.0	2.6
Headache	7.4	4.7	4.7	7.2
Vomiting	0.6	0.8	1.8	2.6
Body as a Whole				
Accident household	0.0	0.0	0.6	2.9
Edema	0.6	2.0	2.4	1.6
Pain	0.9	2.0	3.6	5.2
Central and Peripheral Nervous System				
Dizziness	1.1	1.6	2.4	2.6
Headache	2.4	2.7	3.6	2.6
Hematologic				
Anemia	0.1	0.0	4.1	2.9
Musculoskeletal				
Arthralgia	0.5	0.0	5.3	1.3
Back pain	0.5	0.4	3.0	0.7
Psychiatric				
Insomnia	0.4	0.0	3.6	1.6
Respiratory				
Cough	0.3	0.8	2.4	1.0
Upper respiratory tract infection	0.2	0.0	8.3	7.5
Skin				
Pruritus	0.4	1.2	2.4	0.0
Rash	0.5	1.2	3.0	1.3
Urinary				
Micturition frequency	0.1	0.4	2.4	1.3
Urinary tract infection	0.3	0.4	4.7	6.9

^a WHO preferred terms rash, rash erythematous, and rash maculopapular combined

^b Increased risk of serious GI events; therefore, the daily dose of Meloxicam should not exceed 15 mg.

Pediatrics

Pauciarticular and Polyarticular Course Juvenile Rheumatoid Arthritis (JRA)

Three hundred and eighty-seven patients with pauciarticular and polyarticular course JRA were exposed to Meloxicam with doses ranging from 0.125 to 0.375 mg/kg per day in three controlled trials. These studies consisted of one 12-week controlled trial and two randomized trials (one with a 12-week open-label extension and one with a 40-week extension) and one 1-year open-label PK study. The adverse events observed in these pediatric studies with Meloxicam were similar in nature to the adult clinical trial experience, although the frequency of adverse events was lower. The following most common adverse events, abdominal pain, vomiting, diarrhea, headache, and pyrexia, were more common in the pediatric than in the adult trials. Rash was reported in a similar frequency in the pediatric and adult trials. No unexpected adverse events were identified during the course of the trials. The adverse events did not demonstrate an age or gender-specific subgroup effect.

The following is a list of adverse drug reactions occurring in $> 2\%$ of patients receiving Meloxicam in clinical trials involving approximately 16,200 patients.

Body as a Whole	allergic reaction, edema, fatigue, fever, hot flushes, malaise, syncope, weight decrease, weight increase
Cardiovascular	angina pectoris, cardiac failure, hypertension, hypotension, myocardial infarction, vasculitis
Central and Peripheral Nervous System	convulsions, paresis, paresthesia, vertigo
Gastrointestinal	colitis, dry mouth, duodenal ulcer, eructation, esophagitis, gastritis, gastritis/gastroesophageal reflux, gastrointestinal hemorrhage, hematemesis, hemorrhagic duodenal ulcer, hemorrhagic gastric ulcer, intestinal perforation, melena, pancreatitis, perforated duodenal ulcer, perforated gastric ulcer, stomatitis ulcerative
Heart Rate and Rhythm	tachycardia
Hematologic	leukopenia, purpura, thrombocytopenia
Liver and Biliary System	ALT increased, AST increased, bilirubinemia, GGT increased, hepatitis
Metabolic and Nutritional	dehydration
Psychiatric	delirium, drowsiness, anxiety, appetite increased, confusion, depression, nervousness, somnolence
Respiratory	asthma, bronchospasm, dyspnea
Skin and Appendages	alopecia, angioedema, bullous eruption, photosensitivity reaction, pruritus, sweating increased, urticaria
Special Senses	abnormal vision, conjunctivitis, taste perversion, tinnitus
Urinary System	albuminuria, BUN increased, creatinine increased, hematuria, renal failure

6.2 Post Marketing Experience

The following adverse reactions have been identified during post-approval use of Meloxicam. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship. The following reactions have been selected to illustrate the types of adverse events that are being monitored.

Serotonin release by platelets plays an important role in hemostasis. Case reports and postmarketing studies showed that concomitant use of drugs that interfere with serotonin reuptake and an NSAID may potentiate the risk of bleeding more than an NSAID alone.

Intervention: Adverse events with concomitant use of Meloxicam with anti-coagulants (e.g., warfarin), antiplatelet agents (e.g., aspirin), selective serotonin reuptake inhibitors (SSRIs), and serotonin reuptake inhibitors (e.g., SNRIs) for signs of bleeding [see Warnings and Precautions (5.2)].

Aspirin: Controlled clinical studies showed that the concomitant use of NSAIDs and analgesic doses of aspirin does not produce any greater therapeutic effect than the use of NSAIDs alone. In a clinical study, the concomitant use of an NSAID and aspirin was associated with a significant increase of the risk of serious GI adverse events compared to use of the NSAID alone [see Warnings and Precautions (5.2)].

Intervention: Concomitant use of Meloxicam and low dose aspirin or analgesic doses of aspirin does not generally recommend due to the increased risk of bleeding [see Warnings and Precautions (5.2)]. Meloxicam is not a substitute for low dose aspirin for cardiovascular protection.

ACE Inhibitors, Angiotensin Receptor Blockers, or Beta-Blockers: NSAIDs may diminish the antihypertensive effect of angiotensin converting enzyme (ACE) inhibitors, angiotensin receptor blockers (ARBs), or beta-blockers (including propranolol).

In patients who are elderly, volume-depleted, or have impaired renal function, more prominent signs of worsening renal function [see Warnings and Precautions (5.2)].

When these drugs are administered concomitantly, patients should be monitored for signs of acute renal failure. Assess function at the beginning of the concomitant treatment and periodically thereafter.

Durkets: Clinical studies, as well as post-marketing observations, showed that NSAIDs reduced the natriuretic effect of loop diuretics (e.g.,

frusemide, torsemide, and bumetanide).

During concurrent use of Meloxicam and ACE inhibitors, ARBs, or loop diuretics, the blood pressure should be monitored to ensure that the desired blood pressure is obtained.

During concomitant use of Meloxicam and ACE inhibitors or ARBs in patients who are elderly, volume-depleted, or have impaired renal function, more prominent signs of worsening renal function [see Warnings and Precautions (5.2)].

	<p>furosemide) and thiazide diuretics in some patients. This effect has been attributed to inhibition of prostaglandin synthesis. However, studies with furosemide and mebxicam have not demonstrated a reduction in natriuretic effect. Furosemide single and multiple dose pharmacodynamics and pharmacokinetics are not affected by the use of mebxicam.</p>
Intervention:	During concurrent use of Mebxicam and diuretics, observe patients for signs of worsening renal function, in addition to assuring diuretic efficacy including antihypertensive effects (see Warnings and Precautions (5)).
Lithium	
Clinical Impact:	NSAIDs have produced elevations in plasma lithium levels and reductions in renal lithium clearance. The mean increase in lithium concentrations is up to 15%, and the renal clearance is reduced by approximately 20%. This effect has been attributed to NSAID inhibition of renal prostaglandin synthesis (see Clinical Pharmacology (12.3)).
Intervention:	During concurrent use of Mebxicam and lithium, monitor patients for signs of lithium toxicity.
Methotrexate	
Clinical Impact:	Concomitant use of NSAIDs and methotrexate may increase the risk for methotrexate toxicity (e.g., neutropenia, thrombocytopenia, renal dysfunction).
Intervention:	During concurrent use of Mebxicam and methotrexate, monitor patients for methotrexate toxicity.
Cyclosporine	
Clinical Impact:	Concomitant use of Mebxicam and cyclosporine may increase cyclosporine levels.
Intervention:	During concurrent use of Mebxicam and cyclosporine, monitor patients for signs of worsening renal function.
NSAIDs and Salicylates	
Clinical Impact:	Concomitant use of mebxicam with other NSAIDs or salicylates (e.g., aspirin, salsalate) increases the risk of GI toxicity, with little or no increase in efficacy (see Warnings and Precautions (5.2)).
Intervention:	The concomitant use of mebxicam with other NSAIDs or salicylates is not recommended.
Pemetrexed	
Clinical Impact:	Concomitant use of Mebxicam and pemetrexed may increase the risk of pemetrexed-induced nephrotoxicity, rash, and GI toxicity (see Warnings and Precautions (5.1) and Dosage and Administration).
Intervention:	During concurrent use of Mebxicam and pemetrexed, in patients with renal impairment whose creatinine clearance ranges from 45 to 79 mL/min, the dose of pemetrexed should be reduced by 50% (see Dosage and Administration). Patients taking mebxicam should interrupt dosing for at least five days before, the day of, and two days following pemetrexed administration. In patients with creatinine clearance below 45 mL/min, the concomitant administration of mebxicam with pemetrexed is not recommended.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Use of NSAIDs, including Mebxicam, can cause premature closure of the fetal ductus arteriosus and fetal renal impairment, including oligohydramnios, and, in some cases, neonatal renal impairment. Because of these risks, limit dose and duration of Mebxicam use between about 20 and 30 weeks of gestation, and avoid Mebxicam use at about 30 weeks of gestation and later in pregnancy (see Clinical Considerations, Data).

Premature Closure of Fetal Ductus Arteriosus

Use of NSAIDs, including Mebxicam, at about 30 weeks gestation or later in pregnancy increases the risk of premature closure of the fetal ductus arteriosus.

Oligohydramnios/Neonatal Renal Impairment

Use of NSAIDs at about 20 weeks gestation or later in pregnancy has been associated with cases of fetal renal dysfunction leading to oligohydramnios, and, in some cases, neonatal renal impairment.

Data from observational studies regarding potential embryofetal risks of NSAID use in women in the first or second trimesters of pregnancy are inconclusive.

In animal reproduction studies, embryofetal death was observed in rats and rabbits treated during the period of organogenesis with mebxicam at oral doses equivalent to 0.65 and 5.5 times the MRHD, respectively. No teratogenic effects were observed in rats. Increased incidence of septal heart defects were observed in rabbits treated throughout embryogenesis with mebxicam at an oral dose equivalent to 78-times the MRHD. In pre- and post-implantation studies, mebxicam was an oral dose equivalent to 26-times the MRHD, decreased parturition, and decreased offspring survival at 0.08-times MRHD of mebxicam. No teratogenic effects were observed in rats and rabbits treated with mebxicam during organogenesis at an oral dose equivalent to 2.6 and 26-times the MRHD (see Data).

Based on animal data, pregnancies have been shown to have an increased risk in embryo viability, pre-implantation, and post-implantation, decapsulation. In animal studies, administration of prostaglandin synthesis inhibitors, such as mebxicam, resulted in increased pre- and post-implantation loss. Prostaglandins also have been shown to be important for normal parturition. In published animal studies, prostaglandin synthesis inhibitors have been reported to impair kidney development when administered at clinically relevant doses.

The estimated background risk of major birth defects and miscarriage for the general population is approximately 3% to 4% for birth defects and 15% to 20% for miscarriage. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

Clinical Considerations

- Fetal/Neonatal Adverse Reactions

Premature Closure of Fetal Ductus Arteriosus:

Avoid use of NSAIDs in women at about 30 weeks gestation and later in pregnancy, because of risks, including mebxicam, can cause premature closure of the fetal ductus arteriosus (see Data).

Oligohydramnios/Neonatal Renal Impairment:

If an NSAID is necessary at about 20 weeks gestation or later in pregnancy, limit the use to the lowest effective dose and shortest duration possible. If mebxicam treatment extends beyond 48 hours, consider monitoring with ultrasound for oligohydramnios. If oligohydramnios occurs, discontinue mebxicam and follow up according to clinical practice (see Data).

Labor or Delivery

There are no studies on the effects of Mebxicam during labor or delivery. In animal studies, NSAIDs, including mebxicam, inhibit prostaglandin synthesis, cause delayed parturition, and increase the incidence of stillbirth.

Data

Human Data

Premature Closure of Fetal Ductus Arteriosus:

Published literature reports that the use of NSAIDs at about 30 weeks of gestation and later in pregnancy may cause premature closure of the fetal ductus arteriosus (see Data).

Oligohydramnios/Neonatal Renal Impairment:

Published studies and postmarketing reports describe maternal NSAID use at about 20 weeks of gestation and later in pregnancy with fetal renal dysfunction leading to oligohydramnios, and in some cases, neonatal renal impairment. These adverse outcomes are seen, on average, after days to weeks of treatment, although oligohydramnios has been reported as early as 48 hours after NSAID initiation. In many cases, but not all, the decrease in amniotic fluid was transient and reversible with cessation of the drug. There have been a limited number of case reports of maternal NSAID use and neonatal renal dysfunction without oligohydramnios, some of which involved oligohydramnios that required treatment with invasive procedures, such as exchange transfusion or dialysis.

Methodological limitations of these postmarketing studies and reports include lack of a control group; limited information regarding dose, duration, and timing of drug exposure; and lack of information on the timing of drug initiation, which precludes establishing a reliable estimate of the risk of adverse fetal and neonatal outcomes with maternal NSAID use. Because the published safety data on neonatal outcomes involved mostly term infants, the generalizability of certain reported risks to the full-term infant exposed to NSAIDs through maternal use is uncertain.

Animal Data

Mebxicam was not teratogenic when administered to pregnant rats during fetal organogenesis at oral doses up to 4 mg/kg/day (2.6-fold greater than the MRHD of 15 mg of Mebxicam based on BSA comparison). Administration of mebxicam to pregnant rabbits at oral doses up to 10 mg/kg/day (7.5-fold greater than the MRHD based on BSA comparison) resulted in increased incidence of septal heart defects of the heart at an oral dose of 60 mg/kg/day (78-fold greater than the MRHD based on BSA comparison). The no effect level was 20 mg/kg/day (26-fold greater than the MRHD based on BSA comparison). Administration of mebxicam at oral doses of 1 mg/kg/day and 5 mg/kg/day, respectively (0.65 and 6.5-fold greater, respectively, than the MRHD based on BSA comparison) when administered through oral gavage.

Oral administration of mebxicam to pregnant rats, during late gestation through lactation increased the incidence of dystocia, delayed parturition, and decreased offspring survival at mebxicam doses of 0.125 mg/kg/day or greater (0.08-times MRHD based on BSA comparison).

8.2 Lactation

Risk Summary

There are no human data available on whether mebxicam is present in human milk, or on the effects on breastfed infants, or on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Mebxicam and any potential adverse effects on the breastfed infant from the Mebxicam or from the underlying maternal condition.

Data

Animal Data

Mebxicam was present in the milk of lactating rats at concentrations higher than those in plasma.

8.3 Females and Males of Reproductive Potential

Infertility

Based on the mechanism of action, the use of prostaglandin-mediated NSAIDs, including Mebxicam, may delay or prevent rupture of ovarian follicles, which has been associated with reversible infertility in some women. Published animal studies have shown that administration of NSAIDs, including mebxicam, to female rats during the pre-ovulatory prostaglandin-mediated follicular rupture required for ovulation. Small studies in women treated with NSAIDs have also shown a reversible delay in ovulation. Consider withdrawal of NSAIDs, including Mebxicam, in women who have difficulties conceiving or who are undergoing investigation of infertility.

8.4 Pediatric Use

The safety and effectiveness of mebxicam in pediatric JRA patients from 2 to 17 years of age has been evaluated in three clinical trials (see Dosage and Administration, Adverse Reactions (1) and Clinical Studies (1)).

8.5 Geriatric Use

Elderly patients, compared to younger patients, are at greater risk for NSAID-associated serious cardiovascular, gastrointestinal, and/or renal adverse reactions. If the anticipated benefit for the elderly patient outweighs these potential risks, start dosing at the lowest recommended dose and titrate slowly, monitor closely, and monitor patients for adverse effects [see Warnings and Precautions (5.2)].

8.6 Hepatic Impairment

No dose adjustment is necessary in patients with mild to moderate hepatic impairment. Patients with severe hepatic impairment have not been adequately studied. Meloxicam is significantly metabolized in the liver and hepatotoxicity may occur, use meloxicam with caution in patients with hepatic impairment [see Warnings and Precautions (5.1) and Clinical Pharmacology (12)].

8.7 Renal Impairment

No dose adjustment is necessary in patients with mild to moderate renal impairment. Patients with severe renal impairment have not been adequately studied. The use of meloxicam in subjects with severe renal impairment is not recommended. In patients on hemodialysis, meloxicam should not exceed 7.5 mg per day. Meloxicam is not dialyzable [see Dosage and Administration (1) and Clinical Pharmacology (12)].

10 OVERDOSE

Symptoms following acute NSAID overdoses have been typically limited to lethargy, drowsiness, nausea, vomiting, and epigastric pain, which have been generally reversible with supportive care. Gastrointestinal bleeding has occurred. Hypertension, acute renal failure, respiratory depression, and coma have occurred, but were rare [see Warnings and Precautions (5.2, 5.3)].

Manage patients with symptomatic and supportive care following an NSAID overdose. There are no specific antidotes. Consider emesis and/or activated charcoal (60 to 100 grams in adults, 1 to 2 grams per kg of body weight in pediatric patients) and/or osmotic cathartics. In symptomatic patients, seen within four hours of ingestion or in patients with a large overdose (5 to 10 times the recommended dosage), forced diuresis, alkalinization of urine, hemodialysis, or hemoperfusion may not be useful due to high protein binding.

There is limited experience with meloxicam overdoses. Cholestyramine is known to accelerate the clearance of meloxicam. Accelerated removal of meloxicam by 4 g oral doses of cholestyramine given three times a day was demonstrated in a clinical trial. Administration of cholestyramine may be useful following an overdose.

For additional information about overdosage treatment, call a poison control center (1-800-222-1222).

11 DESCRIPTION

Meloxicam tablets are a nonsteroidal anti-inflammatory drug (NSAID). Each tablet contains 7.5 mg or 15 mg meloxicam for oral administration. Meloxicam is chemically designated as 4-hydroxy-2-methyl-N-(5-methyl-2-thiazolyl)-2H-1,2-benzothiazine-3-carboxamide-1,1-dioxide. The molecular weight is 351.4. Its empirical formula is $C_{14}H_{19}N_2O_5S_2$ and it has the following structural formula:



Chemical Structure

Meloxicam is a pastel yellow solid, practically insoluble in water, with higher solubility observed in strong acids and bases. It is very slightly soluble in methanol. Meloxicam has an apparent partition coefficient (log Papp) = 0.1 in n-octanol/buffer pH 7.4. Meloxicam has a pKa of 1.1 and 4.2.

Meloxicam is available as a tablet for oral administration containing 7.5 mg or 15 mg meloxicam.

The inactive ingredients in Meloxicam tablets USP include colloidal silicon dioxide, crospovidone, lactose monohydrate, magnesium stearate, microcrystalline cellulose, povidone and sodium citrate dihydrate.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Meloxicam has analgesic, anti-inflammatory, and antipyretic properties. The mechanism of action of Meloxicam, like that of other NSAIDs, is not completely understood but involves inhibition of cyclooxygenase (COX-1 and COX-2). Meloxicam is a potent inhibitor of prostaglandin synthesis in vitro. Meloxicam concentrations required to inhibit prostaglandin synthesis are similar to those concentrations that sensitize afferent nerves and potentiate the action of bradykinin in inducing pain in animal models. Prostaglandins are mediators of inflammation. Because meloxicam is an inhibitor of prostaglandin synthesis, its mode of action may be due to a decrease of prostaglandins in peripheral tissues.

12.3 Pharmacokinetics

Absorption

The absolute bioavailability of meloxicam capsules was 89% following a single oral dose of 30 mg compared with 30 mg IV bolus injection. Following a single oral meloxicam dose, dose-proportional pharmacokinetics were shown in the range of 5 mg to 60 mg. After multiple oral doses the pharmacokinetics of meloxicam capsules were dose-proportional over the range of 7.5 mg to 15 mg. Mean Cmax was achieved within four to five hours after oral administration. The pharmacokinetics of meloxicam capsules were dose-prolonged drug absorption. With multiple dosing, steady-state concentrations were reached by Day 5. A second meloxicam concentration peak occurs around 12 to 14 hours post-dose, suggesting biliary recycling.

Meloxicam capsules have been shown to be bioequivalent to Meloxicam tablets.

Table 4 Single Dose and Steady-State Pharmacokinetic Parameters for Oral 7.5 mg and 15 mg Meloxicam (Mean and % CV)*.

Pharmacokinetic Parameters (%CV)	Steady State		Single Dose		
	7.5 mg tablets	15 mg tablets	15 mg capsules	15 mg capsules	15 mg capsules
N			12		
C _{max}	[μ g/ml]	1.05 (20)	2.3 (59)	3.2 (24)	0.84 (29)
T _{max}	[h]	4.9 (8)	5 (12)	6 (27)	4 (65)
T _{1/2}	[h]	20.1 (29)	21 (34)	24 (34)	16 (29)
CL _{app}	[mL/min]	8.8 (29)	9.9 (76)	5.1 (22)	19 (43)
V _d	[L]	14.7 (32)	15 (42)	10 (50)	26 (44)
V _d /F					14 (29)

* The parameter values in the table are from various studies.

† Not under high fat conditions.

‡ Mean \pm SD.

* V_d/F = Dose/(AUC_{0-t}/Ke_t)

Food and Antacid Effects

Administration of meloxicam capsules following a high fat breakfast (75 g of fat) resulted in mean peak drug levels (i.e., C_{max}) being increased by approximately 22% while the extent of absorption (AUC) was unchanged. The time to maximum concentration (T_{max}) was increased by approximately 1.5 hours. Meloxicam capsules may be administered with concomitant administration of antacids. Based on these results, Meloxicam can be administered without regard to timing of meals or concomitant administration of antacids.

Distribution

The mean volume of distribution (V_d) of meloxicam is approximately 10 L. Meloxicam is 99% bound to human plasma proteins (primarily albumin) within the therapeutic dose range. The fraction of protein binding is independent of drug concentration, over the clinically relevant concentration range, but decreases to ~99% in patients with renal disease. The fraction of drug bound to albumin is approximately 99% at concentrations less than 10%. Following a radiolabeled dose, over 90% of the radioactivity detected in the plasma was present as unchanged meloxicam.

Meloxicam concentrations in synovial fluid, after a single oral dose, range from 40% to 50% of those in plasma. The free fraction in synovial fluid is 2.5 times higher than in plasma due to the lower albumin content in synovial fluid as compared to plasma. The significance of this penetration is unknown.

Elimination

Metabolism

Meloxicam is extensively metabolized in the liver. Meloxicam metabolites include 5'-carboxy meloxicam (60% of dose), from P-450 mediated metabolism formed by oxidation of the 5'-hydroxymethyl group. The 5'-carboxy meloxicam is further excreted to a lesser extent (9% of dose). In vitro studies indicate that CYP2C9 (cytochrome P450 2C9) metabolizing enzyme plays an important role in this metabolic pathway with a minor contribution of the CYP3A4 isozyme. Patients' peroxidase activity is proportional to the sum of the other two metabolites which account for 18% and 4% of the administered dose, respectively. All the four metabolites are not known to have any *in vivo* pharmacological activity.

Excretion

Meloxicam excretion is predominantly in the form of metabolites, and occurs to equal extent in urine and feces. The mean total body clearance of meloxicam is 10.1 L/h. Meloxicam is excreted in the urine (0.2%) and feces (1.6%). The extent of the urinary excretion was confirmed for unlabeled multiple 7.5 mg doses. 0.5%, 6%, and 13% of the dose were found in urine in the form of meloxicam, and the 5'-hydroxymethyl and 5'-carboxy meloxicam. The remaining 93% of the dose was excreted in the feces as the total secreted of the drug. This was demonstrated when oral administration of cholestyramine following a single IV dose of meloxicam decreased the AUC of meloxicam by 50%.

The mean elimination half-life (T_{1/2}) ranges from 15 hours to 20 hours. The elimination half-life is dose independent across dose levels indicating linear metabolism within the therapeutic dose range. Plasma clearance ranges from 7 to 9 mL/min.

Specific Population

Pediatric

After single (0.25 mg/kg) dose administration and after achieving steady state (0.375 mg/kg/day), there was a general trend of approximately 30% lower exposure in younger patients (2 to 6 years old) as compared to the older patients (7 to 16 years old). The older patients had approximately 30% higher exposure (AUC) than the younger patients (at steady state) to those in the adult patients, when using AUC values normalized to a dose of 0.25 mg/kg [see Dosage and Administration (1)]. The meloxicam mean (SD) elimination half-life was 15.2 (10.1) to 13.0 hours (3.0) for the 2 to 6 year old patients, and 7 to 15 hours for the older patients.

In a covariate analysis, utilizing population pharmacokinetics body weight, but not age, was the single predictive covariate for differences in the meloxicam apparent oral plasma clearance. The body-weight normalized apparent oral clearance values were adequate predictors of meloxicam exposure in pediatric patients.

The pharmacokinetics of Meloxicam in pediatric patients under 2 years of age have not been investigated.

Geriatric

Elderly males (≥65 years of age) exhibited meloxicam plasma concentrations and steady-state pharmacokinetics similar to those of young males. Elderly females (≥65 years of age) had a 47% higher AUC_{ss} and 22% higher Cmax_{ss} as compared to young females (≤55 years of age) after body weight normalization. Despite the increased total concentrations in the elderly females, the adverse event profile was comparable for both elderly patient populations. A smaller free fraction was found in elderly female patients in comparison to elderly male patients.

Sex

Young females exhibited slightly lower plasma concentrations relative to young males. After single doses of 7.5 mg Meloxicam, the mean elimination half-life was 19.5 hours for the female group as compared to 23.4 hours for the male group. At steady state, the total AUC_{ss} was 22% higher in females as compared to males. This difference due to gender is likely to be of little clinical importance. There was linearity of pharmacokinetics and no appreciable difference in the Cmax or Tmax across genders.

Hepatic Impairment

Following a single 15 mg dose of meloxicam there was no marked difference in plasma concentrations in healthy volunteers with mild (Child-Pugh Class II) or moderate (Child-Pugh Class III) hepatic impairment compared to healthy volunteers. Protein binding of meloxicam was not affected by hepatic impairment. No dosage adjustment is necessary in patients with mild to moderate hepatic impairment. Patients with severe hepatic impairment (Child-Pugh Class IV) have not been adequately studied (see Warnings and Precautions () and Use in Specific Populations ()).

Renal Impairment

Meloxicam pharmacokinetics have been investigated in subjects with mild and moderate renal impairment. Total drug plasma concentrations of meloxicam decreased and total clearance of meloxicam increased with the degree of renal impairment, while free AUC values were similar. The decrease in total drug plasma concentrations with moderate renal impairment may be due to increased fraction of unbound meloxicam which is available for hepatic metabolism and subsequent excretion. No dosage adjustment is necessary in patients with mild to moderate renal impairment. Patients with severe renal impairment have not been adequately studied. The use of Meloxicam in subjects with severe renal impairment is not recommended (see Dosage and Administration (), Warnings and Precautions () and Use in Specific Populations ()).

Hemodialysis

Following a single dose of meloxicam, the free Cmax_{ss} concentrations were higher in patients receiving hemodialysis (1% free fraction) in comparison to healthy volunteers (0.3% free fraction). Hemodialysis did not lower the total drug concentration in plasma; therefore, additional doses are not necessary after hemodialysis. Meloxicam is not dialyzable (see Dosage and Administration () and Use in Specific Populations ()).

Drug Interaction Studies

Aspirin: When NSAIDs were administered with aspirin, the protein binding of NSAIDs were reduced, although the clearance of free NSAID was not altered. When Meloxicam is administered with aspirin (1000 mg) four times daily to healthy volunteers, there was no effect on the AUC (10%) or Cmax (24%) of meloxicam. The clinical significance of this interaction is not known. See Table 3 for clinically significant drug interactions of NSAIDs with aspirin (see Drug Interactions ()).

Cholestyramine: Pretreatment for 7 days with cholestyramine significantly increased the AUC of meloxicam by 22%. This resulted in a decrease in t_{1/2} from 19.2 hours to 12.5 hours, and a 35% reduction in AUC. This suggests the existence of a recirculation pathway for meloxicam in the gastrointestinal tract. The clinical relevance of this interaction is not known.

Cimetidine: Concomitant administration of 200 mg cimetidine four times daily did not alter the single-dose pharmacokinetics of 30 mg meloxicam.

Doxivir: Meloxicam 15 mg once daily for 7 days did not alter the plasma concentration profile of doxivir after β-acetyldoxivir administration for 7 days at clinical doses. *In vitro* testing found a protein binding drug interaction between doxivir and meloxicam.

Lithium: In a study conducted in healthy subjects, mean pre-dose lithium concentration and AUC were increased in subjects receiving lithium doses ranging from 804 to 1072 mg twice daily with meloxicam 15 mg QD every day as compared to subjects receiving lithium alone (see Drug Interactions ()).

Methotrexate: A study in 13 rheumatoid subjects (RA) patients evaluated the effects of multiple doses of meloxicam on the pharmacokinetics of methotrexate taken once weekly. Meloxicam did not have a significant effect on the pharmacokinetics of single doses of methotrexate. *In vitro*, methotrexate did not displace meloxicam from ³H human serum binding sites (see Drug Interactions ()).

Warfarin: The concomitant effect of warfarin was studied in 9 pairs of healthy subjects receiving 4 doses of warfarin that produced an INR (International Normalized Ratio) between 1.2 and 1.8. In these subjects, meloxicam did not alter warfarin pharmacokinetics and the average anticoagulant effect of warfarin as determined by INR was not changed. However, one subject showed an increased INR from 1.5 to 2.1. Caution should be used when administering Meloxicam with warfarin since patients on warfarin may experience changes in INR and an increased risk of bleeding complications when a new medication is introduced (see Drug Interactions ()).

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

There was no increase in tumor incidence in long-term carcinogenicity studies in rats (104 weeks) and mice (91 weeks) administered meloxicam at oral doses up to 0.8 mg/kg/day, 1.6 mg/kg/day, and up to 8.0 mg/kg/day (up to 0.5- and 2.6-times, respectively; the maximum recommended human dose (MRHD) of 15 mg/day Meloxicam based on body surface area [BSA] comparison).

Mutagenesis

Meloxicam was not mutagenic in an Ames assay, or clastogenic in a chromosome aberration assay with human lymphocytes and an *in vivo* micronucleus test in mouse bone marrow.

Impairment of Fertility

Meloxicam did not impair male and female fertility in rats at oral doses up to 9 mg/kg/day in males and 5 mg/kg/day in females (up to 5.8- and 3.2-times greater, respectively, than the MRHD based on BSA comparison).

14 CLINICAL STUDIES

14.1 Osteoarthritis and Rheumatoid Arthritis

The use of Meloxicam for the treatment of the signs and symptoms of osteoarthritis of the knee and hip was evaluated in a 12-week, double-blind, controlled trial. Meloxicam (3.75 mg, 7.5 mg, and 15 mg daily) was compared to placebo. The four primary endpoints were: pain (assessed by a self-administered questionnaire), patient pain assessment, and total WOMAC score (a self-administered questionnaire addressing pain, function, and stiffness). Patients on Meloxicam 7.5 mg daily and Meloxicam 15 mg daily showed significant improvement in each of these endpoints compared with placebo.

The use of Meloxicam for the management of signs and symptoms of osteoarthritis was evaluated in six double-blind, active-controlled trials outside the U.S. ranging from 4 weeks' to 6 months' duration. In these trials, the efficacy of Meloxicam, in doses of 7.5 mg/day and 15 mg/day, was compared to ibuprofen, naproxen, and diclofenac SR 100 mg/day and consistent with the efficacy seen in the U.S. trial.

The use of Meloxicam for the treatment of the signs and symptoms of rheumatoid arthritis was evaluated in a 12-week, double-blind, controlled trial. Meloxicam (7.5 mg, 15 mg, and 22.5 mg daily) was compared to placebo. The primary endpoint in this study was the ACR20 response rate, a composite measure of clinical, laboratory, and functional measures of RA response. Patients receiving Meloxicam 7.5 mg and 15 mg daily showed significant improvement in the primary endpoint compared with placebo. An incremental benefit was observed with the 22.5 mg dose compared to the 15 mg dose.

14.2 Juvenile Rheumatoid Arthritis (JRA) Pauciarticular and Polyarticular Course

The use of Meloxicam for the treatment of the signs and symptoms of pauciarticular or polyarticular course Juvenile Rheumatoid Arthritis in patients 2 years of age and older was evaluated in two 12-week, double-blind, parallel-arm, active-controlled trials.

Both studies included three arms: naproxen and two doses of meloxicam. In both studies, meloxicam dosing began at 0.125 mg/kg/day (7.5 mg maximum) or 0.25 mg/kg/day (15 mg maximum), and naproxen dosing began at 10 mg/kg/day. One study used these doses throughout the 12-week dosing period, while the other incorporated a dose increase after 6 weeks to 0.25 mg/kg/day (7.5 mg) or 0.375 mg/kg/day (22.5 mg maximum) of meloxicam and 15 mg/kg/day of naproxen.

The efficacy analysis used the ACR Pediatric 30 responder definition, a composite of parent and investigator assessments, counts of active joints and joints with limited range of motion, and erythrocyte sedimentation rate. The response rate was similar in all three groups in both studies, and no difference was observed between the meloxicam dose groups.

16 HOW SUPPLIED/STORAGE AND HANDLING

Meloxicam tablets USP are available as a light yellow, round, flat, uncoated tablet containing meloxicam 7.5 mg, as light yellow, oblong, biconvex, uncoated tablet, containing meloxicam 15 mg. The 7.5 mg tablet is imprinted with letter U and L on one side and tablet code 7.5 on the other side. The 15 mg tablet is imprinted with letter U and L on one side and tablet code 15 on the other side.

Meloxicam Tablets USP 7.5 mg are available as follows:

NDC 68788-8758-1 Bottles of 15

NDC 68788-8758-2 Bottles of 20

NDC 68788-8758-8 Bottles of 28

NDC 68788-8758-3 Bottles of 30

NDC 68788-8758-6 Bottles of 60

NDC 68788-8758-9 Bottles of 90

NDC 68788-8758-0 Bottles of 105 (Storage: Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Keep Meloxicam Tablets USP in a dry place. Dispense tablets in a tight container.

Keep this and all medications out of the reach of children.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide) that accompanies each prescription dispensed.

Additional Medication Guides can be obtained by calling Unichem at 1-866-562-4616.

Inform patients, families or their caregivers of the following information before initiating therapy with an NSAID and periodically during the course of ongoing therapy.

Cardiovascular/Thrombotic Events

Advise patients to be alert for the symptoms of cardiovascular thrombotic events, including chest pain, shortness of breath, weakness, or slurring of speech, and to report any of these symptoms to their healthcare provider immediately (see Warnings and Precautions ()).

Gastrointestinal Bleeding, Ulceration, and Perforation

Advise patients to report symptoms of ulcerations and bleeding, including epigastric pain, dyspepsia, melena, and hematemesis to their healthcare provider. In the setting of concomitant use of low-dose aspirin for cardiac prophylaxis, inform patients of the increased risk of the signs and symptoms of GI bleeding [see Warnings and Precautions (5.2)].

• **Hepatotoxicity:**

Inform patients of the warning signs and symptoms of hepatotoxicity (e.g., nausea, fatigue, diarrhea, pruritus, jaundice, right upper quadrant tenderness), and "flu-like" symptoms. If these occur, instruct patients to stop Meloxicam tablets and seek immediate medical therapy [see Warnings and Precautions (5)].

• **Heart Failure and Edema:**

Advise patients to be alert for the symptoms of congestive heart failure including shortness of breath, especially at night, pain, or edema and to contact their healthcare provider if such symptoms occur [see Warnings and Precautions (5)].

• **Anaphylactic Reactions:**

Inform patients of the signs of an anaphylactic reaction (e.g., difficulty breathing, swelling of the face or throat). Instruct patients to seek immediate emergency help if these occur [see Contraindications (4) and Warnings and Precautions (5)].

• **Serious Skin Reactions, including DRESS:**

Advise patients to stop taking Meloxicam tablets immediately if they develop any type of rash or fever, and to contact their healthcare provider as soon as possible [see Warnings and Precautions (5)].

• **Female Fertility:**

Advise females of reproductive potential who desire pregnancy that NSAIDs, including Meloxicam tablets, may be associated with a reversible delay in ovulation [see Use in Specific Populations (8.1)].

• **Fetal Toxicity:**

Inform pregnant women to avoid use of Meloxicam tablets and other NSAIDs starting at 20 weeks of gestation because of the risk of the premature closing of the fetal ductus arteriosus. If treatment with Meloxicam tablets is needed for a pregnant woman between about 20 to 30 weeks gestation, advise her that she may need to be monitored for oligohydramnios, if treatment continues for longer than 48 hours [see Warnings and Precautions (5) and Use in Specific Populations (8.1)].

• **Avoid Concomitant Use of NSAIDs:**

Inform patients that the concomitant use of Meloxicam tablets with other NSAIDs or salicylates (e.g., diflunisal, salsalate) is not recommended due to the increased risk of gastrointestinal toxicity, and little or no increase in efficacy [see Warnings and Precautions (5.1) and Drug Interactions (7)]. Alert patients that NSAIDs may be present in "over the counter" medications for treatment of colds, fever, or insomnia.

• **Use of NSAIDs and Low-Dose Aspirin:**

Inform patients not to use low-dose aspirin concomitantly with Meloxicam tablets until they talk to their healthcare provider [see Drug Interactions (7)].

For current prescribing information, call Unichem at 1-866-562-4616.

Manufactured by:

UNICHEM LABORATORIES LTD.

Piplane Ind. Estate,

Piplane, Bardez, Goa 403511, India

Manufactured for:



East Brunswick, NJ 08816

12-07/2024

13015145

Repackaged By: Preferred Pharmaceuticals Inc.

SPL MEDGUIDE

Medication Guide for Nonsteroidal Anti-inflammatory Drugs (NSAIDs)

What is the most important information I should know about medicines called Nonsteroidal Anti-inflammatory Drugs (NSAIDs)?

NSAIDs can cause serious side effects, including:

- Increased risk of a heart attack or stroke that can lead to death. This risk may happen in patients who may already have a heart attack risk, or with increasing doses of NSAIDs.
- Do not take NSAIDs right before or after a heart surgery called a "coronary artery bypass graft (CABG)."
- Avoid taking NSAIDs after a recent heart attack unless your healthcare provider says it is okay. There is a higher risk of another heart attack if you take NSAIDs after a recent heart attack.
- Increased risk of bleeding, ulcers, and tears (perforation) of the esophagus (tube leading from the mouth to the stomach), stomach and intestines.
- anytime during use
- without warning symptoms
- that may cause death.

The risk of getting an ulcer or bleeding increases with:

- past history of stomach ulcers, or stomach or intestinal bleeding with use of NSAIDs
- taking other called "corticosteroids", "anticoagulants", "SSRIs", or "SNRIs"
- increasing doses of NSAIDs
- smoking
- drinking alcohol
- older age
- poor health
- advanced liver disease
- bleeding problems

NSAIDs should not be used:

- Do not use as prescribed
- at the lowest dose possible for your treatment
- for the shortest time needed

What are NSAIDs?

NSAIDs are used to treat pain and redness, swelling, and heat (inflammation) from medical conditions such as different types of arthritis, menstrual cramps, and other types of pain.

Who should not take NSAIDs?

Do not take NSAIDs:

- If you have had an asthma attack, hives, or other allergic reaction with aspirin or any other NSAID.
- right before or after heart bypass surgery.

Before taking NSAIDs, tell your healthcare provider about all of your medical conditions, including:

- have liver or kidney problems
- have high blood pressure
- have asthma
- are pregnant or plan to become pregnant. Taking NSAIDs at about 20 weeks of pregnancy or later may harm your unborn baby. If you need to take NSAIDs for more than 2 days when you are between 20 and 30 weeks of pregnancy, your healthcare provider should check the risks to your baby before you take NSAIDs.

You should not take NSAIDs after about 30 weeks of pregnancy.

- are breastfeeding or plan to breast feed

Tell your healthcare provider about all of the medicines you take, including prescription or over-the-counter medicines, vitamins or herbal supplements. NSAIDs and some other medicines can interact with each other and cause serious side effects. Do not start taking any new medicine without taking to your healthcare provider first.

What are the possible side effects of NSAIDs?

NSAIDs can cause serious side effects, including:

See the "What is the most important information I should know about medicines called Nonsteroidal Anti-inflammatory Drugs (NSAIDs)?"

new or worse high blood pressure

• heart problems

• liver problems including liver failure

• kidney problems including kidney failure

• low blood sodium levels

• life-threatening skin reactions

• life-threatening allergic reactions

• Other side effects of NSAIDs include: stomach pain, constipation, diarrhea, gas, heartburn, nausea, vomiting, and dizziness.

Get emergency help right away if you get any of the following symptoms:

- shortness of breath or trouble breathing
- chest pain
- weakness in one part or side of your body
- slurred speech
- swelling of the face or throat

Stop taking your NSAID and call your healthcare provider right away if you get any of the following symptoms:

- nausea
- more tired or weaker than usual
- diarrhea
- constipation
- your skin or eyes look yellow
- indigestion or stomach pain
- flu-like symptoms
- vomit blood
- there is blood in your bowel movement or it is black and sticky like tar
- unusual weight gain
- skin rash or blisters with fever
- swelling of the arms, legs, hands and feet

If you take too much of your NSAID, call your healthcare provider or get medical help right away.

These are not all the possible side effects of NSAIDs. For more information, ask your healthcare provider or pharmacist about NSAIDs.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Other information about NSAIDs:

• Aspirin is an NSAID but it does not increase the chance of a heart attack. Aspirin can cause bleeding in the brain, stomach, and intestines. Aspirin can also cause ulcers in the stomach and intestines.

• Some NSAIDs are sold in lower doses without a prescription (over-the-counter). Talk to your healthcare provider before using over-the-counter NSAIDs for more than 10 days.

General Information about the safe and effective use of NSAIDs

Medicines sometimes prescribed for purposes other than those listed in a Medication Guide. Do not take NSAIDs to treat symptoms not listed on your prescription. Do not give NSAIDs to other people, even if they have the same symptoms that you have. It may harm them.

If you want more information about NSAIDs, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about NSAIDs that is

written for health professionals.
Additional Medication Guides can be obtained by calling Unichem at 1-866-562-9116.
 The other trademarks referenced are owned by third parties not affiliated with Unichem Laboratories Limited
 Manufactured by:
UNICHEM LABORATORIES LTD.
 Plerne Ind. Estate,
 Plerne, Bardez, Goa 403511, India
 Manufactured for:
 **UNICHEM**
 PHARMACEUTICALS (USA) INC.
 East Brunswick, NJ 08816
 13031918
 Repackaged By: Preferred Pharmaceuticals Inc.

This Medication Guide has been approved by the U.S. Food and Drug Administration.
 Revised: July 2024

PACKAGE LABEL-PRINCIPAL DISPLAY PANEL



MELOXICAM
 meloxicam tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC-68788-8758(NDC-2930-124)
Route of Administration	ORAL		

Active Ingredient/Active Mixture	Ingredient Name	Strength	Strength
MELOXICAM (UNII: VGZQF3CGU) (MELOXICAM - UNII:VGZQF3CGU)	MELOXICAM	7.5 mg	

Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1H32OD6U)	
COPRIMELT (UNII: 00000000000000000000000000000000)	
LACTOSE MONOHYDRATE (UNII: EWQ557Q8W3)	
MAGNESIUM STEARATE (UNII: 7009TM65D0)	
POVIDONE K30 (UNII: U723QW2X2L)	
SILICON DIOXIDE (UNII: ET7Z7ZB0U4)	
TRISODIUM CITRATE Dihydrate (UNII: B2247B95K)	

Product Characteristics

Color	YELLOW	Score	no score
Shape	ROUND	Size	7mm
Mark		Imprint Code	U.L/7.5
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	68788-8785-1	15 in 1 BOTTLE, Type O: Not a Combination Product	10/17/2024	
2	68788-8785-2	20 in 1 BOTTLE, Type O: Not a Combination Product	10/17/2024	
3	68788-8785-3	28 in 1 BOTTLE, Type O: Not a Combination Product	10/17/2024	
4	68788-8785-5	30 in 1 BOTTLE, Type O: Not a Combination Product	10/17/2024	
5	68788-8785-6	60 in 1 BOTTLE, Type O: Not a Combination Product	10/17/2024	
6	68788-8785-7	70 in 1 BOTTLE, Type O: Not a Combination Product	10/17/2024	
7	68788-8785-0	100 in 1 BOTTLE, Type O: Not a Combination Product	10/17/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077927	10/17/2024	

Labeler - Preferred Pharmaceuticals Inc. (99111902)

Registrant - Preferred Pharmaceuticals Inc. (99111902)

Establishment

Name	Address	ID/FEI	Business Operations
Preferred Pharmaceuticals Inc.	791119022	REPACK068788-8758	Preferred Pharmaceuticals Inc.

Revised: 1/2026