

**HP PRE AND POST- hydrogen peroxide, lactic acid, dodecyl benzene sulfonic acid solution
Animal Health International, Inc.**

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).

FOR INDUSTRIAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

NOT FOR HUMAN USE

WARNING

Causes eye irritation. Wash skin thoroughly after handling. Do not mix with bleach or other chlorinated products - will cause chlorine gas

DIRECTIONS:

IMPORTANT: Do not further dilute with water or mix with any other teat dips. If product in dip cup becomes visibly dirty, discard contents and replenish with fresh product. Do not reuse or return any unused product to the original container. Udder Prep: When using an udder wash step before milking, make sure to wash teats with appropriate udder wash solution using proper cleaning procedures. Teats should then be dried with single-service towels.

Directions for Teat Dipping

Pre-Milk Dipping: Before each cow is milked, and using fresh HP Pre & Post, dip each teat full-length into the teat dip cup. Wipe teats after dipping using single-service towels to avoid contamination of milk.

Post-Milk Dipping: Using fresh HP Pre & Post, dip each teat full-length into the teat dip cup. Do not wipe. Allow to air dry. Do not turn cows out in freezing weather until the product is completely dry.

Directions for Teat Spraying

Pre-Milk Spraying: Before each cow is milked, and using fresh HP Pre & Post, spray entire teat. Wipe teats after dipping using single-service towels to avoid contamination of milk.

Post-Milk Spraying: Using fresh HP Pre & Post, spray entire teat immediately after each milking. Do not wipe. Allow to air dry. Do not turn cows out in freezing weather until the product is completely dry.

Expanded Usage: When freshening cows, begin dipping teats twice daily for about 10 days before calving. **PRECAUTION:** HP Pre & Post is not intended to cure or help the healing of chapped or irritated teats. As with any germicide, irritation or sensitization may occur in sensitive animals. In case of teat irritation or chapping, have the condition examined and, if necessary, treated by a veterinarian.

FIRST AID

Eye Contact: Rinse with plenty of water.

Skin Contact: Rinse with water.

Ingestion: Rinse mouth. Get medical attention if symptoms occur.

Inhalation: Get medical attention if symptoms occur.

READ SAFETY DATA SHEET (SDS) BEFORE USING THIS PRODUCT

EMERGENCY HEALTH INFORMATION: Contact local poison control.

Representative Label and Principal Display

aspen
DAIRY SOLUTIONS

HP PRE & POST™

Sanitizing Teat Dip

- An aid in reducing the spread of organisms which may cause mastitis

ACTIVE INGREDIENTS:

Hydrogen peroxide.....	0.5%
Lactic acid.....	1.7%
Dodecyl benzene sulfonic acid.....	0.5%

SKIN CONDITIONING SYSTEM:

(contains sorbitol, glycerin).....8.0%

INERT INGREDIENTS:..... 89.3%

TOTAL:.....100.0%

Manufactured for:

Aspen Veterinary Resources®, Ltd.
3155 Heartland Drive, Liberty, MO 64068
1-800-203-5620
761356/5300/0315 6301558

NET CONTENTS: 208 L (55 US GAL)

6301558



HP PRE & POST™



ACTIVE INGREDIENTS:	
Hydrogen peroxide.....	0.5%
Lactic acid.....	1.7%
Dodecyl benzene sulfonic acid.....	0.5%
SKIN CONDITIONING SYSTEM:	
(contains sorbitol, glycerin).....	8.0%
INERT INGREDIENTS:	89.3%
TOTAL:	100.0%

Sanitizing Teat Dip

• An aid in reducing the spread of organisms which may cause mastitis

FOR INDUSTRIAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

NOT FOR HUMAN USE

DIRECTIONS:

IMPORTANT: Do not further dilute with water or mix with any other teat dips. If product in dip cup becomes visibly dirty, discard contents and replenish with fresh product. Do not reuse or return any unused product to the original container.
Udder Prep: When using an udder wash step before milking, make sure to wash teats with appropriate udder wash solution using proper cleaning procedures. Teats should then be dried with single-service towels.

Directions for Teat Dipping

Pre-Milk Dipping: Before each cow is milked, and using fresh HP Pre & Post, dip each teat full-length into the teat dip cup. Wipe teats after dipping using single-service towels to avoid contamination of milk.
Post-Milk Dipping: Using fresh HP Pre & Post, dip each teat full-length into the teat dip cup. Do not wipe. Allow to air dry. Do not turn cows out in freezing weather until the product is completely dry.

Directions for Teat Spraying

Pre-Milk Spraying: Before each cow is milked, and using fresh HP Pre & Post, spray entire teat. Wipe teats after dipping using single-service towels to avoid contamination of milk.
Post-Milk Spraying: Using fresh HP Pre & Post, spray entire teat immediately after each milking. Do not wipe. Allow to air dry. Do not turn cows out in freezing weather until the product is completely dry.

Expanded Usage: When freshening cows, begin dipping teats twice daily for about 10 days before calving.

PRECAUTION: HP Pre & Post is not intended to cure or help the healing of chapped or irritated teats. As with any germicide, irritation or sensitization may occur in sensitive animals. In case of teat irritation or chapping, have the condition examined and, if necessary, treated by a veterinarian.

WARNING

Causes eye irritation. Wash skin thoroughly after handling. Do not mix with bleach or other chlorinated products - will cause chlorine gas.

FIRST AID

Eye Contact: Rinse with plenty of water.

Skin Contact: Rinse with water.

Ingestion: Rinse mouth. Get medical attention if symptoms occur.

Inhalation: Get medical attention if symptoms occur.

READ SAFETY DATA SHEET (SDS) BEFORE USING THIS PRODUCT

EMERGENCY HEALTH INFORMATION: Contact local poison control.

Desinfectante sellador de pezones

• Ayuda a evitar la proliferación de organismos que pueden causar mastitis

PARA USO INDUSTRIAL SOLAMENTE

MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS

NO ES PARA USA HUMANO

INSTRUCCIONES:

IMPORTANTE: No lo vuelva a diluir con agua ni lo mezcle con otros productos para baño de pezones. Si el producto del recipiente para realizar el baño está visiblemente sucio, deseche el contenido y rellene el recipiente con producto fresco. No vuelva a utilizar el producto ni vuelva a colocar producto sin usar en el contenedor original.

Preparado para ubres: Si se realiza un paso de limpieza de ubres antes del ordeño, asegúrese de limpiar los pezones con una solución para limpieza de ubres adecuada y procedimientos de limpieza apropiados. Los pezones deben secarse con toallas de uso único.

Instrucciones para el sellado de pezones

Sellado del pezón previo al ordeño: Antes de ordeñar cada vaca, bañe toda la superficie de cada pezón en un recipiente que contenga HP Pre & Post fresco. Limpie los pezones con toallas de uso único luego del baño para evitar la contaminación de la leche.

Sellado del pezón posterior al ordeño: Bañe toda la superficie de cada pezón en un recipiente que contenga HP Pre & Post fresco. No utilice paños. Deje secar al aire. No deje las vacas en un clima helado hasta que el producto se haya secado por completo.

Instrucciones para el rociado de pezones

Rociado previo al ordeño: Antes de ordeñar cada vaca, rocíe toda la superficie de cada pezón con HP Pre & Post fresco. Limpie los pezones con toallas de uso único luego del baño para evitar la contaminación de la leche.

Rociado de pezones posterior al ordeño: Inmediatamente después del ordeño, rocíe toda la superficie de cada pezón con HP Pre & Post fresco. No utilice paños. Deje secar al aire. No deje las vacas en un clima helado hasta que el producto se haya secado por completo.

Uso Expandido: Al refrescar a las vacas, comience con un baño antiséptico de sus pezones dos veces por día por aproximadamente 10 días antes de parir.

PRECAUCIÓN: HP Pre & Post no está diseñado para curar o ayudar con la cicatrización de los pezones agrietados o irritados. Como cualquier otro germicida, puede provocar irritación o sensibilización en animales sensibles. En caso de irritación o agrietamiento de los pezones, haga examinar dichas afecciones y, de ser necesario, hágalas tratar con un veterinario.

ATENCIÓN

Provoca irritación ocular. Lave la piel minuciosamente después del manejo. No se mezcle con el blanqueo o otros productos tratados con cloro - causará el gas de cloro.

PRIMEROS AUXILIOS

Contacto con los ojos: Enjuague con mucha agua.

Contacto con la piel: Enjuague con agua.

Ingestión: Enjuague la boca. Consultar un médico si los síntomas aparecen.

Inhalación: Consultar un médico si los síntomas aparecen.

LEA LA FICHA DE DATOS DE SEGURIDAD ANTES DE USAR EL PRODUCTO

INFORMACIÓN DE EMERGENCIA SOBRE SALUD: Contacto local de veneno.

NET CONTENTS: 208 L (55 US GAL)

6301558



Manufactured for:
Aspen Veterinary Resources®, Ltd.
3155 Heartland Drive, Liberty, MO 64068
1-800-203-5620
761356/5300/0315

HP PRE AND POST

hydrogen peroxide, lactic acid, dodecyl benzene sulfonic acid solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:46066-404
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROGEN PEROXIDE (UNII: BBX060AN9V) (HYDROGEN PEROXIDE - UNII:BBX060AN9V)	HYDROGEN PEROXIDE	5 mg in 1 mL
LACTIC ACID (UNII: 33X04XA5AT) (LACTIC ACID - UNII:33X04XA5AT)	LACTIC ACID	17 mg in 1 mL
DODECYLBENZENESULFONIC ACID (UNII: 60NSK897G9) (DODECYLBENZENESULFONIC ACID - UNII:60NSK897G9)	DODECYLBENZENESULFONIC ACID	5 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SORBITOL (UNII: 506T60A25R)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46066-404-17	56800 mL in 1 DRUM		
2	NDC:46066-404-21	208000 mL in 1 DRUM		
3	NDC:46066-404-44	1150000 mL in 1 PACKAGE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		06/04/2015	

Labeler - Animal Health International, Inc. (627265361)

Revised: 12/2018

Animal Health International, Inc.