

TALZENNA- talazoparib capsule, liquid filled

Pfizer Laboratories Div Pfizer Inc

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use TALZENNA safely and effectively. See full prescribing information for TALZENNA.

TALZENNA® (talazoparib) capsules, for oral use
Initial U.S. Approval: 2018

INDICATIONS AND USAGE

TALZENNA is a poly (ADP-ribose) polymerase (PARP) inhibitor indicated for:
Breast Cancer

- As a single agent, for the treatment of adult patients with deleterious or suspected deleterious germline *BRCA*-mutated (g*BRCA*m) HER2-negative locally advanced or metastatic breast cancer. Select patients for therapy based on an FDA-approved companion diagnostic for TALZENNA. (1.1)

HRR Gene-mutated Metastatic Castration-Resistant Prostate Cancer (mCRPC)

- In combination with enzalutamide for the treatment of adult patients with HRR gene-mutated metastatic castration-resistant prostate cancer (mCRPC). (1.2)

DOSAGE AND ADMINISTRATION

- Take TALZENNA with or without food. (2.4)

Breast Cancer

- The recommended dosage of TALZENNA is 1 mg taken orally once daily until disease progression or unacceptable toxicity. (2.2)
- For adverse reactions, consider dosing interruption or dose reduction. (2.5)

HRR Gene-mutated Metastatic Castration-Resistant Prostate Cancer (mCRPC)

- The recommended dosage of TALZENNA is 0.5 mg taken orally once daily with enzalutamide until disease progression or unacceptable toxicity. (2.3)
- Patients should also receive a gonadotropin-releasing hormone (GnRH) analog concurrently or should have had bilateral orchiectomy. (2.3)

DOSAGE FORMS AND STRENGTHS

Capsules: 0.1 mg, 0.25 mg, 0.35 mg, 0.5 mg, 0.75 mg, and 1 mg (3)

CONTRAINDICATIONS

None. (4)

WARNINGS AND PRECAUTIONS

- Myelodysplastic Syndrome/Acute Myeloid Leukemia (MDS/AML): MDS/AML occurred in patients exposed to TALZENNA, and some cases were fatal. Monitor patients for hematological toxicity and discontinue if MDS/AML is confirmed. (5.1)
- Myelosuppression: TALZENNA may affect hematopoiesis and can cause anemia, neutropenia, and/or thrombocytopenia. (5.2)
- Embryo-Fetal Toxicity: TALZENNA can cause fetal harm. Advise of the potential risk to the fetus and to use effective contraception. (5.3, 8.1, 8.3)

ADVERSE REACTIONS

Most common adverse reactions ($\geq 20\%$) as a single agent, including laboratory abnormalities, are:

- Hemoglobin decreased, neutrophils decreased, lymphocytes decreased, platelets decreased, fatigue, glucose increased, aspartate aminotransferase increased, alkaline phosphatase increased, alanine aminotransferase increased, calcium decreased, nausea, headache, vomiting, alopecia, diarrhea, and decreased appetite. (6.1)

Most common adverse reactions ($\geq 10\%$) in combination with enzalutamide, including laboratory abnormalities, are:

- Hemoglobin decreased, neutrophils decreased, lymphocytes decreased, fatigue, platelets decreased, calcium decreased, nausea, decreased appetite, sodium decreased, phosphate decreased, fractures, magnesium decreased, dizziness, bilirubin increased, potassium decreased, and dysgeusia. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Pfizer Inc. at 1-800-438-1985 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

----- **DRUG INTERACTIONS** -----

- P-gp Inhibitors: Reduce the dose when coadministered with certain P-gp inhibitors. Monitor for increased adverse reactions. (2.7, 7.1)
- BCRP Inhibitors: Monitor for potential increased adverse reactions. (7.1)

----- **USE IN SPECIFIC POPULATIONS** -----

- Lactation: Advise women not to breastfeed. (8.2)
- Renal Impairment: Reduce the dose and monitor for increased adverse reactions for patients with moderate or severe renal impairment. (2.6, 8.7)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 6/2025

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

- 1.1 *BRCA*-mutated (g*BRCAM*) HER2-negative Locally Advanced or Metastatic Breast Cancer
- 1.2 HRR Gene-mutated Metastatic Castration-Resistant Prostate Cancer (mCRPC)

2 DOSAGE AND ADMINISTRATION

- 2.1 Patient Selection
- 2.2 Recommended Dosage for g*BRCAM* HER2-negative Locally Advanced or Metastatic Breast Cancer
- 2.3 Recommended Dosage for HRR Gene-mutated mCRPC
- 2.4 Administration
- 2.5 Dosage Modifications for Adverse Reactions
- 2.6 Recommended Dosage in Patients with Renal Impairment
- 2.7 Dosage Modifications for P-glycoprotein Inhibitors

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Myelodysplastic Syndrome/Acute Myeloid Leukemia
- 5.2 Myelosuppression
- 5.3 Embryo-Fetal Toxicity

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience

7 DRUG INTERACTIONS

- 7.1 Effect of Other Drugs on TALZENNA

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation

8.3 Females and Males of Reproductive Potential

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Hepatic Impairment

8.7 Renal Impairment

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

14.1 Deleterious or Suspected Deleterious Germline *BRCA*-mutated HER2-negative Locally Advanced or Metastatic Breast Cancer

14.2 HRR Gene-mutated mCRPC

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 *BRCA*-mutated (g*BRCA*m) HER2-negative Locally Advanced or Metastatic Breast Cancer

TALZENNA is indicated as a single agent for the treatment of adult patients with deleterious or suspected deleterious germline breast cancer susceptibility gene (*BRCA*)-mutated (g*BRCA*m) human epidermal growth factor receptor 2 (HER2)-negative locally advanced or metastatic breast cancer. Select patients for therapy based on an FDA-approved companion diagnostic for TALZENNA [see *Dosage and Administration* (2.1)].

1.2 HRR Gene-mutated Metastatic Castration-Resistant Prostate Cancer (mCRPC)

TALZENNA is indicated in combination with enzalutamide for the treatment of adult patients with homologous recombination repair (HRR) gene-mutated metastatic castration-resistant prostate cancer (mCRPC) [see *Dosage and Administration* (2.3)].

2 DOSAGE AND ADMINISTRATION

2.1 Patient Selection

Information on the FDA-approved tests for the detection of genetic mutations is

available at <http://www.fda.gov/companiondiagnostics>.

gBRCAm HER2-negative Locally Advanced or Metastatic Breast Cancer

Select patients for the treatment of advanced breast cancer with TALZENNA based on the presence of germline *BRCA* mutations [see *Indications and Usage (1.1)*, *Clinical Studies (14.1)*].

HRR Gene-mutated Metastatic Castration-Resistant Prostate Cancer

Select patients for the treatment of HRR gene-mutated mCRPC with TALZENNA based on the presence of alterations in genes directly or indirectly involved in HRR (*ATM*, *ATR*, *BRCA1*, *BRCA2*, *CDK12*, *CHEK2*, *FANCA*, *MLH1*, *MRE11A*, *NBN*, *PALB2*, or *RAD51C*) [see *Indications and Usage (1.2)*, *Clinical Studies (14.2)*].

An FDA-approved test for the detection of HRR gene mutations for use with TALZENNA is not currently available.

2.2 Recommended Dosage for gBRCAm HER2-negative Locally Advanced or Metastatic Breast Cancer

The recommended dosage of TALZENNA is 1 mg taken orally once daily, until disease progression or unacceptable toxicity.

2.3 Recommended Dosage for HRR Gene-mutated mCRPC

The recommended dosage of TALZENNA is 0.5 mg taken orally once daily with enzalutamide until disease progression or unacceptable toxicity.

Refer to the enzalutamide prescribing information for recommended enzalutamide dosing information.

Patients receiving TALZENNA and enzalutamide should also receive a gonadotropin-releasing hormone (GnRH) analog concurrently or should have had bilateral orchiectomy.

2.4 Administration

Take TALZENNA with or without food.

Swallow TALZENNA capsules whole. Do not open or dissolve.

If a patient vomits or misses a dose of TALZENNA, instruct them to take the next prescribed dose at the usual time.

2.5 Dosage Modifications for Adverse Reactions

To manage adverse reactions, consider interruption of treatment with or without dose reduction based on severity and clinical presentation. Recommended dose reductions are indicated in Table 1 and Table 2. Treatment with TALZENNA should be discontinued if more than 3 dose reductions are required.

gBRCAm HER2-negative Locally Advanced or Metastatic Breast Cancer

**Table 1. Dose Reduction Levels for Adverse Reactions—
Breast Cancer**

Dose Reductions	Dose Level
Recommended starting dose	1 mg once daily
First dose reduction	0.75 mg once daily
Second dose reduction	0.5 mg once daily
Third dose reduction	0.25 mg once daily

HRR Gene-mutated mCRPC

Table 2. Dose Reduction Levels for Adverse Reactions—mCRPC

Dose Reductions	Dose Level
Recommended starting dose	0.5 mg once daily
First dose reduction	0.35 mg once daily
Second dose reduction	0.25 mg once daily
Third dose reduction	0.1 mg once daily

Refer to the enzalutamide prescribing information for dose modifications for adverse reactions associated with enzalutamide.

gBRCAm HER2-negative Locally Advanced or Metastatic Breast Cancer and HRR Gene-mutated mCRPC

Monitor complete blood counts monthly and as clinically indicated [see *Warnings and Precautions (5.2)*].

Table 3. Dose Modification and Management for Adverse Reactions

Adverse Reactions	Withhold TALZENNA Until Levels Resolve to	Resume TALZENNA
Hemoglobin <8 g/dL	≥9 g/dL	Resume TALZENNA at a reduced dose
Platelet count <50,000/μL	≥75,000/μL	
Neutrophil count <1,000/μL	≥1500/μL	
Non-hematologic Grade 3 or Grade 4	≤Grade 1	Consider resuming TALZENNA at a reduced dose or discontinue

2.6 Recommended Dosage in Patients with Renal Impairment

gBRCAm HER2-negative Locally Advanced or Metastatic Breast Cancer

The recommended dosage of TALZENNA for patients with moderate renal impairment (CLcr 30 - 59 mL/min) is 0.75 mg taken orally once daily [see *Use in Specific Populations (8.7)*].

The recommended dosage of TALZENNA for patients with severe renal impairment (CLcr 15 - 29 mL/min) is 0.5 mg taken orally once daily [see *Use in Specific Populations (8.7)*].

HRR Gene-mutated mCRPC

The recommended dosage of TALZENNA for patients with moderate renal impairment (CLcr 30 - 59 mL/min) is 0.35 mg taken orally once daily with enzalutamide [see *Use in Specific Populations (8.7)*].

The recommended dosage of TALZENNA for patients with severe renal impairment (CLcr 15 - 29 mL/min) is 0.25 mg taken orally once daily with enzalutamide [see *Use in Specific Populations (8.7)*].

2.7 Dosage Modifications for P-glycoprotein Inhibitors

gBRCAm HER2-negative Locally Advanced or Metastatic Breast Cancer

Avoid coadministration of TALZENNA with the following P-glycoprotein (P-gp) inhibitors: itraconazole, amiodarone, carvedilol, clarithromycin, itraconazole, and verapamil. If coadministration of TALZENNA with these P-gp inhibitors cannot be avoided, reduce the dose of TALZENNA to 0.75 mg taken orally once daily. When the P-gp inhibitor is discontinued, increase the dose of TALZENNA (after 3 - 5 half-lives of the P-gp inhibitor) to the dose of TALZENNA that was used before starting the P-gp inhibitor [see *Drug Interactions (7.1)*].

Monitor for increased adverse reactions and modify the dosage as recommended for adverse reactions when TALZENNA is coadministered with other P-gp inhibitors [see *Dosage and Administration (2.5)*].

3 DOSAGE FORMS AND STRENGTHS

Capsules: Each capsule is a liquid-filled soft-gelatin capsule with the following description.

Table 4. Dosage Forms and Strengths

Strength	Description
0.1 mg	Red oval capsule (printed with "TLZ" and "010")
0.25 mg	Yellow oblong capsule (printed with "TLZ" and "025")
0.35 mg	Pink oblong capsule (printed with "TLZ" and "035")
0.5 mg	Pale orange oblong capsule (printed with "TLZ" and "050")
0.75 mg	Red oblong capsule (printed with "TLZ" and "075")
1 mg	Reddish brown oblong capsule (printed with "TLZ" and "1")

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Myelodysplastic Syndrome/Acute Myeloid Leukemia

Myelodysplastic Syndrome/Acute Myeloid Leukemia (MDS/AML), including cases with a fatal outcome, has been reported in patients who received TALZENNA.

Overall, MDS/AML has been reported in 0.4% (3 out of 788) of solid tumor patients treated with TALZENNA as a single agent in clinical studies. In TALAPRO-2, MDS/AML occurred in 2 out of 511 (0.4%) patients treated with TALZENNA and enzalutamide and in 0 out of 517 (0%) patients treated with placebo and enzalutamide [see *Adverse Reactions (6.1)*]. The durations of TALZENNA treatment in these 5 patients prior to developing MDS/AML were 0.3, 1, 2, 3, and 5 years. Most of these patients had received previous chemotherapy with platinum agents and/or other DNA damaging agents including radiotherapy.

Do not start TALZENNA until patients have adequately recovered from hematological toxicity caused by previous chemotherapy. Monitor blood counts monthly during treatment with TALZENNA. For prolonged hematological toxicities, interrupt TALZENNA and monitor blood counts weekly until recovery. If counts do not recover within 4 weeks, refer the patient to a hematologist for further investigations including bone marrow analysis and blood sample for cytogenetics. If MDS/AML is confirmed, discontinue TALZENNA.

5.2 Myelosuppression

Myelosuppression consisting of anemia, neutropenia, and/or thrombocytopenia, have been reported in patients treated with TALZENNA [see *Adverse Reactions (6.1)*].

Grade ≥ 3 anemia, neutropenia, and thrombocytopenia were reported, respectively, in 39%, 21%, and 15% of patients receiving TALZENNA as a single agent. Discontinuation due to anemia, neutropenia, and thrombocytopenia occurred, respectively, in 0.7%, 0.3%, and 0.3% of patients.

In TALAPRO-2, Grade ≥ 3 anemia, neutropenia, and thrombocytopenia were reported, respectively, in 48%, 19%, and 9% of patients receiving TALZENNA and enzalutamide. Forty-two percent of patients (216/511) required a red blood cell transfusion, including 25% (127/511) who required more than one transfusion. Discontinuation due to anemia, neutropenia, and thrombocytopenia occurred, respectively, in 8%, 3%, and 0.4% of patients.

Withhold TALZENNA until patients have adequately recovered from hematological toxicity caused by previous therapy. Monitor blood counts monthly during treatment with TALZENNA. If hematological toxicities do not resolve within 28 days, discontinue TALZENNA and refer the patient to a hematologist for further investigations including bone marrow analysis and blood sample for cytogenetics [see *Dosage and Administration (2.5)*].

5.3 Embryo-Fetal Toxicity

Based on its mechanism of action and findings from animal data, TALZENNA can cause fetal harm when administered to a pregnant woman. In an animal reproduction study, administration of talazoparib to pregnant rats during the period of organogenesis caused fetal malformations and structural skeletal variations, and embryo-fetal death at

exposures that were 0.24 times the area under the concentration-time curve (AUC) in patients receiving the recommended human dose of 1 mg daily. Advise pregnant women and females of reproductive potential of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment and for 7 months following the last dose of TALZENNA [see *Use in Specific Populations (8.1, 8.3)*, *Clinical Pharmacology (12.1)*].

Based on findings from genetic toxicity and animal reproduction studies, advise male patients with female partners of reproductive potential or who are pregnant to use effective contraception during treatment and for 4 months following the last dose of TALZENNA [see *Use in Specific Populations (8.1, 8.3)*, *Nonclinical Toxicology (13.1)*].

6 ADVERSE REACTIONS

The following clinically significant adverse reactions are described elsewhere in the labeling:

- Myelodysplastic Syndrome/Acute Myeloid Leukemia [see *Warnings and Precautions (5.1)*]
- Myelosuppression [see *Warnings and Precautions (5.2)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The data described in the WARNINGS AND PRECAUTIONS section reflect exposure to single agent TALZENNA in solid tumor clinical studies, including 286 patients enrolled in EMBRACA trial and to TALZENNA 0.5 mg daily with enzalutamide in 511 patients enrolled in the TALAPRO-2 trial that included 197 patients with HRR gene-mutated mCRPC.

gBRCAm HER2-negative Locally Advanced or Metastatic Breast Cancer EMBRACA

The safety of TALZENNA as a single agent was evaluated in gBRCAm patients with HER2-negative locally advanced or metastatic breast cancer who had previously received no more than 3 lines of chemotherapy for the treatment of locally advanced/metastatic disease [see *Clinical Studies (14.1)*]. EMBRACA was a randomized, open-label, multi-center study in which 412 patients received either TALZENNA 1 mg once daily (N=286) or a chemotherapy agent (capecitabine, eribulin, gemcitabine, or vinorelbine) of the healthcare provider's choice (N=126) until disease progression or unacceptable toxicity. The median duration of study treatment was 6.1 months in patients who received TALZENNA and 3.9 months in patients who received chemotherapy.

Serious adverse reactions of TALZENNA occurred in 32% of patients. Serious adverse reactions reported in >2% of patients included anemia (6%) and pyrexia (2%). Fatal adverse reactions occurred in 1% of patients, including cerebral hemorrhage, liver disorder, veno-occlusive liver disease, and worsening neurological symptoms (1 patient each).

Permanent discontinuation due to adverse reactions occurred in 5% of TALZENNA patients. Dosing interruptions due to an adverse reaction of any grade occurred in 65%

of patients receiving TALZENNA; dose reductions due to any cause occurred in 53% of TALZENNA patients.

The most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were hemoglobin decreased, neutrophils decreased, lymphocytes decreased, platelets decreased, fatigue, glucose increased, aspartate aminotransferase increased, alkaline phosphatase increased, alanine aminotransferase increased, calcium decreased, nausea, headache, vomiting, alopecia, diarrhea, and decreased appetite.

Table 5 and Table 6 summarize the most common adverse reactions and laboratory abnormalities, respectively, in patients treated with TALZENNA or chemotherapy in the EMBRACA study.

Table 5. Adverse Reactions* ($\geq 20\%$) in Patients Receiving TALZENNA in EMBRACA

Adverse Reactions	TALZENNA N=286 (%)			Chemotherapy N=126 (%)		
	Grades 1-4	Grade 3	Grade 4	Grades 1-4	Grade 3	Grade 4
General Disorders and Administration Site Conditions						
Fatigue [†]	62	3	0	50	5	0
Gastrointestinal Disorders						
Nausea	49	0.3	0	47	2	0
Vomiting	25	2	0	23	2	0
Diarrhea	22	1	0	26	6	0
Nervous System Disorders						
Headache	33	2	0	22	1	0
Skin and Subcutaneous Tissue Disorders						
Alopecia	25	0	0	28	0	0
Metabolism and Nutrition Disorders						
Decreased appetite	21	0.3	0	22	1	0

Abbreviation: N=number of patients.

* Graded according to NCI CTCAE 4.03.

† Includes fatigue and asthenia.

Clinically relevant adverse reactions in $< 20\%$ of patients who received TALZENNA included abdominal pain (19%), dizziness (17%), dysgeusia (10%), dyspepsia (10%), stomatitis (8%), and febrile neutropenia (0.3%).

Table 6. Select Laboratory Abnormalities ($\geq 25\%$) of Patients in EMBRACA

Parameter	TALZENNA N*=286 (%)			Chemotherapy N*=126 (%)		
	Grades 1- 4	Grade 3	Grade 4	Grades 1- 4	Grade 3	Grade 4
Hemoglobin decreased	90	39	0	77	6	0
Neutrophils	68	17	3	70	21	17

decreased						
Lymphocytes decreased	76	17	0.7	53	8	0.8
Platelets decreased	55	11	4	29	2	0
Glucose increased [†]	54	2	0	51	2	0
Aspartate aminotransferase increased	37	2	0	48	3	0
Alkaline phosphatase increased	36	2	0	34	2	0
Alanine aminotransferase increased	33	1	0	37	2	0
Calcium decreased	28	1	0	16	0	0

Abbreviation: N=number of patients.

* This number represents the safety population. The derived values in the table are based on the total number of evaluable patients for each laboratory parameter.

† This number represents non-fasting glucose.

HRR Gene-mutated mCRPC

The safety of TALZENNA with enzalutamide was evaluated in patients with HRR gene-mutated mCRPC enrolled in TALAPRO-2 [see *Clinical Studies (14.2)*]. Patients were randomized to receive either TALZENNA 0.5 mg with enzalutamide 160 mg once daily (N=197), or placebo with enzalutamide 160 mg once daily (N=199) until disease progression or unacceptable toxicity. Among patients receiving TALZENNA, 86% were exposed for 6 months or longer, 60% were exposed for greater than one year, and 18% were exposed for greater than two years.

Serious adverse reactions of TALZENNA with enzalutamide occurred in 30% of patients. Serious adverse reactions reported in >2% of patients included anemia (9%) and fracture (3%). Fatal adverse reactions occurred in 1.5% of patients, including pneumonia, COVID infection, and sepsis (1 patient each).

Permanent discontinuation of TALZENNA due to adverse reactions occurred in 10% of patients treated in the TALZENNA with enzalutamide arm. The most common adverse reactions which resulted in permanent discontinuation of TALZENNA were anemia (4%), fatigue, bone fracture, ischemic heart disease, and spinal cord compression (1% each).

Dosage interruption of TALZENNA due to adverse reactions occurred in 58% of patients treated in the TALZENNA with enzalutamide arm. The most common adverse reactions which resulted in dose interruption of TALZENNA were anemia (42%), neutropenia (15%), and platelet count decreased (9%) and fatigue (5%).

Dose reduction of TALZENNA due to adverse reactions occurred in 52% of patients treated in the TALZENNA with enzalutamide arm. The most common adverse reactions which resulted in dose reduction of TALZENNA were anemia (43%), neutrophil count decreased (15%), platelet count decreased (6%), and fatigue (4%).

The most common adverse reactions ($\geq 10\%$), including laboratory abnormalities, in patients who received TALZENNA with enzalutamide were hemoglobin decreased,

neutrophils decreased, lymphocytes decreased, fatigue, platelets decreased, calcium decreased, nausea, decreased appetite, sodium decreased, phosphate decreased, fractures, magnesium decreased, dizziness, bilirubin increased, potassium decreased, and dysgeusia.

Table 7 and Table 8 summarize the most common adverse reactions and laboratory abnormalities, respectively, in the TALAPRO-2 study.

Table 7. Adverse Reactions* ($\geq 10\%$) in Patients Receiving TALZENNA (with a Difference Between Arms of $\geq 2\%$) in TALAPRO-2

	TALZENNA with Enzalutamide N=197			Placebo with Enzalutamide N=199		
	Grades 1-4 %	Grade 3 %	Grade 4 %	Grades 1-4 %	Grade 3 %	Grade 4 %
Fatigue [†]	49	4	0	40	1	0
Nausea	21	2	0	17	1	0.5
Decreased appetite	20	1	0	14	1	1
Fractures [‡]	14	3	0	10	1.5	0
Dizziness [§]	13	1.5	0	9	1.5	0
Dysgeusia [¶]	10	0	0	4.5	0	0

Abbreviation: N=number of patients.

* Graded according to NCI CTCAE 4.03.

† Includes fatigue and asthenia.

‡ Fractures include multiple similar terms.

§ Includes dizziness, dizziness postural, vertigo.

¶ Includes ageusia, anosmia, dysgeusia.

Clinically relevant adverse reactions in $< 10\%$ of patients who received TALZENNA with enzalutamide included abdominal pain (9%), vomiting (9%), alopecia (7%), dyspepsia (4%), venous thromboembolism (3%) and stomatitis (2%).

Table 8. Select Laboratory Abnormalities ($\geq 10\%$) That Worsened from Baseline in Patients Who Received TALZENNA in TALAPRO-2

Laboratory Abnormality	TALZENNA with Enzalutamide N=197*			Placebo with Enzalutamide N=199*		
	Grades 1-4 %	Grade 3 %	Grade 4 %	Grades 1-4 %	Grade 3 %	Grade 4 %
Hemoglobin decreased	79	41	0	34	6	0
Neutrophils decreased	60	18	1	18	0	1
Lymphocytes decreased	58	13	0	36	7	0
Platelets decreased	45	6	3	8	0.5	0
Calcium	25	0	1	11	0	1

decreased						
Sodium decreased	22	3	0	20	1.5	0
Phosphate decreased	17	3	1	13	2	0
Magnesium decreased	14	0	1	12	0	0.5
Bilirubin increased	11	0.5	0	7	0	0
Potassium decreased	11	0	1	7	1	0.5

Abbreviation: N=number of patients.

* The denominator used to calculate the rate varied from 198 to 199 in the placebo with enzalutamide arm based on the number of patients with a baseline value and at least one post-treatment value.

7 DRUG INTERACTIONS

7.1 Effect of Other Drugs on TALZENNA

Effect of P-gp Inhibitors

Breast Cancer

Avoid coadministration of TALZENNA with the following P-gp inhibitors: itraconazole, amiodarone, carvedilol, clarithromycin, itraconazole, and verapamil. If coadministration of TALZENNA with these P-gp inhibitors cannot be avoided, reduce the dose of TALZENNA [see *Dosage and Administration (2.7)*]. When the P-gp inhibitor is discontinued, increase the dose of TALZENNA [see *Dosage and Administration (2.7)*].

Coadministration of TALZENNA with these P-gp inhibitors increased talazoparib concentrations [see *Clinical Pharmacology (12.3)*], which may increase the risk of adverse reactions.

Monitor for increased adverse reactions and modify the dosage as recommended for adverse reactions when TALZENNA is coadministered with other P-gp inhibitors [see *Dosage and Administration (2.5)*].

HRR Gene-mutated mCRPC

The effect of coadministration of P-gp inhibitors on talazoparib exposure when TALZENNA is taken with enzalutamide has not been studied. Monitor patients for increased adverse reactions and modify the dosage as recommended for adverse reactions when TALZENNA is coadministered with a P-gp inhibitor [see *Dosage and Administration (2.5)*].

Effect of Breast Cancer Resistance Protein (BCRP) Inhibitors

Monitor patients for increased adverse reactions and modify the dosage as recommended for adverse reactions when TALZENNA is coadministered with a BCRP inhibitor [see *Dosage and Administration (2.5)*].

Coadministration of TALZENNA with BCRP inhibitors may increase talazoparib exposure [*see Clinical Pharmacology (12.3)*], which may increase the risk of adverse reactions.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Based on findings from animal studies and its mechanism of action [*see Clinical Pharmacology (12.1)*], TALZENNA can cause embryo-fetal harm when administered to a pregnant woman. There are no available data on TALZENNA use in pregnant women to inform a drug-associated risk. In an animal reproduction study, the administration of talazoparib to pregnant rats during the period of organogenesis caused fetal malformations and structural skeletal variations and embryo-fetal death at maternal exposures that were 0.24 times the AUC in patients receiving the recommended dose of 1 mg daily (*see Data*). Apprise pregnant women and females of reproductive potential of the potential risk to a fetus.

The background risk of major birth defects and miscarriage for the indicated population is unknown. In the general U.S. population, the estimated background risks of major birth defects and miscarriage in clinically recognized pregnancies are 2% to 4% and 15% to 20%, respectively.

Data

Animal Data

In an embryo-fetal development toxicity study, pregnant rats received oral doses of 0.015, 0.05, and 0.15 mg/kg/day talazoparib during the period of organogenesis. Talazoparib caused embryo-fetal death at doses ≥ 0.015 mg/kg/day (approximately 0.24 times the AUC in patients at the recommended dose of 1 mg daily). A dose of 0.015 mg/kg/day caused decreased fetal body weights and an increased incidence of fetal malformations (depressed eye bulge, small eye, split sternebra, and fused cervical vertebral arch) and structural variations including misshapen or incomplete ossification of the sternebra, skull, rib, and vertebra.

8.2 Lactation

Risk Summary

There are no data on the presence of talazoparib in human milk, the effects of the drug on milk production, or the effects of the drug on the breastfed child. Because of the potential for serious adverse reactions in a breastfed child from talazoparib, advise lactating women not to breastfeed during treatment with TALZENNA and for 1 month after the last dose.

8.3 Females and Males of Reproductive Potential

TALZENNA can cause fetal harm when administered to pregnant women [*see Use in Specific Populations (8.1)*].

Pregnancy Testing

Verify pregnancy status in females of reproductive potential prior to initiating TALZENNA treatment.

Contraception

Females

Advise females of reproductive potential to use effective contraception during treatment and for 7 months following the last dose of TALZENNA.

Males

Based on genotoxicity and animal reproduction studies, advise male patients with female partners of reproductive potential and pregnant partners to use effective contraception during treatment with TALZENNA and for 4 months following the last dose [see *Use in Specific Populations (8.1)*, *Nonclinical Toxicology (13.1)*].

Infertility

Males

Based on animal studies, TALZENNA may impair fertility in males of reproductive potential [see *Nonclinical Toxicology (13.1)*].

8.4 Pediatric Use

The safety and effectiveness of TALZENNA have not been established in pediatric patients.

8.5 Geriatric Use

In clinical trials of TALZENNA enrolling 494 patients with advanced solid tumors who received TALZENNA 1 mg daily as a single agent, 85 (17%) patients were ≥ 65 years of age, and this included 19 (4%) patients who were ≥ 75 years old. There were 5 patients ≥ 85 years old. In the TALAPRO-2 trial, of 197 patients who received TALZENNA, 77% were ≥ 65 years of age, while 30% were ≥ 75 years of age. No overall differences in safety or effectiveness of TALZENNA were observed between these patients and younger patients.

8.6 Hepatic Impairment

No dosage modification is recommended for patients with hepatic impairment [see *Clinical Pharmacology (12.3)*].


8.7 Renal Impairment

Reduce the recommended dosage of TALZENNA in patients with moderate (CLcr 30 – 59 mL/min) and severe (CLcr 15 – 29 mL/min) renal impairment [see *Dosage and Administration (2.7)*]. Monitor patients with severe renal impairment for increased adverse reactions and modify the dosage as recommended for adverse reactions [see *Dosage and Administration (2.5)*].

No dose adjustment is recommended for patients with mild renal impairment (CLcr 60 – 89 mL/min). TALZENNA has not been studied in patients requiring hemodialysis.

11 DESCRIPTION

Talazoparib is an inhibitor of mammalian polyadenosine 5'-diphosphoribose (ADP-ribose) polymerase (PARP) enzymes. TALZENNA contains talazoparib tosylate. The chemical name of talazoparib tosylate is (8*S*,9*R*)-5-Fluoro-8-(4-fluorophenyl)-9-(1-methyl-1*H*-1,2,4-triazol-5-yl)-2,7,8,9-tetrahydro-3*H*-pyrido[4,3,2-*de*]phthalazin-3-one 4-methylbenzenesulfonate (1:1). The chemical formula of talazoparib tosylate is C₂₆H₂₂F₂N₆O₄S, and the relative molecular mass is 552.56 Daltons. The chemical structure of talazoparib tosylate is shown below:



Talazoparib tosylate is a white to yellow solid. TALZENNA capsules for oral use are available as liquid-filled soft gelatin capsules. Each capsule contains 0.1 mg, 0.25 mg, 0.35 mg, 0.5 mg, 0.75, or 1 mg of talazoparib equivalent to 0.145 mg, 0.363 mg, 0.509 mg, 0.727 mg, 1.09 mg, or 1.453 mg talazoparib tosylate, respectively.

TALZENNA capsules contain the following inactive ingredients: gelatin, glycerol, iron oxide (red), iron oxide (yellow), polyethylene glycol 400, purified water, sorbitol, titanium dioxide, and tocopherol.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Talazoparib is an inhibitor of PARP enzymes, including PARP1 and PARP2, which play a role in DNA repair. In vitro studies with cancer cell lines that harbored defects in DNA repair genes, including *BRCA1* and *BRCA2*, have shown that talazoparib-induced cytotoxicity may involve inhibition of PARP enzymatic activity and increased formation of PARP-DNA complexes resulting in DNA damage, decreased cell proliferation, and apoptosis. Talazoparib anti-tumor activity was observed in patient-derived xenograft breast cancer models bearing mutated *BRCA1* or mutated *BRCA2* or wild type *BRCA1* and *BRCA2*.

12.2 Pharmacodynamics

The exposure-response relationship and time course of pharmacodynamic response for the safety and effectiveness of TALZENNA have not been fully characterized.

Cardiac Electrophysiology

At a dose of 1 mg (the recommended dosage for treatment of breast cancer), TALZENNA had no large QTc prolongation (i.e., >20 ms).

12.3 Pharmacokinetics

After administration of TALZENNA 1 mg orally once daily as a single agent (the recommended dosage for breast cancer), the mean [% coefficient of variation (CV%)] AUC and maximum observed plasma concentration (C_{max}) of talazoparib at steady-state was 173 (32%) ng.hr/mL and 19 (32%) ng/mL, respectively. The mean (CV%) steady-state C_{trough} was 3.5 (61%) ng/mL.

After administration of TALZENNA 0.5 mg orally once daily (the recommended dosage for prostate cancer) with enzalutamide, the mean (CV%) steady-state C_{trough} ranged from 3.3 to 3.7 ng/mL (45% to 48%).

The pharmacokinetics (PK) of talazoparib is linear from 0.025 mg to 2 mg (2 times the recommended dose for breast cancer). The median accumulation ratio of talazoparib following 1 mg orally once daily is 2.3 to 5.2. Talazoparib plasma concentrations reached steady-state within 2 to 3 weeks when administered as a single agent and within 9 weeks when coadministered with enzalutamide.

Absorption

The median time to C_{max} (T_{max}) was approximately 1 hour (range: 0.4 to 4 hours) after dosing.

Food Effect

Following administration of a TALZENNA 1 mg dose soft gelatin capsule with high-fat, high-calorie food (approximately 800 to 1,000 calories with 150, 250, and 500 to 600 calories from protein, carbohydrate, and fat, respectively), the mean steady-state C_{max} was decreased by 42%, the median T_{max} was delayed from 1 to 4 hours, and AUC was not affected as compared to those under fasted conditions.

Distribution

The mean apparent volume of distribution of talazoparib is 420 L. *In vitro*, protein binding of talazoparib is 74% and is independent of talazoparib concentration.

Elimination

The mean terminal plasma half-life (\pm standard deviation) is 90 (\pm 58) hours and the mean apparent oral clearance (inter-subject variability) is 6.5 L/h (31%).

Metabolism

Talazoparib undergoes minimal hepatic metabolism. The identified metabolic pathways include mono-oxidation, dehydrogenation, cysteine conjugation of mono-desfluoro-talazoparib, and glucuronide conjugation.

Excretion

Excretion of talazoparib in urine was the major route of elimination. Approximately 69% (55% unchanged) of the total administered radiolabeled dose of talazoparib was recovered in urine, and 20% (14% unchanged) was recovered in feces.

Specific Populations

Age (18 to 88 years), sex, race (361 White, 41 Asian, 16 Black, 9 Others, and 63 Not Reported), body weight (36 to 162 kg), and mild to severe hepatic impairment had no clinically significant effect on the PK of talazoparib.

Patients with Renal Impairment

Mild (eGFR 60 – 89 mL/min/1.73 m²) renal impairment had no clinically significant effect on talazoparib pharmacokinetics. Talazoparib steady-state total exposure (AUC) increased by 43% in subjects with moderate (eGFR 30 – 59 mL/min/1.73 m²) renal impairment and 163% in patients with severe (eGFR 15 – 29 mL/min/1.73 m²) renal impairment relative to subjects with normal renal function (eGFR ≥ 90 mL/min/1.73 m²). Talazoparib steady-state peak concentration (C_{max}) increased by 32% in subjects with moderate renal impairment and 89% in subjects with severe renal impairment, relative to subjects with normal renal function. Similar increases in AUC were observed with talazoparib when given with enzalutamide for patients with moderate and severe renal impairment. The PK of talazoparib has not been studied in patients requiring hemodialysis. There was no evidence of a relationship between the protein binding of talazoparib and renal function.

Drug Interaction Studies

Clinical Studies

Effect of P-gp Inhibitors: Coadministration of a P-gp inhibitor (itraconazole) with a single 0.5 mg dose of TALZENNA increased talazoparib AUC and C_{max} by approximately 56% and 40%, respectively. Coadministration with the following other P-gp inhibitors: amiodarone, carvedilol, clarithromycin, itraconazole, and verapamil increased talazoparib exposure by 45%.

Coadministration with other P-gp inhibitors (including azithromycin, atorvastatin, diltiazem, felodipine, fluvoxamine, and quercetin) had no clinically significant effect on talazoparib pharmacokinetics.

Effect of P-gp Inducers: Coadministration of a P-gp inducer (rifampin) with a single 1 mg dose of TALZENNA increased talazoparib C_{max} by 37% with no effect on talazoparib AUC.

Effect of Acid-Reducing Agents: Coadministration of acid-reducing agents including proton pump inhibitors (PPI), histamine receptor 2 antagonists (H₂RA), or other acid-reducing agents has no effect on the absorption of talazoparib.

Enzalutamide: Coadministration of enzalutamide with TALZENNA increased talazoparib exposure approximately 2-fold.

In Vitro Studies

Transporters: Talazoparib is a substrate of P-gp and BCRP transporters, but not a substrate of OATP1B1, OATP1B3, OCT1, OCT2, OAT1, OAT3, BSEP, MATE1, or MATE2-K.

Talazoparib is not an inhibitor of P-gp, BCRP, OATP1B1, OATP1B3, OCT1, OCT2, OAT1, OAT3, BSEP, MATE1, or MATE2-K.

CYP Enzymes: Talazoparib is not an inhibitor of CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, or CYP3A4/5.

Talazoparib is not an inducer of CYP1A2, CYP2B6, or CYP3A4.

UGT: Talazoparib is not an inhibitor of UGT isoforms (1A1, 1A4, 1A6, 1A9, 2B7, and

2B15).

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenicity studies have not been conducted with talazoparib.

Talazoparib was clastogenic in an in vitro chromosomal aberration assay in human peripheral blood lymphocytes and in an in vivo bone marrow micronucleus assay in rats. This clastogenicity is consistent with genomic instability resulting from the primary pharmacology of talazoparib, indicating the potential for genotoxicity in humans. Talazoparib was not mutagenic in a bacterial reverse mutation (Ames) test.

Fertility studies in animals have not been conducted with talazoparib. In repeat-dose toxicity studies up to 3-months duration, talazoparib-related findings in the testis and epididymis at doses ≥ 0.04 mg/kg/day in rats and ≥ 0.01 mg/kg/day in dogs included decreased organ weights, luminal cellular debris, reduced sperm, and degeneration/atrophy. These doses in rats and dogs resulted in approximately 1.0 times and 0.2 times, respectively, the exposure (AUC) in humans at the recommended dose of 1 mg daily. Follicular atresia of the ovary was observed in rats at doses ≥ 1 mg/kg/day talazoparib, approximately 9.5 times the AUC in patients at the recommended dose of 1 mg daily.

14 CLINICAL STUDIES

14.1 Deleterious or Suspected Deleterious Germline *BRCA*-mutated HER2-negative Locally Advanced or Metastatic Breast Cancer

EMBRACA (NCT01945775) was an open-label study in which patients (N=431) with g*BRCA*m HER2-negative locally advanced or metastatic breast cancer were randomized 2:1 to receive TALZENNA 1 mg or healthcare provider's choice of chemotherapy (capecitabine, eribulin, gemcitabine, or vinorelbine) until disease progression or unacceptable toxicity. Randomization was stratified by prior lines of chemotherapy for metastatic disease (0 versus 1, 2, or 3), by triple-negative disease status [triple-negative breast cancer (TNBC) versus non-TNBC], and history of central nervous system (CNS) metastasis (yes versus no).

Patients received no more than 3 prior cytotoxic chemotherapy regimens for their metastatic or locally advanced disease. Patients were required to have received treatment with an anthracycline and/or a taxane (unless contraindicated) in the neoadjuvant, adjuvant, and/or metastatic treatment setting. First-line treatment for advanced or metastatic disease with no prior adjuvant chemotherapy was allowed if the investigator determined that 1 of the 4 chemotherapy choices in the control arm would be an appropriate treatment option for the patient. Patients with prior platinum therapy for advanced disease were required to have no evidence of disease progression during platinum therapy. No prior treatment with a PARP inhibitor was permitted. Of the 431 patients randomized in the EMBRACA study, 408 (95%) were centrally confirmed to have

a deleterious or suspected deleterious gBRCAm using a clinical trial assay; out of which 354 (82%) were confirmed using the BRCAAnalysis CDx[®]. BRCA mutation status [breast cancer susceptibility gene 1 (BRCA1)-positive or breast cancer susceptibility gene 2 (BRCA2)-positive] was similar across both treatment arms.

The median age of patients treated with TALZENNA was 46 years (range 28 to 84) and 51 years (range 24 to 89) among patients treated with chemotherapy. Among all randomized patients, 1% versus 2% were males, 67% versus 75% were White; 11% versus 11% were Asian, and 4% versus 1% were Black or African American in the TALZENNA and chemotherapy arms, respectively. Almost all patients (98%) in both arms had an Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1. Approximately 56% of patients had estrogen receptor-positive and/or progesterone receptor-positive disease; 44% of patients had triple-negative disease, and the proportions were balanced across both treatment arms. Fifteen percent (15%) of patients in the TALZENNA arm and 14% of patients in the chemotherapy arm had a history of CNS metastases. Ninety-one percent (91%) of patients in the TALZENNA arm had received prior taxane therapy, and 85% had received prior anthracycline therapy in any setting. Sixteen percent (16%) of patients in the TALZENNA arm and 21% of patients in the chemotherapy arm had received prior platinum treatment in any setting. The median number of prior cytotoxic regimens for patients with advanced breast cancer was one; 38% received no prior cytotoxic regimens for advanced or metastatic disease, 37% received one, 20% received two, and 5% received three or more prior cytotoxic regimens.

The major efficacy outcome measure was progression-free survival (PFS) evaluated according to Response Evaluation Criteria in Solid Tumors (RECIST) version 1.1, as assessed by blinded independent central review (BICR). A statistically significant improvement in PFS was demonstrated for TALZENNA compared to chemotherapy. A sensitivity analysis of investigator-assessed PFS was consistent with the BICR-assessed PFS results. Consistent PFS results were observed across patient subgroups defined by study stratification factors (prior lines of chemotherapy, TNBC status, and history of CNS metastases). Efficacy data from the EMBRACA study are summarized in Table 9, and the Kaplan-Meier curves for PFS are shown in Figure 1 and final overall survival (OS) in Figure 2.

Table 9. Efficacy Results - EMBRACA Study

	TALZENNA	Chemotherapy
PFS by BICR	N=287	N=144
Disease progression or deaths, n (%)	186 (65)	83 (58)
Median months (95% CI)	8.6 (7.2, 9.3)	5.6 (4.2, 6.7)
Hazard ratio (95% CI)*	0.54 (0.41, 0.71)	
p-value [†]	p<0.0001	
Patients with Measurable Disease by Investigator[‡]	N=219	N=114
ORR, % (95% CI) [§]	50.2 (43.4, 57.0)	18.4 (11.8, 26.8)
Median [¶] DOR months (95% CI)	6.4 (5.4, 9.5)	3.9 (3.0, 7.6)
OS	N=287	N=144
Deaths, n (%)	216 (75)	108 (75)

Median months (95% CI)	19.3 (16.6, 22.5)	19.5 (17.4, 22.4)
Hazard ratio (95% CI)*	0.85 (0.67, 1.07)	
p-value†	p=0.1693	

Abbreviations: BICR=blinded independent central review; CI=confidence interval; DOR=duration of response; ITT=intent-to-treat; N=number of patients; ORR=objective response rate; OS=overall survival; PFS=progression-free survival.

* Hazard ratio is estimated from a Cox proportional hazards model stratified by prior use of chemotherapy for metastatic disease (0 versus 1, 2, or 3), by triple-negative disease status [triple-negative breast cancer (TNBC) versus non TNBC], and by history of central nervous system metastasis (yes versus no) and is relative to overall chemotherapy with <1 favoring talazoparib.


† p-values (2-sided) from the log-rank test stratified by number of prior cytotoxic chemotherapy regimens, triple-negative status and history of central nervous system metastasis.

‡ Conducted in ITT population with measurable disease at baseline.

§ Response rate based on confirmed responses.

¶ Median estimated from Kaplan-Meier probabilities.

Figure 1. Kaplan-Meier Curves of PFS - EMBRACA Study




Number of patients at risk

TALZENNA	287	229	148	91	55	42	29	23	16	12	5	3	1
Chemotherapy	144	68	34	22	9	8	4	2	2	1			

Abbreviation: PFS=progression-free survival.

Figure 2. Kaplan-Meier Curves of OS - EMBRACA Study (ITT Population)



Abbreviations: ITT=intent-to-treat; OS=overall survival.

14.2 HRR Gene-mutated mCRPC

The efficacy of TALZENNA with enzalutamide was evaluated in TALAPRO-2 (NCT03395197), a randomized, double-blind, placebo-controlled, multi-cohort trial in which 399 patients with HRR gene-mutated (HRRm) mCRPC were randomized 1:1 to receive enzalutamide 160 mg daily plus either TALZENNA 0.5 mg or placebo daily until unacceptable toxicity or disease progression. Mutation status of HRR genes was determined prospectively using solid tumor tissue or circulating tumor DNA (ctDNA)-based next generation sequencing assays. Patients were required to have a mutation in at least one of 12 genes involved directly or indirectly in the HRR pathway (*ATM*, *ATR*, *BRCA1*, *BRCA2*, *CDK12*, *CHEK2*, *FANCA*, *MLH1*, *MRE11A*, *NBN*, *PALB2*, or *RAD51C*). All patients received a GnRH analog or had prior bilateral orchiectomy and needed to have progressed on prior androgen deprivation therapy. Prior treatment with a CYP17 inhibitor or docetaxel for metastatic castration-sensitive prostate cancer (mCSPC) was permitted. Randomization was stratified by previous treatment with a CYP17 inhibitor or docetaxel (yes/no).

The median age was 70 years (range: 41 to 90); 100% were male; 68% were White, 21% Asian, 2.8% Black, 0.8% Other, 7% unknown/not reported; 12% were Hispanic/Latino; and baseline ECOG performance status was 0 (62%) or 1 (38%). Thirty-nine percent of patients had bone-only disease; 15% had visceral disease. In the mCSPC setting, 29% percent of patients had received docetaxel and 9% had received a prior CYP17 inhibitor. The most commonly mutated HRR genes (>5%), including co-occurring mutations, were: *BRCA2* (34%), *ATM* (22%), *CDK12* (19%), *CHEK2* (18%), and *BRCA1* (6%).

The major efficacy outcome measure was radiographic progression-free survival (rPFS) evaluated according to RECIST, version 1.1 and Prostate Cancer Working Group (PCWG3) (bone) criteria, assessed by BICR. An additional efficacy outcome measure was OS.

A statistically significant improvement in rPFS and OS were demonstrated in patients randomized to TALZENNA with enzalutamide compared with placebo with enzalutamide. Efficacy results are presented in Table 10, and Figures 3 and 4.

Table 10. Efficacy Results - TALAPRO-2 (HRR Gene-mutated mCRPC)


	TALZENNA with Enzalutamide N=200	Placebo with Enzalutamide N=199
Radiographic Progression-free Survival (rPFS) by BICR		
Number of events, n (%)	66 (33)	104 (52)
Median months (95% CI)	NR (21.9, NR)	13.8 (11.0, 16.7)
Hazard ratio (95% CI)*	0.45 (0.33, 0.61)	
p-value†	p<0.0001	
Overall Survival (OS)		
Deaths, n (%)	93 (47)	126 (63)
Median months (95% CI)	45.1 (35.4, NR)	31.1 (27.3, 35.4)
Hazard ratio (95% CI)*	0.62 (0.48, 0.81)	
p-value†	p=0.0005	

Abbreviations: BICR=blinded independent central review; CI=confidence interval; CSPC=castration-sensitive prostate cancer; HRR=homologous recombination repair; mCRPC=metastatic castration-resistant prostate cancer; N=number of patients; NR=not reached.

* Hazard ratio and CI are based on Cox PH model stratified by previous treatment for CSPC.

† p-value (2-sided) is based on log-rank test stratified by previous treatment for CSPC.

Figure 3. Kaplan-Meier Curves of rPFS - TALAPRO-2 (HRR Gene-mutated mCRPC)




Number of patients at risk

TALZENNA + Enzalutamide	200	191	180	168	163	131	107	86	82	60	49	45	34	26	21	19	9	4	2	1	0
Placebo + Enzalutamide	199	171	149	131	126	96	67	51	47	38	29	25	21	11	7	7	4	0	0	0	0

Abbreviations: HRR=homologous recombination repair; mCRPC=metastatic castration-resistant prostate cancer; rPFS=radiographic progression-free survival.

Figure 4. Kaplan-Meier Curves of OS - TALAPRO-2 (HRR Gene-mutated

mCRPC)



Abbreviations: HRR=homologous recombination repair; mCRPC=metastatic castration-resistant prostate cancer; OS=overall survival.

Exploratory subgroup analyses of rPFS and OS for patients with *BRCA*-mutated (*BRCA*m) and non-*BRCA*m HRRm are presented in Table 11.

Table 11. Exploratory rPFS and OS Subgroup Analyses by *BRCA*m Status for TALAPRO-2 (HRR Gene-mutated mCRPC)

	<i>BRCA</i> m		Non- <i>BRCA</i> m HRRm*	
	TALZENNA with Enzalutamide N=71	Placebo with Enzalutamide N=84	TALZENNA with Enzalutamide N=129	Placebo with Enzalutamide N=115
rPFS				
Number of events, n (%)	15 (21)	54 (64)	51 (40)	50 (43)
Median months (95% CI)	NR (NR, NR)	11.0 (8.3, 11.1)	24.7 (16.4, NR)	16.7 (13.8, 27.7)
Hazard ratio (95% CI)	0.20 (0.11, 0.36)		0.74 (0.50, 1.09)	
Overall Survival (OS)				
Deaths, n (%)	30 (42)	56 (67)	63 (49)	70 (61)
Median months (95% CI)	NR (35.4, NR)	28.5 (24.5, 34.4)	42.4 (34.2, NR)	32.6 (28.0, 42.2)

Hazard ratio (95% CI) [†]	0.48 (0.31, 0.75)	0.71 (0.51, 1.00)
------------------------------------	-------------------	-------------------

Abbreviations: *BRCAM*=breast cancer susceptibility gene-mutated; CI=confidence interval; *CSPC*=castration-sensitive prostate cancer; *HRRm*=homologous recombination repair gene-mutated; NR=not reached; rPFS=radiographic progression-free survival.

* Includes 4 patients who were incorrectly randomized in the *HRRm* stratum who did not have *HRR* gene mutations.

† Hazard ratio and CI are based on Cox PH model unstratified by previous treatment for *CSPC*.

16 HOW SUPPLIED/STORAGE AND HANDLING

TALZENNA is supplied in strengths and package configurations as described in Table 12:

Table 12. TALZENNA Capsules

Package Configuration	Capsule Strength (mg)	NDC	Print
Bottles of 30 capsules	0.1	0069-0252-30	Red oval liquid-filled soft gelatin capsule printed with "TLZ" and "010".
Bottles of 30 capsules	0.25	0069-0353-30	Yellow oblong liquid-filled soft gelatin capsule printed with "TLZ" and "025".
Bottles of 30 capsules	0.35	0069-0454-30	Pink oblong liquid-filled soft gelatin capsule printed with "TLZ" and "035".
Bottles of 30 capsules	0.5	0069-0546-30	Pale orange oblong liquid-filled soft gelatin capsule printed with "TLZ" and "050".
Bottles of 30 capsules	0.75	0069-0655-30	Red oblong liquid-filled soft gelatin capsule printed with "TLZ" and "075".
Bottles of 30 capsules	1	0069-0757-30	Reddish brown oblong liquid-filled soft gelatin capsule printed with "TLZ" and "1".

Storage

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C to 30°C (59°F to 86°F). [See USP Controlled Room Temperature].

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

- **MDS/AML:** Advise patients to contact their healthcare provider if they experience weakness, feeling tired, fever, weight loss, frequent infections, bruising, bleeding easily, breathlessness, blood in urine or stool, and/or laboratory findings of low blood cell counts, or a need for blood transfusions. This may be a sign of

hematological toxicity or a more serious uncommon bone marrow problem called MDS or AML, which have been reported in patients who received PARP inhibitors [see *Warnings and Precautions (5.1)*].

- **Myelosuppression:** Advise patients that TALZENNA may affect hematopoiesis and can cause anemia, leukopenia/neutropenia, and/or thrombocytopenia [see *Warnings and Precautions (5.2)*].
- **Administration Instructions:** Advise patients that TALZENNA can be taken once daily with or without food. Instruct patients that if they miss a dose of TALZENNA, they should take their next normal dose at the usual time. Also advise patients to swallow each capsule whole, and that capsules must not be opened or dissolved [see *Dosage and Administration (2.4)*].
- **Embryo-Fetal Toxicity:** Advise females to inform their healthcare provider if they are pregnant or become pregnant. Inform female patients of the risk to a fetus and potential loss of the pregnancy [see *Use in Specific Populations (8.1)*]. Advise females of reproductive potential to use effective contraception during treatment with TALZENNA and for 7 months after the last dose. Advise male patients with female partners of reproductive potential or who are pregnant to use effective contraception during treatment and for 4 months after receiving the last dose of TALZENNA [see *Warnings and Precautions (5.3)*, *Use in Specific Populations (8.1, 8.3)*].
- **Lactation:** Advise patients not to breastfeed while taking TALZENNA and for 1 month after receiving the last dose [see *Use in Specific Populations (8.2)*].

This product's labeling may have been updated. For the most recent prescribing information, please visit <http://www.pfizer.com>.



Distributed by
Pfizer Labs
Division of Pfizer Inc.
New York, NY 10001

LAB-1561-2.0

PATIENT INFORMATION
TALZENNA® (Tal-ZEN-ah)
(talazoparib)
capsules

What is the most important information I should know about TALZENNA?
TALZENNA may cause serious side effects, including:

- **Bone marrow problems called Myelodysplastic Syndrome (MDS) or Acute Myeloid Leukemia (AML).** Some people who have cancer and who have received previous treatment with chemotherapy or certain other medicines for their cancer have developed MDS or AML during or after treatment with TALZENNA. MDS or AML may lead to death. If you develop MDS or AML, your healthcare provider will stop treatment with TALZENNA.

Symptoms of low blood cell counts are common during treatment with TALZENNA, but can be a sign of serious problems, including MDS or AML. Tell your healthcare provider if you have any of the following symptoms during treatment with TALZENNA:

- weakness
- weight loss
- fever
- frequent infections
- blood in urine or stool
- shortness of breath
- feeling very tired
- bruising or bleeding more easily

Your healthcare provider will do blood tests to check your blood cell counts:

- every month during treatment with TALZENNA.
- weekly if you have low blood cell counts that last a long time.

Your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with TALZENNA if you get certain side effects.

See “What are the possible side effects of TALZENNA?” below for other side effects of TALZENNA.

What is TALZENNA?

TALZENNA is a prescription medicine used:

- alone to treat adults with human epidermal growth factor receptor 2 (HER2)-negative breast cancer:
 - o who have a certain type of an abnormal inherited *BRCA* gene, **and**
 - o whose cancer has spread to other parts of the body (locally advanced or metastatic).

Your healthcare provider will perform a test to make sure that TALZENNA is right for you.

- in combination with a medicine called enzalutamide, to treat adults with prostate cancer
 - o with certain abnormal inherited or acquired genes called homologous recombination repair (HRR genes) **and**
 - o which no longer responds to hormone therapy or surgical treatment to lower testosterone and has spread to other parts of the body (metastatic).

Your healthcare provider will perform a test to make sure that TALZENNA is right for you.

It is not known if TALZENNA is safe and effective in children.

Before taking TALZENNA, tell your healthcare provider about all of your medical conditions, including if you:

- have kidney problems.
- are pregnant or plan to become pregnant. TALZENNA can harm your unborn baby and may cause loss of pregnancy (miscarriage). You should not become pregnant during treatment with TALZENNA.

Females who are able to become pregnant:

- o Your healthcare provider may do a pregnancy test before you start treatment with TALZENNA.
- o You should use effective birth control (contraception) during treatment with TALZENNA and for 7 months after the last dose of TALZENNA. Talk to your healthcare provider about forms of birth control that may be right for you.
- o Tell your healthcare provider right away if you are pregnant or think you are pregnant during treatment with TALZENNA.

Males with female partners who are pregnant or are able to become pregnant:

- o You should use effective birth control during treatment with TALZENNA and for 4 months after the last dose of TALZENNA.
- are breastfeeding or plan to breastfeed. It is not known if TALZENNA passes into your breast milk. Do not breastfeed during treatment with TALZENNA and for 1 month after the last dose of TALZENNA. Talk to your healthcare provider about the best way to feed your baby during this time.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking TALZENNA with certain other medicines can affect how TALZENNA works and may cause side effects.

Know the medicines you take. Keep a list of your medicines and show it to your healthcare provider and pharmacist when you get a new medicine.

How should I take TALZENNA?

- Take TALZENNA exactly as your healthcare provider tells you.
- Do not change your dose or stop taking TALZENNA without first talking with your healthcare provider.
- If you are prescribed TALZENNA for breast cancer, take TALZENNA 1 time a day.
- If you are prescribed TALZENNA for prostate cancer:
 - o Take TALZENNA with enzalutamide 1 time a day.
 - o You should start or continue a gonadotropin-releasing hormone (GnRH) analog therapy during your treatment with TALZENNA and enzalutamide unless you have had a surgery to lower the amount of testosterone in your body (surgical castration).
- Take TALZENNA with or without food.
- Swallow TALZENNA capsules whole. Do not dissolve or open TALZENNA capsules.
- If you vomit or miss a dose of TALZENNA, take your next dose at your regular time. Do not take an extra dose to make up for a missed dose.

What are the possible side effects of TALZENNA?

TALZENNA may cause serious side effects, including:

- **See “What is the most important information I should know about TALZENNA?”**

The most common side effects of TALZENNA when taken alone include:

- decreased red blood cell counts
- decreased white blood cell counts
- tiredness or weakness
- decreased platelet counts
- increased blood sugar levels
- increased liver function tests
- nausea
- headache
- decreased blood calcium levels
- vomiting
- hair loss
- diarrhea
- decreased appetite

The most common side effects of TALZENNA when taken in combination with enzalutamide include:

- decreased red blood cell counts
- decreased white blood cell counts
- tiredness or weakness
- decreased platelet counts
- decreased calcium in the blood
- nausea
- decreased appetite
- decreased sodium in the blood
- decreased phosphate in the blood
- bone injuries
- decreased magnesium in the blood
- dizziness
- increased bilirubin in the blood
- decreased potassium in the blood
- changes in your sense of taste

TALZENNA may cause fertility problems in males. This may affect your ability to father a child. Talk to your healthcare provider if this is a concern for you.

These are not all of the possible side effects of TALZENNA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store TALZENNA?

- Store TALZENNA at 68°F to 77°F (20°C to 25°C).

Keep TALZENNA and all medicines out of the reach of children.

General information about the safe and effective use of TALZENNA.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use TALZENNA for a condition for which it is not prescribed. Do not give TALZENNA to other people, even if they have the same symptoms that you have. It may harm them. You can ask your healthcare provider or pharmacist for information about TALZENNA that is written for health professionals.

What are the ingredients in TALZENNA?

Active ingredient: talazoparib tosylate

Inactive ingredients: gelatin, glycerol, iron oxide (red), iron oxide (yellow), polyethylene glycol 400, purified water, sorbitol, titanium dioxide, and tocopherol



Distributed by

Pfizer Labs

Division of Pfizer Inc.

New York, NY 10001

For more information, go to www.pfizer.com or call 1-800-438-1985.

LAB-1562-2.0

This Patient Information has been approved by the U.S. Food and Drug

PRINCIPAL DISPLAY PANEL - 0.1 mg Capsule Bottle Label

NDC 0069-0252-30

Pfizer

Talzenna[®]
(talazoparib) capsules

0.1 mg

30 Capsules

Rx only

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C to 30°C (59°F to 86°F).
Keep out of reach of children.
RECOMMENDED DOSAGE:
See Prescribing Information.
Each capsule contains 0.1 mg of talazoparib (equivalent to 0.145 mg talazoparib tosylate).
Distributed by Pfizer Labs
Division of Pfizer Inc.
New York, NY 10001
MADE IN GERMANY

Pfizer NDC 0069-0252-30

Talzenna[®]
(talazoparib) capsules

0.1 mg

30 Capsules **Rx only**

3 00690 25230 8
131674
GTIN: 00300690252308

Unvarnished Area for
3 lines of Variable Text
Font - 6pt OCRB

PRINCIPAL DISPLAY PANEL - 0.25 mg Capsule Bottle Label

NDC 0069-0353-30

Pfizer

Talzenna[®]
(talazoparib) capsules

0.25 mg

30 Capsules

Rx only

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C to 30°C (59°F to 86°F).

Keep out of reach of children.

RECOMMENDED DOSAGE:
See Prescribing Information.

Each capsule contains 0.25 mg of talazoparib (equivalent to 0.363 mg talazoparib tosylate).

Distributed by Pfizer Labs
Division of Pfizer Inc.
New York, NY 10001
MADE IN GERMANY

 NDC 0069-0353-30

Talzenna[®]
(talazoparib) capsules

0.25 mg

30 Capsules **Rx only**


3 00690 35330 2
131675
GTIN: 00300690353302

Unvarnished Area for
3 lines of Variable Text
Font - 6pt OCRB

PRINCIPAL DISPLAY PANEL - 0.35 mg Capsule Bottle Label

NDC 0069-0454-30

Pfizer

Talzenna[®]
(talazoparib) capsules

0.35 mg

30 Capsules

Rx only

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C to 30°C (59°F to 86°F).

Keep out of reach of children.

RECOMMENDED DOSAGE:
See Prescribing Information.

Each capsule contains 0.35 mg of talazoparib (equivalent to 0.509 mg talazoparib tosylate).

Distributed by Pfizer Labs
Division of Pfizer Inc.
New York, NY 10001
MADE IN GERMANY

 NDC 0069-0454-30

Talzenna[®]
(talazoparib) capsules

0.35 mg

30 Capsules **Rx only**


3 00690 45430 6
131676
GTIN: 00300690454306

Unvarnished Area for
3 lines of Variable Text
Font - 6pt OCRB

PRINCIPAL DISPLAY PANEL - 0.5 mg Capsule Bottle Label

NDC 0069-0546-30

Pfizer

Talzenna[®]
(talazoparib) capsules

0.5 mg

30 Capsules

Rx only

<p>Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C to 30°C (59°F to 86°F). Keep out of reach of children. RECOMMENDED DOSAGE: See Prescribing Information. Each capsule contains 0.5 mg of talazoparib (equivalent to 0.727 mg talazoparib tosylate). Distributed by Pfizer Labs Division of Pfizer Inc. New York, NY 10001 MADE IN GERMANY</p>	<p>NDC 0069-0546-30  Talzenna[®] (talazoparib) capsules 0.5 mg 30 Capsules Rx only</p>	 <p>3 0069054630 8 131677 GTIN: 00300690546308</p>	<p>Unvarnished Area for 3 lines of Variable Text Font - 6pt OCRB</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------

PRINCIPAL DISPLAY PANEL - 0.75 mg Capsule Bottle Label

NDC 0069-0655-30

Pfizer

Talzenna[®]
(talazoparib) capsules

0.75 mg

30 Capsules

Rx only

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C to 30°C (59°F to 86°F).

Keep out of reach of children.

RECOMMENDED DOSAGE:
See Prescribing Information.

Each capsule contains 0.75 mg of talazoparib (equivalent to 1.09 mg talazoparib tosylate).

Distributed by Pfizer Labs
Division of Pfizer Inc.
New York, NY 10001
MADE IN GERMANY

 NDC 0069-0655-30

Talzenna[®]
(talazoparib) capsules

0.75 mg

30 Capsules **Rx only**


3 00690 65530 7
131678
GTIN: 00300690655307

Unvarnished Area for
3 lines of Variable Text
Font - 6pt OCRB

PRINCIPAL DISPLAY PANEL - 1 mg Capsule Bottle Label

NDC 0069-0757-30

Pfizer

Talzenna[®]
(talazoparib) capsules

1 mg

30 Capsules

Rx only

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C to 30°C (59°F to 86°F).

Keep out of reach of children.

RECOMMENDED DOSAGE:
See Prescribing Information.

Each capsule contains 1 mg of talazoparib (equivalent to 1.453 mg talazoparib tosylate).

Distributed by Pfizer Labs
Division of Pfizer Inc.
New York, NY 10001
MADE IN GERMANY

 NDC 0069-0757-30

Talzenna[®]
(talazoparib) capsules

1 mg

30 Capsules **Rx only**


3 00690 75730 8
131679
GTIN: 00300690757308

Unvarnished Area for
3 lines of Variable Text
Font - 6pt OCRB

TALZENNA

talazoparib capsule, liquid filled

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0069-0252
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TALAZOPARIB TOSYLATE (UNII: 02WK9U5NZC) (TALAZOPARIB - UNII:9QHX048FRV)	TALAZOPARIB	0.1 mg

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
GLYCERIN (UNII: PDC6A3C0OX)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
GELATIN TYPE A PORCINE (195 BLOOM) (UNII: 6EG4DCS5TJ)	
SORBITOL (UNII: 506T60A25R)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	RED	Score	no score
Shape	OVAL	Size	9mm
Flavor		Imprint Code	TLZ;010
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0069-0252-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	04/17/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA217439	04/17/2024	

TALZENNA

talazoparib capsule, liquid filled

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0069-0353
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TALAZOPARIB TOSYLATE (UNII: 02WK9U5NZC) (TALAZOPARIB - UNII:9QHX048FRV)	TALAZOPARIB	0.25 mg

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
GLYCERIN (UNII: PDC6A3C0OX)	
GELATIN TYPE A PORCINE (195 BLOOM) (UNII: 6EG4DCS5TJ)	
SORBITOL (UNII: 506T60A25R)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	YELLOW	Score	no score
Shape	OVAL (Oblong)	Size	11mm
Flavor		Imprint Code	TLZ;025
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0069-0353-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	04/17/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA217439	04/17/2024	

TALZENNA

talazoparib capsule, liquid filled

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0069-0454
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TALAZOPARIB TOSYLATE (UNII: 02WK9U5NZC) (TALAZOPARIB - UNII:9QHX048FRV)	TALAZOPARIB	0.35 mg

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
GLYCERIN (UNII: PDC6A3C0OX)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
GELATIN TYPE A PORCINE (195 BLOOM) (UNII: 6EG4DCS5TJ)	
SORBITOL (UNII: 506T60A25R)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	PINK	Score	no score
Shape	OVAL (Oblong)	Size	14mm
Flavor		Imprint Code	TLZ;035
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0069-0454-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	04/17/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA217439	04/17/2024	

TALZENNA

talazoparib capsule, liquid filled

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0069-0546
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TALAZOPARIB TOSYLATE (UNII: 02WK9U5NZC) (TALAZOPARIB - UNII:9QHX048FRV)	TALAZOPARIB	0.5 mg

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
GLYCERIN (UNII: PDC6A3C0OX)	
GELATIN TYPE A PORCINE (195 BLOOM) (UNII: 6EG4DCS5TJ)	
SORBITOL (UNII: 506T60A25R)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WATER (UNII: 059QF0KO0R)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

Product Characteristics

Color	ORANGE (Pale Orange)	Score	no score
Shape	OVAL (Oblong)	Size	17mm
Flavor		Imprint Code	TLZ;050
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0069-0546-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	04/17/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA217439	04/17/2024	

TALZENNA

talazoparib capsule, liquid filled

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0069-0655
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TALAZOPARIB TOSYLATE (UNII: 02WK9U5NZC) (TALAZOPARIB - UNII:9QHX048FRV)	TALAZOPARIB	0.75 mg

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
GLYCERIN (UNII: PDC6A3C0OX)	
GELATIN TYPE A PORCINE (195 BLOOM) (UNII: 6EG4DCS5TJ)	
SORBITOL (UNII: 506T60A25R)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	RED	Score	no score
Shape	OVAL (Oblong)	Size	19mm
Flavor		Imprint Code	TLZ;075
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0069-0655-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	04/17/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA217439	04/17/2024	

TALZENNA

talazoparib capsule, liquid filled

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0069-0757
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TALAZOPARIB TOSYLATE (UNII: 02WK9U5NZC) (TALAZOPARIB - UNII:9QH048FRV)	TALAZOPARIB	1.0 mg

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
GLYCERIN (UNII: PDC6A3C0OX)	
GELATIN TYPE A PORCINE (195 BLOOM) (UNII: 6EG4DCS5TJ)	
SORBITOL (UNII: 506T60A25R)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	BROWN (Reddish brown)	Score	no score
Shape	OVAL (Oblong)	Size	22mm
Flavor		Imprint Code	TLZ;1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0069-0757-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	04/17/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA217439	04/17/2024	

Labeler - Pfizer Laboratories Div Pfizer Inc (134489525)

Establishment

Name	Address	ID/FEI	Business Operations
Pfizer Ireland Pharmaceuticals Unlimited Company		985052076	API MANUFACTURE(0069-0252, 0069-0353, 0069-0454, 0069-0546, 0069-0655, 0069-0757)

Establishment

Name	Address	ID/FEI	Business Operations
Eurofins Biolab Srl		429117112	ANALYSIS(0069-0252, 0069-0353, 0069-0454, 0069-0546, 0069-0655, 0069-0757)

Revised: 6/2025

Pfizer Laboratories Div Pfizer Inc