

MOTIONCALM- anamirta cocculus seed, ipecac,strychnos nux-vomica seed,tobacco leaf, kerosene tablet

Boiron

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

MotionCalm

Active ingredients** (in each tablet)

Cocculus indicus 4C HPUS (0.5 mg) (contains less than 10-10 mg picrotoxinin)

Ipecacuanha 4C HPUS (0.5 mg) (contains less than 10-10 mg emetine alkaloids)

Nux vomica 4C HPUS (0.5 mg) (contains less than 10-10 mg strychnine alkaloids)

Tabacum 4C HPUS (0.5 mg) (contains less than 10-10 mg nicotine alkaloids)

Petroleum 4C HPUS (0.5 mg) (contains less than 10-10 mg aromatic hydrocarbons)

The letters "HPUS" indicate that the components in this product are officially monographed in the Homeopathic Pharmacopoeia of the United States.

Purpose*

Cocculus indicus 4C HPUS Relieves nausea or vomiting with dizziness and the need to lie down

Ipecacuanha 4C HPUS Relieves nausea and vomiting with hypersalivation

Nux vomica 4C HPUS Relieves upset stomach, indigestion, nausea, and vomiting

Tabacum 4C HPUS Relieves dizziness, nausea, and vomiting improved by fresh air

Petroleum 4C HPUS Relieves nausea improved by eating

Uses*

for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness due to: travel, amusement rides, video games

Stop use and ask a doctor if symptoms persist for more than 2 days or worsen.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Adults and children 7 years of age and older

- To prevent motion sickness, dissolve 2 tablets in the mouth on the evening before travel and day of travel.
- To treat motion sickness, dissolve 2 tablets in the mouth at the onset of symptoms, then repeat if necessary until symptoms improve, up to 3 times per day.

Children 2 to 6 years of age

- Dissolve 2 tablets in 1 tablespoon of water for easier intake. Follow directions above.

Children under 2 years of age

- Ask a doctor.

croscarmellose sodium, lactose, magnesium stearate

do not use if glued carton end flaps are open or if the blister seal is broken.

store below 86°F (30°C)

60 meltaway tablets unflavored

Motion Sickness*

Nausea Vomiting Dizziness*

Prevents & Relieves Symptoms

Non-Drowsy

*CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE, NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.

**C,K,CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.

BoironUSA.com

info@Boiron.com

1-800-BOIRON-1

(1-800-264-7661)

Made in France

Distributed by Boiron, Inc.

Newtown Square, PA 19073

Drug Facts

Active ingredients* (in each tablet)

Cocculus indicus 4C HPUS (0.5 mg) (contains less than 10 ⁻¹² mg picrotoxinin)	Relieves nausea or vomiting with dizziness and the need to lie down	Purpose*
Ipecacuanha 4C HPUS (0.5 mg) (contains less than 10 ⁻¹² mg emetine alkaloids)	Relieves nausea and vomiting with hypersalivation	
Nux vomica 4C HPUS (0.5 mg) (contains less than 10 ⁻¹² mg strychnine alkaloids)	Relieves upset stomach, indigestion, nausea, and vomiting	
Tabacum 4C HPUS (0.5 mg) (contains less than 10 ⁻¹² mg nicotine alkaloids)	Relieves dizziness, nausea, and vomiting improved by fresh air	
Petroleum 4C HPUS (0.5 mg) (contains less than 10 ⁻¹² mg aromatic hydrocarbons)	Relieves nausea improved by eating	

The letters "HPUS" indicate that the components in this product are officially monographed in the Homeopathic Pharmacopoeia of the United States.

Uses* for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness due to: ■ travel ■ amusement rides ■ video games

Warnings

Stop use and ask a doctor if symptoms persist for more than 2 days or worsen.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Adults and children 7 years of age and older	■ To prevent motion sickness, dissolve 2 tablets in the mouth on the evening before travel and the day of travel.
	■ To treat motion sickness, dissolve 2 tablets in the mouth at the onset of symptoms, then repeat if necessary until symptoms improve, up to 3 times per day.

CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE, NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED. *C, K, CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.

Drug Facts (continued)

Directions (continued)

Children 2 to 6 years of age	Dissolve 2 tablets in 1 tablespoon of water for easier intake. Follow directions above.
Children under 2 years of age	Ask a doctor.

Other information

- do not use if glued carton end flaps are open or if the blister seal is broken
- store below 86°F (30°C)

Inactive ingredients

croscarmellose sodium, lactose, magnesium stearate

Questions or comments?

BoironUSA.com
Info@Boiron.com
1-800-BOIRON-1
(1-800-264-7661)

Made in France
Distributed by Boiron Inc.
Newtown Square, PA 19073



NDC 0220-9234-04

MOTION SICKNESS*

Nausea • Vomiting
Dizziness*



MotionCalm[®] MELTAWAY TABLETS

HOMEOPATHIC MEDICINE

- Prevents & Relieves Symptoms
- Non-Drowsy



BOIRON[®]

60
MELTAWAY
TABLETS
Unflavored

World Leader in
Homeopathic Medicines

MOTION SICKNESS MotionCalm[®]

Adults and children 7 years of age and older: To prevent motion sickness, dissolve 2 tablets in the mouth on the evening before travel and the day of travel. To treat motion sickness, dissolve 2 tablets in the mouth at the onset of symptoms, then repeat if necessary until symptoms improve, up to 3 times per day. Children 2 to 6 years of age: Dissolve 2 tablets in 1 tablespoon of water for easier intake. Follow directions above. Children under 2 years of age: Ask a doctor.

BOIRON[®]

MOTION SICKNESS MotionCalm[™]

Adults and children 7 years of age and older: To prevent motion sickness, dissolve 2 tablets in the mouth on the evening before travel and the day of travel. To treat motion sickness, dissolve 2 tablets in the mouth at the onset of symptoms, then repeat if necessary until symptoms improve, up to 3 times per day. Children 2 to 6 years of age: Dissolve 2 tablets in 1 tablespoon of water for easier intake. Follow directions above. Children under 2 years of age: Ask a doctor.

Drug Facts

Active ingredients* (in each tablet)

Cocculus indicus 4C HPUS (0.5 mg) (contains less than 10 ⁻¹² mg picrotoxinin)	Relieves nausea or vomiting with dizziness and the need to lie down	Purpose*
Ipecacuanha 4C HPUS (0.5 mg) (contains less than 10 ⁻¹² mg emetine alkaloids)	Relieves nausea and vomiting with hypersalivation	
Nux vomica 4C HPUS (0.5 mg) (contains less than 10 ⁻¹² mg strychnine alkaloids)	Relieves upset stomach, indigestion, nausea, and vomiting	
Tabacum 4C HPUS (0.5 mg) (contains less than 10 ⁻¹² mg nicotine alkaloids)	Relieves dizziness, nausea, and vomiting improved by fresh air	
Petroleum 4C HPUS (0.5 mg) (contains less than 10 ⁻¹² mg aromatic hydrocarbons)	Relieves nausea improved by eating	

The letters "HPUS" indicate that the components in this product are officially monographed in the Homeopathic Pharmacopoeia of the United States.

Uses* for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness due to: ■ travel ■ amusement rides ■ video games

Warnings

Stop use and ask a doctor if symptoms persist for more than 2 days or worsen.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Adults and children 7 years of age and older	■ To prevent motion sickness, dissolve 2 tablets in the mouth on the evening before travel and the day of travel.
	■ To treat motion sickness, dissolve 2 tablets in the mouth at the onset of symptoms, then repeat if necessary until symptoms improve, up to 3 times per day.

CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE, NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED. *C, K, CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.

Drug Facts (continued)

Directions (continued)

Children 2 to 6 years of age	Dissolve 2 tablets in 1 tablespoon of water for easier intake. Follow directions above.
Children under 2 years of age	Ask a doctor.

Other information

- do not use if glued carton end flaps are open or if the blister seal is broken
- store below 86°F (30°C)

Inactive ingredients

croscarmellose sodium, lactose, magnesium stearate

Questions or comments?

BoironUSA.com
Info@Boiron.com
1-800-BOIRON-1
(1-800-264-7661)

Made in France
Distributed by Boiron Inc.
Newtown Square, PA 19073



NDC 0220-9234-04

MOTION SICKNESS*

Nausea • Vomiting
Dizziness*



MotionCalm[™] MELTAWAY TABLETS

HOMEOPATHIC MEDICINE

- Prevents & Relieves Symptoms
- Non-Drowsy



BOIRON[®]

60
MELTAWAY
TABLETS
Unflavored

World Leader in
Homeopathic Medicines



Meltaway Tablets Require
No Food or Water
No Artificial Colors,
Flavors, Sweeteners,
or Preservatives
No Known Drug
Interactions

The Boiron Promise

Your health deserves the greatest respect. That's why we're committed to your satisfaction. Please visit BoironUSA.com/promise for details.



MOTIONCALM

anamirta cocculus seed, ipecac,strychnos nux-vomica seed,tobacco leaf, kerosene tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0220-9234
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IPECAC (UNII: 62I3C8233L) (IPECAC - UNII:62I3C8233L)	IPECAC	4 [hp_C]
STRYCHNOS NUX-VOMICA SEED (UNII: 269XH13919) (STRYCHNOS NUX-VOMICA SEED - UNII:269XH13919)	STRYCHNOS NUX-VOMICA SEED	4 [hp_C]
KEROSENE (UNII: 1C89KKC04E) (KEROSENE - UNII:1C89KKC04E)	KEROSENE	4 [hp_C]
ANAMIRTA COCCULUS SEED (UNII: 810258W28U) (ANAMIRTA COCCULUS SEED - UNII:810258W28U)	ANAMIRTA COCCULUS SEED	4 [hp_C]
TOBACCO LEAF (UNII: 6YR2608RSU) (TOBACCO LEAF - UNII:6YR2608RSU)	TOBACCO LEAF	4 [hp_C]

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	Boiron
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0220-9234-04	1 in 1 CARTON	07/23/2021	
1		60 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved homeopathic		07/23/2021	

Labeler - Boiron (282560473)

Registrant - Boiron, Inc. (014892269)

Establishment

Name	Address	ID/FEI	Business Operations
Boiron		282560473	manufacture(0220-9234)

Revised: 5/2023

Boiron