

RECHARGE PROTECT BLUR- titanium dioxide, zinc oxide cream
Rodan & Fields

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active Ingredients

Titanium Dioxide 3.2%

Zinc Oxide 7.7%

Purpose

Sunscreen

Sunscreen

Uses

- Helps prevent sunburn.
- If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

- **For external use only**
- **Do not use** on damaged or broken skin
- **When using this product** keep out of eyes. Rinse with water to remove.
- **Stop use and ask a doctor** if rash occurs.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- After moisturizing, apply generously at least 15 minutes before sun exposure using fingertips. Smooth and blend evenly over entire face in an outward direction.
- Wear alone or under makeup to prime skin.
- Reapply at least every 2 hours.
- Use a water-resistant sunscreen if swimming or sweating
- Children under 6 months of age: ask a doctor.
- **Sun Protection Measures:**
Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
 - Limit time in the sun, especially from 10 a.m.–2 p.m.
 - Wear long-sleeve shirts, pants, hats and sunglasses.

Other Information

Protect the product in this container from excessive heat and direct sun.

Inactive Ingredients

Isododecane, Cyclopentasiloxane, Dimethicone Crosspolymer, Calcium Sodium Borosilicate, Butyloctyl Salicylate, Jojoba Esters, Dimethicone, Glycerin, Sea Water/Maris Aqua/Eau De Mer, Hexyl Laurate, Acrylates/Dimethicone Copolymer, Water, Vaccinium Angustifolium (Blueberry) Fruit Extract, Carnosine, Sclareolide, Zinc PCA, Fragrance/Parfum, Silica, Dimethicone/Polyglycerin-3 Crosspolymer, Polyglyceryl-4 Isostearate, PEG-10 Dimethicone, Cetyl PEG/PPG-10/1 Dimethicone, Polyglyceryl-3 Polydimethylsiloxyethyl Dimethicone, Triethoxysilylethyl Polydimethylsiloxyethyl Hexyl Dimethicone, Stearic Acid, Biosaccharide Gum-4, Methyl Carboxymethylphenyl Aminocarboxypropylphosphonate, Isopropyl Titanium Triisostearate, Tocopherol, Sodium Citrate, Butylene Glycol, Propanediol, Dipropylene Glycol, Phenoxyethanol, Hexyl Cinnamal, Limonene, Linalool, Alumina, Iron Oxides (CI 77499).

Principal Display Panel - 40 mL Carton Label

RODAN + FIELDS

Protect + Blur Broad

Spectrum SPF 30

Sunscreen

3

40 mL / 1.35 Fl. Oz. U.S.

R+F

RODAN + FIELDS

Drug Facts

Active Ingredients Purpose
Titanium Dioxide 3.2% } Sunscreen
Zinc Oxide 7.7% }

Uses

- Helps prevent sunburn.
- If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

- **For external use only**
- **Do not use** on damaged or broken skin
- **When using this product** keep out of eyes. Rinse with water to remove.
- **Stop use and ask a doctor** if rash occurs.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- After moisturizing, apply generously at least 15 minutes before sun exposure using fingertips. Smooth and blend evenly over entire face in an outward direction. • Wear alone or under makeup to prime skin.
- Reapply at least every 2 hours. • Use a water-resistant sunscreen if swimming or sweating • Children under 6 months of age: ask a doctor.
- **Sun Protection Measures:** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including: ▶

Drug Facts (continued)

- Limit time in the sun, especially from 10 a.m.–2 p.m. • Wear long-sleeve shirts, pants, hats and sunglasses.

Other Information

Protect the product in this container from excessive heat and direct sun.

Inactive Ingredients

Isododecane, Cyclopentasiloxane, Dimethicone Crosspolymer, Calcium Sodium Borosilicate, Butyloctyl Salicylate, Jojoba Esters, Dimethicone, Glycerin, Sea Water/Maris Aqua/Eau De Mer, Hexyl Laurate, Acrylates/Dimethicone Copolymer, Water, Vaccinium Angustifolium (Blueberry) Fruit Extract, Carnosine, Sclareolide, Zinc PCA, Fragrance/Parfum, Silica, Dimethicone/Polyglycerin-3 Crosspolymer, Polyglyceryl-4 Isostearate, PEG-10 Dimethicone, Cetyl PEG/PPG-10/1 Dimethicone, Polyglyceryl-3 Polydimethylsiloxyethyl Dimethicone, Triethoxysilylethyl Polydimethylsiloxyethyl Hexyl Dimethicone, Stearic Acid, Biosaccharide Gum-4, Methyl Carboxymethylphenyl Aminocarboxypropylphosphate, Isopropyl Titanium Triisostearate, Tocopherol, Sodium Citrate, Butylene Glycol, Propanediol, Dipropylene Glycol, Phenoxethanol, Hexyl Cinnamal, Limonene, Linalool, Alumina, Iron Oxides (CI 77499).

Step 3 in the RECHARGE Regimen for Balancing, Boosting and Defending Healthy-Looking Skin. Fast-absorbing lotion delivers lightweight, broad spectrum, UVA/UVB protection + blurs imperfections for a smooth, poreless-looking matte finish. The universally flattering formula easily massages into skin. May also be used as a makeup primer to prep your face for the day — with or without makeup. 100% mineral sunscreen is oil-free and non-comedogenic. Use as part of the RECHARGE Regimen for best results.

Developed by Stanford-trained Dermatologists Dr. Katie Rodan and Dr. Kathy Fields.

RECHARGE

Protect + Blur Broad Spectrum SPF 30 Sunscreen

3

40 mL / 1.35 Fl. Oz. U.S.

Dist. Rodan & Fields, LLC
San Francisco, CA 94105
Made in USA with US and Imported Materials
Please Recycle


Questions?
1-888-995-5656
rodanandfields.com

RCSP040 8000-12-0001
6000-12-0001 0A



LOT CODE AREA





Principal Display Panel - 40 mL Tube Label

RODAN + FIELDS

Protect + Blur Broad
Spectrum SPF 30
Sunscreen

3

RODAN+FIELDS

RECHARGE

Protect + Blur Broad Spectrum
SPF 30 Sunscreen

3

ACTIVE INGREDIENTS

Titanium Dioxide 3.2%
Zinc Oxide 7.7%

PURPOSE

100% Mineral UVA and UVB protection. Broad Spectrum SPF 30 protects skin while blurring the appearance of pores and fine lines. The oil-free formula virtually disappears into skin for a translucent, smooth matte finish.

DIRECTIONS

After moisturizing, apply generously at least 15 minutes before sun exposure. Smooth and blend evenly over entire face in an outward direction, avoiding the eye area. Wear alone or as a makeup primer to prep skin.

WARNINGS

For external use only. Do not use on damaged or broken skin. **When using this product** keep out of eyes. Rinse with water to remove. **Stop use and ask a doctor** if rash occurs. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Dist. Rodan & Fields, LLC
San Francisco, CA 94105

Questions

1-888-995-5656
rodanandfields.com

RCSP040 8500-12-0001 6000-12-0002 0A

40 mL / 1.35 Fl. Oz. U.S.

RECHARGE PROTECT BLUR

titanium dioxide, zinc oxide cream

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:14222-80 10
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.032 g in 1 mL
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	0.077 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
ISODODECANE (UNII: A8289P68Y2)	
CYCLOMETHICONE 5 (UNII: 0THT5PC10R)	
BUTYLOCTYL SALICYLATE (UNII: 2EH13UN8D3)	
HYDROGENATED JOJOBA OIL, RANDOMIZED (UNII: Q47ST02F58)	
DIMETHICONE (UNII: 92RU3N3Y10)	
GLYCERIN (UNII: PDC6A3C0OX)	
HEXYL LAURATE (UNII: 4CG9F9W01Q)	
WATER (UNII: 059QF0KO0R)	
BLUEBERRY (UNII: 253RUG1X1A)	
CARNOSINE (UNII: 8HO6PVN24W)	
SCLAREOLIDE (UNII: 37W400O6E6)	
ZINC PIDOLATE (UNII: C32PQ86DH4)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 3) (UNII: G300307ZXP)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
BIO SACCHARIDE GUM-4 (UNII: 9XRL057X90)	
ISOPROPYL TITANIUM TRIISOSTEARATE (UNII: 949E3KBJ1I)	
TOCOPHEROL (UNII: R0ZB2556P8)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PROPANEDIOL (UNII: 5965N8W85T)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
LIMONENE, (+)- (UNII: GFD7C86Q1W)	
LINALOOL, (+/-)- (UNII: D81QY6I88E)	
.BETA.-CITRONELLOL, (R)- (UNII: P01OUT964K)	
GERANIOL (UNII: L837108USY)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:14222-8010-1	1 in 1 CARTON	09/20/2019	
1		30 mL in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	09/20/2019	

