

IBUPROFEN- ibuprofen capsules 200 mg capsule, liquid filled
TIME CAP LABORATORIES, INC.

610T Timely 49483-133 Ibuprofen Capsules 200 mg

DRUG FACTS

Active ingredient (in each capsule)

Solubilized Ibuprofen equal to
200 mg ibuprofen (NSAID)*
(present as the free acid and potassium salt)
* nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

- temporarily relieves minor aches and pains due to :

headache
toothache
backache
menstrual cramps
the common cold
muscular aches
minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alerts: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:

feel faint

vomit blood

have bloody or black stools

have stomach pain that does not get better

- you have symptoms of heart problems or stroke:

chest pain

trouble breathing

weakness in one part or side of body

slurred speech

leg swelling

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a

Poison Control Center right away (1-800-222-1222).

Directions

do not take more than directed

the smallest effective dose should be use

adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist

- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor

children under 12 years: ask a doctor

OTHER INFORMATION

Other information

- **each capsule contains:** potassium 20 mg
- read all warnings and directions before use
- store at 20-25°C(68-77 °F)
- avoid excessive heat above 40°C(104 °F)

Inactive ingredients Ammonium hydroxide, FD&C green #3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, propylene glycol, potassium hydroxide, purified water, shellac, sorbitan monooleate, sorbitol sorbitan

Questions or comments? call **1-877-290-4008**

Adhesive Area

Drug Facts (continued)

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor

Other information ■ each capsule contains:

potassium 20 mg ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F)

Inactive ingredients ammonium hydroxide, FD&C green #3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shellac, sorbitan monoleate, sorbitol sorbitan

Questions or comments? Call 1-877-290-4008

Varnish Omit Area

Lot No.:
Exp. Date:



3
Time-Cap Labs, Inc
7 Michael Avenue Farmingdale,
NY 11735
610T 0923
Made in India

Drug Facts (continued under label)

This product is not manufactured or distributed by Haleon group of companies, owner of the registered trademark Advil® and Catalent Pharma Solutions, Inc. owner of the registered trademark Liqui-Gels®.

Distributed by:
Time-Cap Labs, Inc
7 Michael Avenue Farmingdale,
NY 11735
610T 0923
Made in India

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Uses ■ temporarily relieves minor aches and pains due to:
● headache ● toothache ● backache ● menstrual cramps
● common cold ● muscular aches ● minor pain of arthritis
■ temporarily reduces fever

Drug Facts

Active ingredient (in each capsule)
Solubilized ibuprofen equivalent to
200 mg ibuprofen (NSAID)* Pain reliever/
Fever reducer
(present as the free acid and potassium salt)
*nonsteroidal anti-inflammatory drug

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

NDC 49483-133-37

Compare to the active ingredient in Advil® Liqui-Gels®†



Ibuprofen
solubilized ibuprofen capsules, 200 mg /
pain reliever/fever reducer
(NSAID)

133

ACTUAL SIZE

160 SOFTGELS**
(*LIQUID-FILLED CAPSULES)

Adhesive Area

Drug Facts (continued)

Do not use ■ If you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ● chest pain ● trouble breathing ● weakness in one part or side of body ● slurred speech ● leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ If pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor

Other information ■ each capsule contains: potassium 20 mg ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F)

Inactive ingredients ammonium hydroxide, FD&C green no. 3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, polyethylene glycol, purified water, shellac, sorbitan monooleate, sorbitol, sorbitan

Questions? call 1-877-290-4008

NDC 49483-133-12

timely™ Compare to the active ingredient in Advil® Liqui-Gels®†

Ibuprofen
solubilized ibuprofen capsules, 200 mg / pain reliever/fever reducer (NSAID)



ACTUAL SIZE

120 SOFTGELS**
(*LIQUID-FILLED CAPSULES)

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, MISSING, BROKEN OR MISSING

Drug Facts

Purpose
Pain reliever/
Fever reducer

Active ingredient (in each capsule)
Solubilized ibuprofen (present as the potassium salt)

Uses
Temporarily relieves minor aches and pains due to:
● headache
● toothache
● cold or flu
● sore throat
● menstrual cramps
● muscle aches
● backache
● pain of arthritis

Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people with allergies to aspirin, salicylates, or other NSAIDs. Symptoms may include hives, facial swelling, rash, dizziness, or difficulty breathing. Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ● chest pain ● trouble breathing ● weakness in one part or side of body ● slurred speech ● leg swelling ● pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

Stomach bleeding warning: This product contains an anti-inflammatory that may increase the risk of stomach bleeding. The risk is higher if you use more than directed or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for a longer time than directed.

Drug Facts (continued under label)

This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, owner of the registered trademark Advil® and Cataplan Pharma Solutions, Inc. owner of the registered trademark Liqui-Gels®.

Manufactured by:
Time-Cap Labs, Inc.
17 Michael Avenue Farmingdale,
NY 11735 6107 0622
Made in India

Varnish Omit Area

Lot No.:
Exp. Date:

Adhesive Area

Drug Facts (continued)

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ● chest pain ● trouble breathing ● weakness in one part or side of body ● slurred speech ● leg swelling ● pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ If pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor

Other information ■ each capsule contains: potassium 20 mg ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F)

Inactive ingredients ammonium hydroxide, FD&C green #3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shellac, sorbitan monooleate, sorbitol, sorbitan

Questions? call 1-877-290-4008

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL UNDER THE BOTTLE CAP IS BROKEN OR MISSING

Drug Facts

Purpose
Pain reliever/
Fever reducer

Active ingredient (in each capsule)
Solubilized ibuprofen (present as the potassium salt)

Uses
Temporarily relieves minor aches and pains due to:
● headache
● toothache
● cold or flu
● sore throat
● menstrual cramps
● muscle aches
● backache
● pain of arthritis

Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people with allergies to aspirin, salicylates, or other NSAIDs. Symptoms may include hives, facial swelling, rash, dizziness, or difficulty breathing. Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ● chest pain ● trouble breathing ● weakness in one part or side of body ● slurred speech ● leg swelling ● pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

Stomach bleeding warning: This product contains an anti-inflammatory that may increase the risk of stomach bleeding. The risk is higher if you use more than directed or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued under label)

This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, owner of the registered trademark Advil® and Cataplan Pharma Solutions, Inc. owner of the registered trademark Liqui-Gels®.

Manufactured by:
Time-Cap Labs, Inc.
17 Michael Avenue Farmingdale,
NY 11735 6107 0622
Made in India

Varnish Omit Area

Lot No.:
Exp. Date:

NDC 49483-133-80

timely™ Compare to the active ingredient in Advil® Liqui-Gels®†

Ibuprofen
solubilized ibuprofen capsules, 200 mg / pain reliever/fever reducer (NSAID)



ACTUAL SIZE

80 SOFTGELS**
(*LIQUID-FILLED CAPSULES)

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL UNDER THE BOTTLE CAP IS BROKEN OR MISSING

Drug Facts

Purpose
Pain reliever/
Fever reducer

Active ingredient (in each capsule)
Solubilized ibuprofen (present as the potassium salt)

Uses
Temporarily relieves minor aches and pains due to:
● headache
● toothache
● cold or flu
● sore throat
● menstrual cramps
● muscle aches
● backache
● pain of arthritis

Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people with allergies to aspirin, salicylates, or other NSAIDs. Symptoms may include hives, facial swelling, rash, dizziness, or difficulty breathing. Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ● chest pain ● trouble breathing ● weakness in one part or side of body ● slurred speech ● leg swelling ● pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

Stomach bleeding warning: This product contains an anti-inflammatory that may increase the risk of stomach bleeding. The risk is higher if you use more than directed or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued under label)

This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, owner of the registered trademark Advil® and Cataplan Pharma Solutions, Inc. owner of the registered trademark Liqui-Gels®.

Manufactured by:
Time-Cap Labs, Inc.
17 Michael Avenue Farmingdale,
NY 11735 6107 0622
Made in India

Varnish Omit Area

Lot No.:
Exp. Date:

timely™

Ibuprofen

solubilized ibuprofen capsules, 200 mg /
pain reliever/fever reducer
(NSAID)

20 SOFTGELS**
(*LIQUID-FILLED CAPSULES)

NDC 49483-133-23

DO NOT USE IF UNDER BOTTLE CAP IMPRINTED WITH
"SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

**READ AND KEEP CARTON FOR COMPLETE
WARNINGS AND INFORMATION**

Uses

temporarily relieves minor aches and pains, due to: headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis

- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older have had stomach ulcers or serious bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing aspirin, naproxen, or others have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart

Drug Facts (continued under label)

40136802 Cxide No.: 60/DRUGS/515

Lot No.:

Exp. Date:

Varnish Omit Area

PER
HERE

Exp. Date:

attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use

ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: potassium 20 mg
- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Questions? call 1-877-290-4008

Distributed by:

Time-Cap Labs, Inc.
7 Michael Avenue Farmingdale,
NY 11735
610T 0622
Made in India

Adhesive Area

Varnish/Omit
Lot No.:
Exp Date:



FOR INFORMATION ONLY
This product is a combination of Varnish/Omit and Omit. It is not intended to be used as a substitute for Varnish/Omit or Omit. The use of this product is limited to the treatment of pain and inflammation. It is not intended for use in the treatment of other conditions. For more information, please contact your healthcare provider.

1549590005900

Drug Facts

Active ingredient
(in each capsule)

Solubilized ibuprofen equal to 200 mg the potent NSAID (present as the free acid and potassium salt) (monomer and anti-inflammatory drug)

Pur pose

Uses • temporary relief of minor aches and pains due to: • headache • toothache • backache • menstrual cramps • the common cold • muscular aches • minor pain of arthritis • temporarily reduces fever

Warnings • Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, rash, wheezing, • shock • skin reddening • dizziness • fainting • If an allergic reaction occurs, stop use and seek medical help right away. Symptoms include: wheezing, The product contains an NSAID. Warnings: causes severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs such as aspirin, prescription or nonprescription NSAIDs (naproxen, ibuprofen, naproxen, or others) • have 3 or more

Drug Facts (continued)

• alcoholic drinks every day while using this product • the more or for a longer time that alcohol is consumed, the more serious the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use • if you have ever had an allergic reaction to any other pain reliever/fever reducer • right before or after heart surgery

Ask a doctor or pharmacist before use if you are • under a doctor's care for any serious condition • taking aspirin for heart attack or stroke, because ibuprofen may decrease its benefit for aspirin • taking any other drug

Other information • each capsule contains 200 mg of ibuprofen • read all warnings and directions • keep out of reach of children • do not use if the seal is broken • see the back of the box for more information • see the back of the box for more information • see the back of the box for more information

Drug Facts (continued)

any of the following signs of stomach bleeding: • belching • vomiting • knee body or back • stools • have stomach pain that does not get better • you have symptoms of heart problems or stroke • chest pain • trouble breathing • weakness in one part or side of body • slurred speech • tingling or numbness • pain, redness or swelling in the joints • more than 3 days of fever • pain, redness or swelling in the joints • more than 3 days of fever • pain, redness or swelling in the joints • more than 3 days of fever

Directions • Do not take more than directed • If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions • Do not take more than directed • If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

NDC 49483-133-23

Compare to the active ingredient in Advil® Liqui-Gels®



Ibuprofen

solubilized ibuprofen capsules, 200 mg / pain reliever/fever reducer

(NSAID)

20 SOFTGELS** (MULTI-DOSAGE CAPSULES)



ACTUAL SIZE

Varnish/Print Omit Area

C6011R2

Adhesive Area

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery.

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke ■ you are taking a diuretic.

Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug.

When using this product ■ take with food or milk if stomach upset occurs.

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ● chest pain ● trouble breathing ● weakness in one part of side of body ● slurred speech ● leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later. In pregnancy, unless definitely directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor.

Other information ■ each capsule contains: potassium 20 mg ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F).

Inactive ingredients ammonium hydroxide, FD&C green #3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, silica, sorbitan monoleate, sorbitol sorbitan.

Questions or comments? call 1-877-290-4008

NDC 49483-133-20



Compare to the active ingredient in Advil® Liqui-Gels®††

Ibuprofen

solubilized ibuprofen capsules, 200 mg / pain reliever/fever reducer (NSAID)



ACTUAL SIZE

200 SOFTGELS** (**LIQUID-FILLED CAPSULES)

DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to Pain reliever/ Fever reducer
200 mg ibuprofen (NSAID)*Fever reducer
(present as the free acid and potassium salt)
* nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to: ● headache ● toothache ● backache ● menstrual cramps ● the common cold
- muscular aches ● minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Drug Facts (continued under label)

††This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, owner of the registered trademark Advil® and Catalent Pharma Solutions, Inc., owner of the registered trademark Liqui-Gels®.

Manufactured by:
Time-Cap Labs, Inc.
7 Michael Avenue
Farmingdale, NY 11735
MADE IN INDIA
810T 0622



3 49483 13320 2

Lot No.:

Exp. Date:

Varnish Omit Area

PEEL HERE

VALUE SIZE

NDC 49483-133-31

timely™ Compare to the Active Ingredient in Advil® Liqui-Gels®

Ibuprofen

IBUPROFEN CAPSULES, 200 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)

SOFTGELS

300 SOFTGELS**
(**LIQUID-FILLED CAPSULES)

DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each capsule)
Soubinized ibuprofen equval to 200 mg ibuprofen (NSAID) (present as the free acid and potassium salt) nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- rash
- blisters

Stop using ibuprofen immediately and get medical help right away if you experience the following symptoms:

Stomach bleeding warning: This soft gel product contains an NSAID, which may cause severe stomach bleeding during treatment. The chance is higher if you:

- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug (ibuprofen, naproxen, or others)
- take other drugs containing prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Purpose
Pain reliever/fever reducer

Directions
Adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist.
If pain or fever does not respond to 1 capsule, 2 capsules may be used.
Do not exceed 6 capsules in 24 hours, unless directed by a doctor.
Children under 12 years: ask a doctor.

Other information

- each capsule contains potassium 20 mg
- read all warnings and directions before use
- store at 20-25°C (68°-77°) ■ avoid excessive heat above 40°C (104°F)

Inactive ingredients Aluminum hydroxide, FD&C green no. 3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, silica, sorbitan monoleate, sorbitol sorbitan

Questions or comments? call 1-877-239-4008

Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
- taking any other drug

When using this product: ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have blood or black stools
 - vomit blood
- you have symptoms of heart problem or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
- taking any other drug

When using this product: ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have blood or black stools
 - vomit blood
- you have symptoms of heart problem or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
- taking any other drug

When using this product: ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have blood or black stools
 - vomit blood
- you have symptoms of heart problem or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. In case of overdose, get medical help or contact a Poison Control Center right away.

3 049483 133 31 18

Varnish Omit Area

Lot No.: Exp. Date:

FREE 10 DAY TRIAL

IBUPROFEN

ibuprofen capsules 200 mg capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49483-133
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
SHELLAC (UNII: 46N107B71O)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
AMMONIA (UNII: 5138Q19F1X)	
GELATIN (UNII: 2G86QN327L)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	green	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	133
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49483-133-31	300 in 1 BOTTLE; Type 0: Not a Combination Product	07/27/2021	
2	NDC:49483-133-23	1 in 1 CARTON	06/16/2022	
2		20 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:49483-133-80	80 in 1 BOTTLE; Type 0: Not a Combination Product	06/16/2022	
4	NDC:49483-133-12	120 in 1 BOTTLE; Type 0: Not a Combination Product	06/16/2022	
5	NDC:49483-133-20	200 in 1 BOTTLE; Type 0: Not a Combination Product	06/16/2022	
6	NDC:49483-133-37	160 in 1 BOTTLE; Type 0: Not a Combination Product	09/16/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079205	07/27/2021	

Labeler - TIME CAP LABORATORIES, INC. (037052099)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(49483-133)

Revised: 9/2023

TIME CAP LABORATORIES, INC.