# ARTHRITIS PAIN ACETAMINOPHEN- acetaminophen tablet TARGET CORPORATION

-----

# 699R Target 11673 673 Acetaminophen Extended Release Tablets USP, 650 mg

## DRUG FACTS

## Active ingredient (in each caplet)

Acetaminophen 650 mg

## Purpose

Pain reliever/fever reducer

# Uses

temporarily relieves minor aches and pains due to:

- minor pain of arthritis
- muscular aches
- backache
- premenstrual and menstrual cramps
- the common cold
- headche
- toothache

temporarily reduces fever

# Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

# Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

# Ask a doctor before use if you have liver disease

# Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

# Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

## Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

# Directions

# Do not take more than directed

See overdose warning

- take 2 caplets every 8 hours with water
- swallow whole; do not crush, chew, split or dissolve

adults

- do not take more than 6 caplets in 24 hours
- do not use for more than 10 days unless directed by a doctor

under 18 years of age ask a doctor

# Other information

- store between 20-25°C (68-77°F)
- The FDA approved Dissolution methods differ from USP

*Inactive ingredients* carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

# **Questions or comments?**

Call **1-877-290-4008** 

OR

Call **1-800-910-6874** 

Questions or comments? ( Contains No Aspirin	<b>igredients</b> magnesium si gelatinized star tin	Other information ■ store between 20-25°C (68- ■ The FDA approved dissolution methods differ from USP	of age	aduits  aduits aduits aduits aduits aduits aduits aduity a	Directions Do not take more than directed. See overdose warning	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Ouick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	Stop use and ask a doctor if pain gets worse or lasts more than 10 days lever gets worse or lasts more than 3 days new symptoms occur redness or swelling is present These could be signs of a serious condition.	Ask a doctor or pharmactst before use If you are taking the blood thinning drug warfarin	Ask a doctor before use if you have liver disease	<ul> <li>Do not use</li> <li>with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>If you are allergic to acetaminophen or any of the inactive ingredients in this product</li> </ul>	Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ bilsters ■ rash If a skin reaction occurs, stop use and seek medical help right away.
Call 1-800-910-6874	camauba wax, hydroxyethyl cellulose, tearate, microcrystalline cellulose, rch, sodium starch glycolate, titanium	■ store between 20-25°C (68-77°F) ution methods differ from USP		<ul> <li>take 2 caplets every 8 hours with water</li> <li>swallow whole; do not crush, chew, split, or dissolve</li> <li>do not take more than 6 caplets in 24 hours</li> <li>do not use for more than 10 days unless directed by a doctor</li> </ul>	than directed.	a health professional before use. Jose, get medical help or contact 1-800-222-1222), Oulck medical as for children even if you do not	an 10 days ian 3 days xondition.	use If you are taking the blood	e liver disease	etaminophen (prescription or sure whether a drug contains sharmadst. ien or any of the inactive	may cause severe skin reactions. ■ skin reddening ■ bilsters ■ rash op use and seek medical help right aw ay.







ARTHRITIS PAIN ACE acetaminophen tablet	TAMINOPHEN						
Product Information							
Product Type	HUMAN OTC DRUG	Item Code (Sou	urce)	NDC:116	73-673		
Route of Administration	ORAL						
Active Ingredient/Active Moiety							
Ingre	Basis of St	trength	Strength				
ACETAMINOPHEN (UNII: 36209ITL	ACETAMINOPHEN		650 mg				
Inactive Ingredients							

	Ingredient N	ame	Strength					
POVIDONE K30 (UNII: U725QWY32X)								
CARNAUBA V	CARNAUBA WAX (UNII: R12CBM0EIZ)							
HYDROXYET	HYL CELLULOSE (140 CPS AT 5%) (UNII	: 8136Y38GY5)						
STARCH, CO	RN (UNII: 08232NY3SJ)							
MAGNESIUM	MAGNESIUM STEARATE (UNII: 70097M6I30)							
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)								
TRIACETIN (UNII: XHX3C3X673)								
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)								
CELLULOSE,	, MICROCRYSTALLINE (UNII: OP1R32D61U	)						
SODIUM STA	ARCH GLYCOLATE TYPE A (UNII: H8AV0SC	)X4D)						
Product Characteristics								
Color	white (White to off white)	Score	no score					
	0.000							

Color	white (White to off white)	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	71
Contains			

# Packaging

#	ltem Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:11673-673- 26	1 in 1 CARTON	12/17/2021			
1		225 in 1 BOTTLE; Type 0: Not a Combination Product				
2	NDC:11673-673- 01	100 in 1 BOTTLE; Type 0: Not a Combination Product	11/30/2022			
Marketing Information						

· · ··································								
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date					
ANDA	ANDA215486	12/17/2021						

# Labeler - TARGET CORPORATION (006961700)

**Registrant -** TIME CAP LABORATORIES, INC. (037052099)

# EstablishmentNameAddressID/FEIBusiness OperationsMARKSANS PHARMA LIMITED925822975manufacture(11673-673)

Revised: 3/2024

TARGET CORPORATION