

**ANTACID ANTIGAS - aluminum hydroxide, magnesium hydroxide,
simethicone liquid
Retail Business Services, LLC.**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Careone Regular Strength Antacid Antigas Liquid

ACTIVE INGREDIENTS (in each 10 mL)

Aluminum hydroxide (equiv. to dried gel, USP) 400 mg
Magnesium hydroxide 400 mg
Simethicone 40 mg

PURPOSE

Antacid
Antacid
Antigas

USE(S)

relieves:

- acid indigestion
- heartburn
- sour stomach
- upset stomach and gas associated with these symptoms

WARNINGS

Do not take more than 80 mL in a 24-hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor.

ASK A DOCTOR BEFORE USE IF YOU HAVE

- kidney disease
- a magnesium-restricted diet

ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.

STOP USE AND ASK DOCTOR

if symptoms last more than two weeks

KEEP OUT OF REACH OF CHILDREN

.

DIRECTIONS

- **shake well before use**
- adults and children 12 years and older: take 10 mL - 20 mL four times a day, or as directed by a doctor
- children under 12 years: consult a doctor
- mL = milliliter

OTHER INFORMATION

- **each 10 mL contains:** magnesium 170 mg, sodium 5 mg
- store at controlled room temperature 20°C-25°C (68°-77°F)
- do not freeze

INACTIVE INGREDIENTS

ethyl alcohol, flavor, glycerin, hydroxyethyl cellulose, methylparaben, propylene glycol, propylparaben, purified water, saccharin sodium, simethicone emulsion, sorbitol

PRINCIPAL DISPLAY PANEL

Careone
NDC 72476-503-05

REGULAR STRENGTH

ANTACID ANTIGAS LIQUID

Aluminum Hydroxide 400mg

Magnesium Hydroxide 400mg

Simethicone 40mg


Mint Flavor

Relieves:

Acid Indigestion, Pressure & Bloating (Gas)

12 FL OZ (355 mL)

Alcohol content 0.4% v/v



DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each 10 mL)	Purpose
Aluminum hydroxide (equiv. to dried gel, USP) 400 mg	Antacid
Magnesium hydroxide 400 mg	Antacid
Simethicone 40 mg	Antigas

Uses relieves: ■ acid indigestion ■ heartburn ■ sour stomach ■ upset stomach and gas associated with these symptoms

Warnings
Do not take more than 80 mL in a 24-hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor.
Ask a doctor before use if you have ■ kidney disease ■ a magnesium-restricted diet
Ask a doctor or pharmacist before use if you are ■ presently taking a prescription drug. Antacids may interact with certain prescription drugs.
Stop use and ask a doctor if symptoms last more than two weeks
Keep out of reach of children.

Directions ■ shake well before use ■ adults and children 12 years and older: take 10 - 20 mL four times a day, or as directed by a doctor ■ children under 12 years: consult a doctor ■ mL = milliliter

Other information ■ each 10 mL contains: magnesium 170 mg, sodium 5 mg ■ store at 20°C-25°C (68°F-77°F) ■ do not freeze

Inactive ingredients ethyl alcohol, flavor, glycerin, hydroxyethyl cellulose, methylparaben, propylene glycol, propylparaben, purified water, saccharin sodium, simethicone emulsion, sorbitol.

DISTRIBUTED BY
FOODHOLD U.S.A., LLC
LANDOVER, MD 20785
1-877-846-9949

Quality guaranteed
or your money back.

MFR# 53041 REV 0421

UNWORNISHED AREA
FOR LOT & EXP
(LINE DOES NOT PRINT)

3 4 1520 00604 3

ANTACID ANTIGAS

aluminum hydroxide, magnesium hydroxide, simethicone liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72476-503
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IU0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IU0)	ALUMINUM HYDROXIDE	400 mg in 10 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM HYDROXIDE	400 mg in 10 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	40 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYDROXYETHYL CELLULOSE (2000 MPA.S AT 1%) (UNII: S38J6RZN16)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	WHITE	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72476-503-05	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/26/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part331	05/26/2021	

Labeler - Retail Business Services, LLC. (967989935)

Establishment

Name	Address	ID/FEI	Business Operations
Guardian Drug Company		119210276	MANUFACTURE(72476-503)

Revised: 5/2021

Retail Business Services, LLC.