

GOOD REMEDIES GENTLE LAXATIVE- bisacodyl tablet, coated
Strive Pharmaceuticals Inc.

GOOD REMEDIES™ GENTLE Laxative BISACODYL (USP) 5 mg

Drug Facts

Active ingredient (in each tablet)

Bisacodyl (USP) 5 mg

Purpose

Stimulant laxative

Use

• for relief of occasional constipation and irregularity • this product generally produces bowel movement in 6-12 hours

Warnings

Do not use • if you cannot swallow without chewing.

Ask a doctor before use if you have

- stomach pain, nausea or vomiting
- noticed a sudden change in bowel habits that lasts more than 2 weeks

When using this product

• it may cause stomach discomfort, faintness and cramps • do not chew or crush tablet(s) • do not use within 1 hour after taking an antacid or milk

Stop use and ask a doctor if

• you have rectal bleeding or fail to have a bowel movement after using this product. These could be signs of a serious condition. • you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
 - take with a glass of water
-

adults and children 12 years of age and over	1 to 3 tablets in a single daily dose. It is recommended to start with the lowest dose (1 tablet) increasing the next day as needed
children 6 to under 12 years of age	1 tablet in a single daily dose
children under 6 years	ask a doctor

Other information

- contains FD&C yellow no. 6
- store at 20°C- 25°C (68°-77°F)
- protect from excessive humidity

Inactive ingredients

acacia gum, calcium carbonate, croscarmellose sodium, D&C yellow #10 aluminum lake, dibasic calcium phosphate, FD&C yellow #6 aluminum lake, gelatin, glycerol monostearate, hypromellose, iron oxide red, magnesium stearate, methacrylic acid copolymer type C, microcrystalline cellulose, polyethylene glycol, precipitated silica, silicon dioxide, sodium methylparaben, sodium propylparaben, sodium starch glycolate, starch, sucrose, talcum, titanium dioxide, triethyl citrate

Questions or comments?

1-888-577-8033 Monday-Friday 8am - 4pm EST

Compare to the active ingredient of **Dulcolax[®]** Laxative Tablets*

Gentle, dependable constipation relief

*This product is not manufactured or distributed by Sanofi-Aventis Deutschland GMBH, owner of the registered trademark Dulcolax[®] Laxative Tablets.

READ AND KEEP OUTER CARTON FOR COMPLETE PRODUCT WARNINGS AND INFORMATION

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

DISTRIBUTED BY:

Strive Pharmaceuticals, Inc.
East Brunswick, NJ 08816

Product of India

Packaging



GENTLE Laxative
BISACODYL (USP) 5 mg
Stimulant Laxative



NDC 70692-167-52

\$1.25

GENTLE Laxative
BISACODYL (USP) 5 mg
Stimulant Laxative


Compare to the active ingredient of
Dulcolax® Laxative Tablets*

Gentle, dependable
constipation relief

25 TABLETS

Actual Size

DISTRIBUTED BY:
Strive Pharmaceuticals, Inc.
East Brunswick, NJ 08816
Product of India
CT77069216752 REV/01-022024



*This product is not manufactured or distributed by Sanofi-Aventis Deutschland GmbH, owner of the registered trademark Dulcolax® Laxative Tablets.

READ AND KEEP OUTER CARTON FOR COMPLETE PRODUCT WARNINGS AND INFORMATION
TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

NO VARNISH

LOT: EXP:

Drug Facts (continued)
magnesium stearate, methacrylic acid co-polymer type C, microcrystalline cellulose, polyethylene glycol, precipitated silica, silicon dioxide, sodium methylparaben, sodium propylparaben, talcum, titanium dioxide, triethyl citrate

Drug Facts (continued)
sodium starch glycolate, starch, sucrose, talcum, titanium dioxide, triethyl citrate

Use
for relief of occasional constipation and irregularity ■ this product generally produces bowel movement in 6-12 hours

Warnings
Do not use ■ if you cannot swallow without chewing.
Ask a doctor before use if you have ■ stomach pain, nausea or vomiting ■ noticed a sudden change in bowel habits that lasts more than 2 weeks

When using this product
■ it may cause stomach discomfort, flatness and cramps ■ do not chew or crush tablet(s) ■ do not use within 1 hour after taking an antacid or milk

Stop use and ask a doctor if
■ you have rectal bleeding or fail to have a bowel movement after using this product. These could be signs of a serious condition. ■ you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Other information
■ contains FD&C yellow no. 6 ■ store at 20°C-25°C (68°-77°F) ■ protect from excessive humidity


Inactive ingredients acacia gum, calcium carbonate, croscarmellose sodium, D&C yellow #10 aluminum lake, dibasic calcium phosphate, FD&C yellow #6 aluminum lake, gelatin, glycerol monostearate, hypromellose, iron oxide red

Directions
adults and children 12 years of age and over: 1 to 3 tablets in a single daily dose. It is recommended to start with the lowest dose (1 tablet) increasing the next day as needed
children 6 to under 12 years of age: 1 tablet in a single daily dose
children under 6 years: ask a doctor

Questions or comments?
1-888-577-8033 Monday-Friday 8am-4pm EST

K/O FOR GLUING

K/O FOR GLUING



DRUG FACTS

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Purpose</p> <p>Bisacodyl (USP) 5 mg.....Stimulant laxative</p>	<p>Drug Facts (continued)</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>						
<p>Use</p> <ul style="list-style-type: none"> for relief of occasional constipation and irregularity this product generally produces bowel movement in 6-12 hours 	<p>Directions</p> <ul style="list-style-type: none"> do not take more than directed take with a glass of water 						
<p>Warnings</p> <p>Do not use if you cannot swallow without chewing.</p>	<table border="1"> <tr> <td>adults and children 12 years of age and over</td> <td>1 to 3 tablets in a single daily dose. It is recommended to start with the lowest dose (1 tablet) increasing the next day as needed</td> </tr> <tr> <td>children 6 to under 12 years of age</td> <td>1 tablet in a single daily dose</td> </tr> <tr> <td>children under 6 years</td> <td>ask a doctor</td> </tr> </table>	adults and children 12 years of age and over	1 to 3 tablets in a single daily dose. It is recommended to start with the lowest dose (1 tablet) increasing the next day as needed	children 6 to under 12 years of age	1 tablet in a single daily dose	children under 6 years	ask a doctor
adults and children 12 years of age and over	1 to 3 tablets in a single daily dose. It is recommended to start with the lowest dose (1 tablet) increasing the next day as needed						
children 6 to under 12 years of age	1 tablet in a single daily dose						
children under 6 years	ask a doctor						
<p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> stomach pain, nausea or vomiting noticed a sudden change in bowel habits that lasts more than 2 weeks 	<p>Other information</p> <ul style="list-style-type: none"> contains FD&C yellow no. 6 store at 20°C-25°C (68°-77°F) protect from excessive humidity 						
<p>When using this product</p> <ul style="list-style-type: none"> it may cause stomach discomfort, faintness and cramps do not chew or crush tablet(s) do not use within 1 hour after taking an antacid or milk 	<p>Inactive ingredients acacia gum, calcium carbonate, croscarmellose sodium, D&C yellow #10 aluminum lake, dibasic calcium phosphate, FD&C yellow #6 aluminum lake, gelatin, glycerol monostearate, hypromellose, iron oxide red, ↓</p>						
<p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> you have rectal bleeding or fail to have a bowel movement after using this product. These could be signs of a serious condition. you need to use a laxative for more than 1 week 							
<p>If pregnant or breast-feeding, ask a health professional before use. →</p>							
<p>Drug Facts (continued)</p> <p>magnesium stearate, methacrylic acid co-polymer type C, microcrystalline cellulose, polyethylene glycol, precipitated silica, silicon dioxide, sodium methylparaben, sodium propylparaben, →</p>	<p>Drug Facts (continued)</p> <p>sodium starch glycolate, starch, sucrose, talcum, titanium dioxide, triethyl citrate</p> <p>Questions or comments?</p> <p>1-888-577-8033 Monday-Friday 8am-4pm EST</p>						

GOOD REMEDIES GENTLE LAXATIVE

bisacodyl tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70692-167
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
CALCIUM CARBONATE (UNII: H0G9379FGK)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)	
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
METHYLPARABEN SODIUM (UNII: CR6K9C2NHK)	
PROPYLPARABEN SODIUM (UNII: 625NNB0G9N)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
STARCH, CORN (UNII: O8232NY3SJ)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	G
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70692-167-52	1 in 1 CARTON	11/10/2023	
-		25 in 1 BLISTER PACK; Type 0: Not a Combination		

Product			
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	505G(a)(3)	11/10/2023	

Labeler - Strive Pharmaceuticals Inc. (080028013)

Revised: 1/2025

Strive Pharmaceuticals Inc.