

GOOD REMEDIES GENTLE LAXATIVE- bisacodyl tablet, coated
Strive Pharmaceuticals Inc.

GOOD REMEDIES™ GENTLE Laxative BISACODYL (USP) 5 mg

Drug Facts

Active ingredient (in each tablet)

Bisacodyl (USP) 5 mg

Purpose

Stimulant laxative

Use

• for relief of occasional constipation and irregularity • this product generally produces bowel movement in 6-12 hours

Warnings

Do not use • if you cannot swallow without chewing.

Ask a doctor before use if you have

- stomach pain, nausea or vomiting
- noticed a sudden change in bowel habits that lasts more than 2 weeks

When using this product

• it may cause stomach discomfort, faintness and cramps • do not chew or crush tablet(s) • do not use within 1 hour after taking an antacid or milk

Stop use and ask a doctor if

• you have rectal bleeding or fail to have a bowel movement after using this product. These could be signs of a serious condition. • you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
 - take with a glass of water
-

adults and children 12 years of age and over	1 to 3 tablets in a single daily dose. It is recommended to start with the lowest dose (1 tablet) increasing the next day as needed
children 6 to under 12 years of age	1 tablet in a single daily dose
children under 6 years	ask a doctor

Other information

- contains FD&C yellow no. 6
- store at 20°C- 25°C (68°-77°F)
- protect from excessive humidity

Inactive ingredients

acacia gum, calcium carbonate, croscarmellose sodium, D&C yellow #10 aluminum lake, dibasic calcium phosphate, FD&C yellow #6 aluminum lake, gelatin, glycerol monostearate, hypromellose, iron oxide red, magnesium stearate, methacrylic acid copolymer type C, microcrystalline cellulose, polyethylene glycol, precipitated silica, silicon dioxide, sodium methylparaben, sodium propylparaben, sodium starch glycolate, starch, sucrose, talcum, titanium dioxide, triethyl citrate

Questions or comments?

1-888-577-8033 Monday-Friday 8am - 4pm EST

Compare to the active ingredient of **Dulcolax[®]** Laxative Tablets*

Gentle, dependable constipation relief

*This product is not manufactured or distributed by Sanofi-Aventis Deutschland GMBH, owner of the registered trademark Dulcolax[®] Laxative Tablets.

READ AND KEEP OUTER CARTON FOR COMPLETE PRODUCT WARNINGS AND INFORMATION

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

DISTRIBUTED BY:

Strive Pharmaceuticals, Inc.
East Brunswick, NJ 08816

Product of India

Packaging



GOOD REMEDIES™

GENTLE

Laxative

BISACODYL (USP) 5 mg
Stimulant Laxative

NDC 70692-167-52

Compare to the active ingredient of
Dulcolax® Laxative Tablets*



GOOD REMEDIES™

GENTLE

Laxative

BISACODYL (USP) 5 mg
Stimulant Laxative

Gentle, dependable
constipation relief



Actual Size

25 TABLETS

LOT: _____

EXP: _____

DISTRIBUTED BY:
Strive Pharmaceuticals, Inc.
East Brunswick, NJ 08816

Product of India

CT7069216752 REV.00-072023

This product is not manufactured or distributed by Sanofi-Aventis Deutschland GmbH, owner of the registered trademark Dulcolax® Laxative Tablets.

READ AND KEEP OUTER CARTON FOR COMPLETE
PRODUCT WARNINGS AND INFORMATION

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING

3 70692 00097 9



*Compare to the active ingredient of Dulcolax® Laxative Tablets

NO VARNISH

Drug Facts (continued)

magnesium stearate, methacrylic acid co-polymer type C, microcrystalline cellulose, polyethylene glycol, precipitated silica, silicon dioxide, sodium methylparaben, sodium propylparaben.

Drug Facts (continued)

sodium starch glycolate, starch, sucrose, talcum, titanium dioxide, triethyl citrate

Questions or comments?

1-888-577-8033 Monday-Friday 8am-4pm EST

Other information

- contains FD&C yellow no. 6
- store at 20°C-25°C (68°-77°F)
- protect from excessive humidity

Inactive ingredients acacia gum, calcium carbonate, croscarmellose sodium, D&C yellow #10 aluminum lake, dibasic calcium phosphate, FD&C yellow #6 aluminum lake, gelatin, glycerol monostearate, hypromellose, iron oxide red

When using this product

that lasts more than 2 weeks

- stomach pain, nausea or vomiting
- noticed a sudden change in bowel habits

Ask a doctor before use if you have

- it may cause stomach discomfort,
- fatiness and cramps ■ do not chew or crush tablet(s) ■ do not use within 1 hour after taking an antacid or milk

Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Chewing.

- noticed a sudden change in bowel habits that lasts more than 2 weeks
- stomach pain, nausea or vomiting
- noticed a sudden change in bowel habits

When using this product

- it may cause stomach discomfort,
- fatiness and cramps ■ do not chew or crush tablet(s) ■ do not use within 1 hour after taking an antacid or milk

Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

start with the lowest dose (1 tablet) increasing the next day as needed

1 tablet in a single daily dose

children 6 to under 12 years of age

ask a doctor

children under 6 years

ask a doctor

K/O FOR GLUING

K/O FOR GLUING

Drug Facts	Drug Facts (continued)
Active ingredient (in each tablet) Bisacodyl (USP) 5 mg.....Stimulant laxative	Purpose Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Use ■ for relief of occasional constipation and irregularity ■ this product generally produces bowel movement in 6-12 hours	Directions ■ do not take more than directed ■ take with a glass of water
Warnings Do not use ■ if you cannot swallow without chewing.	Warnings Do not use ■ if you cannot swallow without chewing
Ask a doctor before use if you have ■ stomach pain, nausea or vomiting ■ noticed a sudden change in bowel habits that lasts more than 2 weeks	Use irregularly ■ this product generally produces bowel movement in 6-12 hours ■ for relief of occasional constipation and irregularity ■ this product generally produces bowel movement in 6-12 hours
When using this product ■ it may cause stomach discomfort, faintness and cramps ■ do not chew or crush tablet(s) ■ do not use within 1 hour after taking an antacid or milk	Do not use ■ if you cannot swallow without chewing
Stop use and ask a doctor if ■ you have rectal bleeding or fail to have a bowel movement after using this product. These could be signs of a serious condition. ■ you need to use a laxative for more than 1 week	adults and children 12 years of age and over 1 to 3 tablets in a single daily dose. It is recommended to start with the lowest dose (1 tablet) increasing the next day as needed
If pregnant or breast-feeding , ask a health professional before use. →	children 6 to under 12 years of age 1 tablet in a single daily dose
	children under 6 years ask a doctor
	Other information ■ contains FD&C yellow no. 6 ■ store at 20°C-25°C (68°-77°F) ■ protect from excessive humidity
	Inactive ingredients acacia gum, calcium carbonate, croscarmellose sodium, D&C yellow #10 aluminum lake, dibasic calcium phosphate, FD&C yellow #6 aluminum lake, gelatin, glycerol monostearate, hypromellose, iron oxide red. ↓

DRUG FACTS TABLE

Drug Facts	Drug Facts (continued)
Active ingredient (in each tablet) Bisacodyl (USP) 5 mg.....Stimulant laxative	Purpose Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Use ■ for relief of occasional constipation and irregularity ■ this product generally produces bowel movement in 6-12 hours	Directions ■ do not take more than directed ■ take with a glass of water
Warnings Do not use ■ if you cannot swallow without chewing.	Warnings Do not use ■ if you cannot swallow without chewing
Ask a doctor before use if you have ■ stomach pain, nausea or vomiting ■ noticed a sudden change in bowel habits that lasts more than 2 weeks	Use irregularly ■ this product generally produces bowel movement in 6-12 hours ■ for relief of occasional constipation and irregularity ■ this product generally produces bowel movement in 6-12 hours
When using this product ■ it may cause stomach discomfort, faintness and cramps ■ do not chew or crush tablet(s) ■ do not use within 1 hour after taking an antacid or milk	Do not use ■ if you cannot swallow without chewing
Stop use and ask a doctor if ■ you have rectal bleeding or fail to have a bowel movement after using this product. These could be signs of a serious condition. ■ you need to use a laxative for more than 1 week	adults and children 12 years of age and over 1 to 3 tablets in a single daily dose. It is recommended to start with the lowest dose (1 tablet) increasing the next day as needed
If pregnant or breast-feeding , ask a health professional before use. →	children 6 to under 12 years of age 1 tablet in a single daily dose
	children under 6 years ask a doctor
	Other information ■ contains FD&C yellow no. 6 ■ store at 20°C-25°C (68°-77°F) ■ protect from excessive humidity
	Inactive ingredients acacia gum, calcium carbonate, croscarmellose sodium, D&C yellow #10 aluminum lake, dibasic calcium phosphate, FD&C yellow #6 aluminum lake, gelatin, glycerol monostearate, hypromellose, iron oxide red. ↓

Drug Facts (continued)	Drug Facts (continued)
magnesium stearate, methacrylic acid co-polymer type C, microcrystalline cellulose, polyethylene glycol, precipitated	sodium starch glycolate, starch, sucrose, talcum, titanium dioxide, triethyl citrate
	Questions or comments?

silica, silicon dioxide, sodium
methylparaben, sodium propylparaben,



QUESTIONS OR COMMENTS?

1-888-577-8033 Monday-Friday 8am-4pm EST

GOOD REMEDIES GENTLE LAXATIVE

bisacodyl tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70692-167
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
CALCIUM CARBONATE (UNII: H0G9379FGK)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
METHYLPARABEN SODIUM (UNII: CR6K9C2NHK)	
PROPYLPARABEN SODIUM (UNII: 625NNB0G9N)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
STARCH, CORN (UNII: O8232NY3SJ)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	G

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70692-167-52	1 in 1 CARTON	11/10/2023	
1		25 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	505G(a)(3)	11/10/2023	

Labeler - Strive Pharmaceuticals Inc. (080028013)

Revised: 11/2023

Strive Pharmaceuticals Inc.