

TYLENOL EXTRA STRENGTH- acetaminophen tablet, film coated
Johnson & Johnson Consumer Inc.

TYLENOL Extra Strength

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or

pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 caplets every 6 hours while symptoms last▪ do not take more than 6 caplets in 24 hours, unless directed by a doctor▪ do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- store between 20-25°C (68-77°F)
- **do not use if blister unit is torn or broken**

Inactive ingredients

carnauba wax, corn starch, FD&C red no. 40 aluminum lake, hypromellose, magnesium stearate, powdered cellulose, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, titanium dioxide

Questions or comments?

Call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

PRINCIPAL DISPLAY PANEL

500mg

NDC 50580-457-11

To re-order reference the NDC Code

TYLENOL[®]

FOR ADULTS

Acetaminophen Pain Reliever - Fever Reducer

Extra Strength

100 Caplets

500 mg each

10 Blister Cards

with

10 Individual Blisters


SMART PACKAGING

DESIGNED FOR HEALTH

CARE PROFESSIONALS

FOR HOSPITAL AND GOVERNMENT USE ONLY

125616



TYLENOL
FOR ADULTS
Acetaminophen Pain Reliever - Fever Reducer
Extra Strength

500mg

NDC 50580-457-11

To re-order reference the NDC Code

TYLENOL
FOR ADULTS
Acetaminophen Pain Reliever - Fever Reducer
Extra Strength



100 Caplets
500 mg each

10 Blister Cards
with
10 Individual Blisters

500mg

Window

SMART PACKAGING
DESIGNED FOR HEALTH
CARE PROFESSIONALS

FOR HOSPITAL AND GOVERNMENT USE ONLY



NDC 50580-457-11


500mg


Blister pack
configuration for
ease of inventory

Clear dosing
information for
ease of identification

Perforation and
rounded corners for
easy dispensing

Scannable LOT/EXP
information on
bottom of carton





30049012/125616

500mg

TYLENOL[®]

FOR ADULTS

Acetaminophen Pain Reliever - Fever Reducer

Extra Strength

Drug Facts

Active ingredient (in each caplet)	Purpose
Acetaminophen 500 mg	Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - backache
 - toothache
 - premenstrual and menstrual cramps
- temporarily reduces fever
- headache
- minor pain of arthritis
- muscular aches

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Drug Facts (continued)

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none">take 2 caplets every 6 hours while symptoms lastdo not take more than 6 caplets in 24 hours, unless directed by a doctordo not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- store between 20-25°C (68-77°F)
- do not use if blister unit is torn or broken

Inactive ingredients

carnauba wax, corn starch, FD&C red no. 40 aluminum lake, hypromellose, magnesium stearate, powdered cellulose, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, titanium dioxide

Questions or comments?

Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Active ingredient made in India

Distributed by:

JOHNSON & JOHNSON CONSUMER INC.

McNeil Consumer Healthcare Division

Fort Washington, PA 19084 USA ©J&JCI 2020

Visit us at www.tylenol.com or call toll-free 1-877-TYLENOL (1-877-895-3665)

125616

TYLENOL EXTRA STRENGTH

acetaminophen tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50580-457
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name		Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		500 mg

Inactive Ingredients	
Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	white (WHITE WITH RED PRINT)	Score	no score
Shape	OVAL	Size	18mm
Flavor		Imprint Code	TYLENOL;500
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-457-11	10 in 1 CARTON	07/17/2020	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:50580-457-10	1 in 1 CARTON	07/10/2020	
2		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:50580-	700 in 1 BOTTLE, PLASTIC; Type 0: Not a	07/17/2020	

457-70	Combination Product	07/17/2020	
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/10/2020	

Labeler - Johnson & Johnson Consumer Inc. (878046358)

Revised: 1/2024

Johnson & Johnson Consumer Inc.