

PAIN RELIEF IBUPROFEN- ibuprofen tablet
Rite Aid Corporation

Rite Aid 44-292

Active ingredient (in each brown caplet)

Ibuprofen USP, 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - backache
 - menstrual cramps
 - toothache
 - headache
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- facial swelling
- shock
- asthma (wheezing)
- skin reddening
- hives
- blisters
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- are age 60 or older
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart

failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - slurred speech
 - leg swelling
 - trouble breathing
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist

- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- see end flap for expiration date and lot number

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

**RITE
AID
PHARMACY®**

****Compare to the active
ingredient of Advil® Caplets**

**pain relief
ibuprofen**

ibuprofen tablets USP, 200 mg
pain reliever • fever reducer
(NSAID)

ACTUAL SIZE

**100 COATED CAPLETS
(CAPSULE-SHAPED TABLETS)**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS
BROKEN OR MISSING**

****This product is not manufactured or distributed by Wyeth LLC, owner of the registered trademark
Advil® Caplets.**

50844 REV0618B29212

**DISTRIBUTED BY: RITE AID
30 HUNTER LANE, CAMP HILL, PA 17011**

44-292

PAIN RELIEF IBUPROFEN			
ibuprofen tablet			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-0292
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength

ITEM 0386501



TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING



Compare to the active ingredient in Advil® Caplets

pain relief ibuprofen

ibuprofen tablets USP, 200 mg
pain reliever • fever reducer
(NSAID)

50 COATED CAPLETS (CAPSULE-SHAPED TABLETS)

ACTUAL SIZE

DISTRIBUTED BY: RITE AID
30 HUNTER LANE, CAMP HILL, PA 17011

50844 REV0618829215
*This product is not manufactured or distributed by Wyeth LLC, owner of the registered trademark Advil® Caplets.

Monday-Friday
Call 1-800-426-9391 8:30 AM-4:00 PM ET

Questions or comments?
microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - leg swelling
 - slurred speech
- weakness in one part or side of body

Drug Facts (continued)

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **do not take more than directed**
- the **smallest effective dose should be used**
- adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- Store between 20-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- see end flap for expiration date and lot number

Inactive ingredients carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate,

Drug Facts (continued)

microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?
Call 1-800-426-9391 8:30 AM-4:00 PM ET
Monday-Friday

*This product is not manufactured or distributed by Wyeth LLC, owner of the registered trademark Advil® Caplets.
50844 REV0618829215
DISTRIBUTED BY: RITE AID
30 HUNTER LANE, CAMP HILL, PA 17011

Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredient
(in each brown caplet)
ibuprofen USP, 200 mg (NSAID) *nonsteroidal anti-inflammatory drug

Purpose
Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - menstrual cramps
 - the common cold
- headache
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- rash
- skin redness
- asthma (wheezing)
- facial swelling
- stock
- blisters
- hives

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older

take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg
---	-----------	--------

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STARCH, CORN (UNII: O8232NY3SJ)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Product Characteristics

Color	BROWN	Score	no score
Shape	OVAL	Size	14mm
Flavor		Imprint Code	44;292
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0292-8	1 in 1 BLISTER PACK	05/24/1988	
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:11822-0292-5	1 in 1 CARTON	05/24/1988	
2		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:11822-0292-2	1 in 1 CARTON	05/24/1988	
3		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:11822-0292-6	1 in 1 CARTON	05/24/1988	10/08/2021
4		16 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
5	NDC:11822-0292-7	1 in 1 CARTON	05/24/1988	10/08/2021
5		125 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
6	NDC:11822-0292-4	500 in 1 BOTTLE; Type 0: Not a Combination Product	05/24/1988	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	05/24/1988	

Labeler - Rite Aid Corporation (014578892)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(11822-0292)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(11822-0292)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	PACK(11822-0292)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	PACK(11822-0292)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	PACK(11822-0292)

Revised: 5/2020

Rite Aid Corporation