

COLD HEAD CONGESTION SEVERE- acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, coated

Harris Teeter

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

HTE - 1175 - 2019-1011

Drug Facts

<i>Active ingredients (in each caplet)</i>	<i>Purpose</i>
Acetaminophen 325 mg	Pain reliever/fever reducer
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- for the temporary relief of the following cold symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"> ▪ take 2 caplets every 4 hours ▪ swallow whole – do not crush, chew, or dissolve ▪ do not take more than 10 caplets in 24 hours
children under 12	<ul style="list-style-type: none"> ▪ ask a doctor

years |

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information

Inactive ingredients

acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, crospovidone, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

Questions or comments?

1-844-705-4384

PRINCIPAL DISPLAY PANEL

Harris Teeter

Cold & Head Congestion Severe

Compare to the Active Ingredients in Tylenol(R) Cold + Head Congestion Severe

Acetaminophen, Phenylephrine HCl, Guaifenesin

PAIN RELIEVER/FEVER REDUCER, NASAL DECONGESTANT, EXPECTORANT

Relieves

Headache, Body Aches, Fever & Sore Throat

Nasal Congestion

Mucus & Chest Congestion

For Adults

24 COOL TASTE CAPLETS

Actual Size

NC

NC

NC

Drug Facts (continued)

Warnings: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours.

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have ■ liver disease ■ diabetes ■ heart disease ■ high blood pressure ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland

Drug Facts (continued)

Directions ■ do not take more than directed (see overdose warning)

adults and children 12 years and over ■ take 2 caplets every 4 hours ■ swallow whole – do not crush, chew, or dissolve

children 6 to 11 years ■ do not take more than 10 caplets in 24 hours ■ ask a doctor

children under 6 years ■ ask a doctor

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Keep out of reach of children.

If pregnant or breast-feeding, ask a health professional before use. These could be signs of a serious condition.

- cough comes back or occurs with rash or headache that lasts
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

When using this product do not exceed recommended dosage

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Drug Facts (continued)

■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Drug Facts

Active Ingredients (in each caplet)

- Acetaminophen 325 mg.....Pain reliever/fever reducer
- Guaifenesin 200 mg.....Expectorant
- Phenylephrine HCl 5 mg.....Nasal decongestant

Purpose

Uses ■ for the temporary relief of the following cold symptoms:

- sore throat
- headache
- minor aches and pains
- nasal congestion
- sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Drug Facts (continued)

Cold & Head Congestion Severe

Acetaminophen, Phenylephrine HCl, Guaifenesin

**PAIN RELIEVER / FEVER REDUCER
NASAL DECONGESTANT, EXPECTORANT**

- Relieves
- Headache, Body Aches, Fever & Sore Throat
 - Nasal Congestion
 - Mucus & Chest Congestion
 - For Adults

24 COOL TASTE CAPLETS



Actual Size

†Compare to the Active Ingredients in Tylenol® Cold + Head Congestion Severe

Drug Facts (continued)

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information

Inactive ingredients acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, crospovidone, flavor, magnesium stearate, maltoedextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

Questions or comments? 1-844-705-4384

yourplusguarantee.
1-800-432-6111 or harris-teeter.com

PROUDLY DISTRIBUTED BY:
HARRIS TEETER, LLC
MATTHEWS, NC 28105

†This product is not manufactured or distributed by Mckel Consumer Healthcare, distributor of Tylenol® Cold + Head Congestion Severe.

DO NOT USE IF BLISTER UNITS ARE TORN OR BROKEN



F11750ZHE_RD

acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72036-175
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTO DEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	19mm
Flavor	MINT	Imprint Code	AAA;1119
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72036-175-02	2 in 1 CARTON	04/11/2017	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	04/11/2017	

Labeler - Harris Teeter (047279351)

Revised: 10/2019

Harris Teeter