

**WALGREEN DAYTIME COLD AND FLU RELIEF- acetaminophen,
dextromethorphan hydrobromide, guaifenesin and phenylephrine hcl liquid
WALGREENS CO.**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Walgreen DayQuil Severe Cold & Flu Honey

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifensin 200 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- minor aches & pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to clear the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
- child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persists or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product,

do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
 - pain, nasal congestion or cough gets worse or lasts more than 5 days (children) 7 days (adults)
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back, or occurs with rash or headache that lasts.
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over 30 mL every 4 hrs

children 6 to under 12 yrs 15 mL every 4 hrs

children 4 to under 6 yrs ask a doctor

children under 4 yrs do not use

Other information

- **each 15 mL contains:** sodium 9 mg
- Store at room temperature
- Do not refrigerate.

Inactive ingredients

citric acid, D&C Yellow No. 10, edetate disodium, FD&C Green No. 3, FD&C Red No. 40, flavors, glycerin, menthol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions?

1-866-467-2748

*This product is not manufactured or distributed by Procter & Gamble, distributor of

Vicks® Dayquil™ Severe Cold & Flu Honey Flavour

Distributed. by:

PRINCIPAL DISPLAY PANEL - 354 mL Bottle Label

*Compare to the active ingredients in Vicks® Dayquil™ Severe Cold & Flu Honey Flavour

NDC 0363-6110-12

DayTime

Cold & Flu

Relief

Acetaminophen - Pain reliever/Fever reducer

Guaifensin - Expectorant

Phenylephrine HCl - Nasal Congestion

Dextromethorphan HBr - Cough suppressant

- Headache, Fever, Sore Throat, Minor Aches & Pains
- Chest Congestion,
- Thins & Loosens Mucus
- Nasal Congestion, Sinus Pressure
- Cough

HONEY FLAVOR

Naturally and Artificially Flavored

12 FL OZ (354 ml)

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults) • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • take only as directed
 • only use the dose cup provided • do not exceed 4 doses per 24 hrs
 adults & children 12 yrs & over 30 mL every 4 hrs
 children 6 to under 12 yrs 15 mL every 4 hrs
 children 4 to under 6 yrs ask a doctor
 children under 4 yrs do not use

Other information • each 15 mL contains: sodium 9 mg • store at room temperature. Do not refrigerate.

Inactive ingredients citric acid, D&C Yellow No.10, edetate disodium, FD&C Green No. 3, FD&C Red No. 40, flavors, glycerin, menthol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum.

Questions or comments? 1-866-467-2748

*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® Dayquil™ Severe Cold & Flu Honey Flavor.

LOT: 61012XXLR Distributed by:

UNVARNISHED AREA

EXP:



*Compare to the active ingredients in Vicks® Dayquil™ Severe Cold & Flu Honey Flavor
NDC 0363-6110-12

DayTime Cold & Flu Relief

Acetaminophen - Pain reliever/Fever reducer
Guaifensin - Expectorant
Phenylephrine HCl - Nasal decongestant
Dextromethorphan HBr - Cough suppressant

- Headache, Fever, Sore Throat, Minor Aches & Pains
- Chest Congestion,
- Thins & Loosens Mucus
- Nasal Congestion, Sinus Pressure
- Cough

Honey Flavor
Naturally and Artificially Flavored
12 FL OZ (354 ml)

Drug Facts **DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN**

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Guaifensin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms: • nasal congestion • sinus congestion & pressure • cough due to minor throat & bronchial irritation • minor aches & pains • headache • fever • sore throat • reduces swelling of nasal passages
 • temporarily restores freer breathing through the nose • promotes nasal and/or sinus drainage • helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if • adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product • child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product
 • taken with other drugs containing acetaminophen • adult has 3 or more alcoholic drinks every day while using this product
Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 • Skin reddening • Blisters • Rash
 If a skin reaction occurs, stop use and seek medical help right away
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

WALGREEN DAYTIME COLD AND FLU RELIEF

acetaminophen, dextromethorphan hydrobromide, guaifenesin and phenylephrine hcl liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-6110
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
MENTHOL, UNSPECIFIED FORM (UNII: L7T10EIP3A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	brown	Score	
Shape		Size	
Flavor	HONEY	Imprint Code	
Contains			

Packaging

Marketing Start Marketing End

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-6110-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/08/2021	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part341	02/08/2021		

Labeler - WALGREENS CO. (008965063)

Revised: 2/2021

WALGREENS CO.