

**IBUPROFEN- ibuprofen tablet, film coated**  
**L.N.K. International, Inc.**

-----  
**Quality Plus 44-353**

***Active ingredient (in each white caplet)***

Ibuprofen USP, 200 mg (NSAID)\*  
\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - backache
  - menstrual cramps
  - the common cold
  - muscular aches
  - minor pain of arthritis
- temporarily reduces fever

***Warnings***

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- skin reddening
- shock
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - weakness in one part or side of body
  - leg swelling
  - slurred speech
  - trouble breathing
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- any new symptoms appear
- redness or swelling is present in the painful area

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen at 20

weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 caplet, 2 caplets may be used
  - do not exceed 6 caplets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

***Other information***

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- use by expiration date on package

***Inactive ingredients***

colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, sodium starch glycolate, stearic acid, titanium dioxide, triacetin

***Questions or comments?***

**Call 1-800-426-9391** 8:30 AM-4:00 PM ET, Monday-Friday

***Principal Display Panel***

**Quality  
+Plus**

NDC 50844-353-16

†Compare to active ingredient  
in Motrin® IB Caplets

**IBUPROFEN  
TABLETS  
USP, 200 mg**

**PAIN RELIEVER/FEVER REDUCER  
(NSAID)**

**1000 Coated Caplets\*\***

(\*\*CAPSULE-SHAPED TABLETS)

**ACTUAL SIZE**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Motrin® IB Caplets. 50844 REV1221C35316

Distributed by  
**LNK INTERNATIONAL, INC.**  
 60 Arkay Drive, Hauppauge, NY 11788  
 USA

NDC 50844-353-16

†Compare to active ingredient  
in Motrin® IB Caplets

**QUALITY PLUS**

**IBUPROFEN TABLETS USP, 200 mg**

**PAIN RELIEVER/FEVER REDUCER (NSAID)**

**1000 Coated Caplets\*\***  
(\*CAPSULE-SHAPED TABLETS)

ACTUAL SIZE

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts	Drug Facts (continued)
<p><b>Active ingredient (in each white caplet)</b></p> <p>Ibuprofen USP, 200 mg (NSAID) ..... Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug</p> <p><b>Purpose</b></p> <p>Pain reliever/fever reducer</p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>■ temporarily relieves minor aches and pains due to:</li> <li>■ headache    ■ toothache    ■ backache    ■ menstrual cramps</li> <li>■ the common cold    ■ muscular aches    ■ minor pain of arthritis</li> <li>■ temporarily reduces fever</li> </ul> <p><b>Warnings</b></p> <p><b>Allergy alert:</b> Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives    ■ facial swelling</p> <p>■ asthma (wheezing)    ■ skin reddening    ■ shock    ■ rash    ■ blisters</p> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older</p> <ul style="list-style-type: none"> <li>■ have had stomach ulcers or bleeding problems</li> <li>■ take a blood thinning (anticoagulant) or steroid drug</li> <li>■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]</li> <li>■ have 3 or more alcoholic drinks every day while using this product</li> <li>■ take more or for a longer time than directed</li> </ul> <p><b>Heart attack and stroke warning:</b> NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p><b>Do not use</b> ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer    ■ right before or after heart surgery</p> <p><b>Ask a doctor before use if</b> ■ stomach bleeding warning applies to you</p> <ul style="list-style-type: none"> <li>■ you have problems or serious side effects from taking pain relievers or fever reducers</li> <li>■ you have a history of stomach problems, such as heartburn</li> <li>■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke</li> <li>■ you are taking a diuretic</li> </ul> <p><b>Ask a doctor or pharmacist before use if you are</b> ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin</p> <ul style="list-style-type: none"> <li>■ under a doctor's care for any serious condition</li> <li>■ taking any other drug</li> </ul>	<p><b>When using this product</b> ■ take with food or milk if stomach upset occurs</p> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>■ you experience any of the following signs of stomach bleeding:</li> <li>■ feel faint    ■ vomit blood    ■ have bloody or black stools</li> <li>■ have stomach pain that does not get better</li> <li>■ you have symptoms of heart problems or stroke:</li> <li>■ chest pain    ■ weakness in one part or side of body    ■ trouble breathing</li> <li>■ slurred speech    ■ leg swelling</li> <li>■ pain gets worse or lasts more than 10 days</li> <li>■ fever gets worse or lasts more than 3 days    ■ any new symptoms appear</li> <li>■ redness or swelling is present in the painful area</li> </ul> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b> ■ do not take more than directed</p> <ul style="list-style-type: none"> <li>■ the smallest effective dose should be used</li> <li>■ adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist</li> <li>■ if pain or fever does not respond to 1 caplet, 2 caplets may be used</li> <li>■ do not exceed 6 caplets in 24 hours, unless directed by a doctor</li> <li>■ children under 12 years: ask a doctor</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ store between 20°-25°C (68°-77°F)</li> <li>■ avoid excessive heat 40°C (104°F)</li> <li>■ use by expiration date on package</li> </ul> <p><b>Inactive ingredients</b> colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, sodium starch glycolate, stearic acid, titanium dioxide, triacetin</p> <p><b>Questions or comments?</b> Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday</p>

0 50844 35316 9

†This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Motrin® IB Caplets. 50844 REV1221C35316

Distributed by  
**LNK INTERNATIONAL, INC.**  
 60 Arkay Drive, Hauppauge, NY 11788  
 USA

**Quality Plus 44-353**

<b>IBUPROFEN</b>			
ibuprofen tablet, film coated			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50844-353
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
	IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg
<b>Inactive Ingredients</b>			
	<b>Ingredient Name</b>		<b>Strength</b>

<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRACETIN</b> (UNII: XHX3C3X673)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	14mm
<b>Flavor</b>		<b>Imprint Code</b>	44;353
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-353-02	1 in 1 CARTON	03/01/1999	
1		12 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:50844-353-16	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/01/1999	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075139	03/01/1999	

**Labeler** - L.N.K. International, Inc. (038154464)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(50844-353)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(50844-353)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(50844-353)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(50844-353)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(50844-353)

Revised: 1/2026

L.N.K. International, Inc.