

**DAYTIME NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hbr, phenylephrine hcl and acetaminophen, dextromethorphan hbr, doxylamine succinate
WALGREENS**

695T 0363 0729 WALGREENS - Daytime Nighttime Cold & Flu

DRUG FACTS

Daytime Cold & Flu

Active ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Nighttime Cold & Flu

Active ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Daytime Cold & Flu

Purpose

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Nighttime Cold & Flu

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion (Daytime only)
- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing (Nighttime only)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4 doses in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease (Daytime only)
- high blood pressure (Daytime only)
- thyroid disease (Daytime only)
- diabetes (Daytime only)
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema (Daytime only)
- glaucoma (Nighttime only)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema (Nighttime only)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (Nighttime only)

When using this product

- **do not use more than directed** (Daytime only)
- excitability may occur, especially in children (Nighttime only)
- marked drowsiness may occur (Nighttime only)
- avoid alcoholic drinks (Nighttime only)
- be careful when driving a motor vehicle or operating machinery (Nighttime only)
- alcohol, sedatives, & tranquilizers may increase drowsiness (Nighttime only)

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless (Daytime only)
- pain, nasal congestion or cough get worse or last more than 7 days (Daytime only)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.
- pain, or cough gets worse or lasts more than 7 days (Nighttime only)
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 4 doses per 24 hrs

adults & children 12yrs & over: 2 softgels with water every 6 hrs(Nighttime only)

2 softgels with water every 4 hrs (Daytime only)

children 4 to under 12 yrs: ask a doctor

children under 4 yrs: do not use

OTHER INFORMATION**Other information**

- store at room temperature

Inactive ingredients (Daytime only)

FD&C red # 40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide.

Inactive ingredients (Nighttime only)

FD&C blue #1, FD&C yellow #10, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide.

Questions? Call 1-877-290-4008

Print/Varnish Omit Area

DAY & NIGHT PACK

Walgreens



Compare to the active ingredients in Vicks® DayQuil® & NyQuil® Cold & Flu LiquidCaps™

DAYTIME Cold & Flu

ACETAMINOPHEN 325 mg / ACHES / FEVER / SORE THROAT
DEXTROMETHORPHAN HBr 10 mg / COUGH
PHENYLEPHRINE HCl 5 mg / NASAL CONGESTION

Multi-Symptom

Relieves aches, fever, sore throat, nasal congestion & cough



ACTUAL SIZE

16 SOFTGELS

NIGHTTIME Cold & Flu

ACETAMINOPHEN 325 mg / ACHES / FEVER / SORE THROAT
DEXTROMETHORPHAN HBr 15 mg / COUGH
DOXYLAMINE SUCCONATE 6.25 mg / SNEEZING / RUNNING NOSE

Multi-Symptom

Relieves aches, fever, sore throat, cough, sneezing & runny nose



ACTUAL SIZE

8 SOFTGELS

24 TOTAL SOFTGELS

Varnish Omit Area

Varnish Omit Area

Drug Facts

Active Ingredients (in each softgel)

Acetaminophen 325 mg Pain reliever/fever reducer
Dextromethorphan HBr 10 mg Cough suppressant
Phenylephrine HCl 5 mg Nasal decongestant

Uses temporarily relieves common cold/flu symptoms: nasal congestion cough due to minor throat & bronchial irritation sore throat headache minor aches & pains fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses in 24 hrs, which is the maximum daily amount for this product with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include skin redness blisters rash If a skin reaction occurs, stop use and seek medical help right away

See blood warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly

Daytime Cold & Flu

Purpose

Pain reliever/fever reducer
Cough suppressant
Nasal decongestant

Drug Facts

Active Ingredients (in each softgel)

Acetaminophen 325 mg Pain reliever/fever reducer
Dextromethorphan HBr 15 mg Cough suppressant
Doxylamine succinate 6.25 mg Antihistamine

Uses temporarily relieves common cold/flu symptoms: cough due to minor throat & bronchial irritation sore throat headache minor aches & pains fever runny nose & sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses in 24 hrs, which is the maximum daily amount for this product with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include skin redness blisters rash If a skin reaction occurs, stop use and seek medical help right away

See blood warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly

Do not use with any other drug containing acetaminophen (prescription or

Drug Facts (continued)

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

Ask a doctor before use if you have liver disease heart disease high blood pressure thyroid disease diabetes trouble urinating due to enlarged prostate gland cough that occurs with too much phlegm (mucus) persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product, do not use more than directed

Stop use and ask a doctor if you get nervous, dizzy or sleepless pain, nasal congestion or cough get worse or last more than 7 days fever gets worse or lasts more than 3 days hoarseness or swelling is present new symptoms occur cough comes back, or occurs with rash or headache that lasts These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-422-1222) right away. Usual medical attention is critical for adults & for children even if you do not notice any signs or symptoms

Directions take only as directed do not exceed 4 doses per 24 hrs adults & children 12 yrs & over 2 softgels with water every 6 hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use

Other information store at room temperature

Inactive ingredients FD&C Red #40, FD&C Yellow #6, gelatin, glycerin, polyethylene glycol, polyethylene glycol, purified water, stearic acid, sorbitol sorbitan solidifier, titanium dioxide

Questions? Call 1-877-299-4888

Drug Facts (continued)

nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

Ask a doctor before use if you have liver disease glaucoma cough that occurs with too much phlegm (mucus) a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers taking the blood thinning drug warfarin

When using this product, drowsiness may occur especially in children marked drowsiness may occur avoid alcoholic drinks be careful when driving a motor vehicle or operating machinery alcohol, sedatives, & tranquilizers may increase drowsiness

Stop use and ask a doctor if pain, or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-422-1222) right away. Usual medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms

Directions take only as directed do not exceed 4 doses per 24 hrs adults & children 12 yrs & over 2 softgels with water every 6 hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use

Other information store at room temperature

Inactive ingredients D&C yellow #10, FD&C blue #1, gelatin, glycerin, polyethylene glycol, polyethylene glycol, purified water, stearic acid, sorbitol sorbitan solidifier, titanium dioxide

Questions? Call 1-877-299-4888

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

SAFETY SEALED. DO NOT USE IF THE INDIVIDUAL BUSTER UNIT IS OPEN OR TORN

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not a generic. It is not distributed by Pfizer. A generic product may contain the same active ingredients as Vicks® DayQuil®, NyQuil®, and LiquidCaps™.



Print/Varnish Omit Area

Lot #: Exp Date:

NDC 093-073-34

PARENTS: Learn about how medicine works www.TopMedicineForKids.org

DISTRIBUTED BY: WALGREEN CO. DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
Refunds limited to unopened and unused products
walgreens.com
©2024 Walgreen Co. MADE IN INDIA



W3090019-6-F
HEM-0423

DAYTIME NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hbr, phenylephrine hcl and acetaminophen, dextromethorphan hbr, doxylamine succinate kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0729
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0729-24	2 in 1 CARTON	10/15/2021	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1		8
Part 2		4

Part 1 of 2

DAYTIME COLD AND FLU

acetaminophine dextromethorphan hbr phenylphrine hcl capsule, liquid filled

Product Information

Item Code (Source)	NDC:0363-0853
---------------------------	---------------

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

POVIDONE (UNII: FZ989GH94E)	
SORBITAN (UNII: 6O92ICV9RU)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
SHELLAC (UNII: 46N107B71O)	
WATER (UNII: 059QF0KO0R)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL (Oblong shaped)	Size	21mm
Flavor		Imprint Code	70
Contains			

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/15/2021	

Part 2 of 2

NIGHTTIME COLD AND FLU

acetaminophen dextromethorphan hbr doxylamine succinate capsule, liquid filled

Product Information

Item Code (Source)	NDC:0363-0850
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	

POVIDONE (UNII: FZ989GH94E)	
WATER (UNII: 059QF0KO0R)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SHELLAC (UNII: 46N107B71O)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SORBITAN (UNII: 6O92ICV9RU)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
GLYCERIN (UNII: PDC6A3C0OX)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL (Oblong shaped)	Size	21mm
Flavor		Imprint Code	71
Contains			

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/15/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/15/2021	

Labeler - WALGREENS (008965063)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(0363-0729)