

**THOMPSON CHEWABLE BISMUTH TABS- bismuth subsalicylate tablet, chewable**

**Thompson Safety LLC**

-----  
**Thompson Chewable Bismuth Tabs**

***Drug Facts***

***Active ingredient***

Bismuth Subsalicylate 262 mg  
(each tablet contains 102 mg salicylate)

***Purpose***

Upset stomach reliever/antidiarrheal

***Uses***

relieves

- traveler's diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including: • heartburn, • indigestion, • nausea, • gas, • belching, • fullness

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are:**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if you have**

- bloody or black stool
- an ulcer
- a bleeding problem

**Ask a doctor before use if you have**

- fever
- mucus in the stool

**Ask a doctor or pharmacist if you are taking any drug for**

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

**When using this product** a temporary and harmless darkening of the tongue and/or stool may occur.

**Stop use and ask a doctor if**

- symptoms get worse or lasts more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

In case of accidental overdose, contact a physician or poison control center right away (1-800-222-1222)

**Directions**

- chew or dissolve in mouth
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Adults and children: (12 years and over)**

- chew 2 tablets every 1/2 to 1 hour as needed
- do not exceed 16 tablets in 24 hours
- use until diarrhea stops but not more than 2 days

**Children under 12 years:** ask a doctor

**Other information**

- **phenylketonurics:** contains phenylalanine 1.1mg per tablet
- **calcium** content per tablet: 73mg
- store at room temperature 59°-86°F (15°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

**Inactive ingredients**

acacia gum, aspartame, calcium carbonate, D&C red #27 aluminum lake, dextrans, magnesium stearate, maltodextrin, microcrystalline cellulose, peppermint flavor, silicon dioxide

**Questions or comments?** 877.506.4291

Chewable Bismuth Tabs

Upset Stomach / Diarrhea Relief

Thompson

Safety Works | Business Thrives™

Bismuth Subsalicylate 262 MG

36 Tablets

Tamper Evident

### Drug Facts

**Active ingredient (in each tablet)** **Purpose**  
Bismuth Subsalicylate 262 mg .....Upset stomach reliever/antidiarrheal (each tablet contains 102 mg salicylate)

**Uses**  
relieves ■ travelers' diarrhea ■ diarrhea  
■ upset stomach due to overindulgence in food and drink, including:  
■ heartburn ■ indigestion ■ nausea  
■ gas ■ belching ■ fullness

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome a rare but serious illness.

**Allergy alert:** Contains salicylate. Do not take if you are: ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

**Do not use if you have**  
■ bloody or black stool ■ an ulcer ■ a bleeding problem

**Ask a doctor before use if you have** ■ fever ■ mucus in the stool

**Ask a doctor or pharmacist if you are taking any drug for**  
■ anticoagulation (thinning of the blood) ■ diabetes ■ gout ■ arthritis

**When using this product** a temporary and harmless darkening of the tongue and/or stool may occur.

**Stop use and ask a doctor if**  
■ symptoms get worse or lasts for more than 2 days  
■ ringing in the ears or loss of hearing occurs  
■ diarrhea lasts more than 2 days

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.** In case of accidental overdose, contact a physician or poison control center right away (1-800-222-1222).

**Directions**  
■ chew or dissolve in mouth  
■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Adults and children:** (12 years and older) ■ chew 2 tablets every 1/2 to 1 hour as needed  
■ do not exceed 16 tablets in 24 hours  
■ use until diarrhea stops, but not more than 2 days

**Children under 12 years:** Ask a doctor

**Other information**  
■ phenylketonurics: contains phenylalanine 1.1 mg per tablet  
■ calcium content per tablet: 73 mg  
■ store at room temperature 59°-86°F (15°-30°C)  
■ tamper-evident sealed packets  
■ do not use any opened or torn packets  
*(continued on opposite panel)*

**Drug Facts (continued)**

**Inactive ingredients**  
acacia gum, aspartame, calcium carbonate, D&C red #27 aluminum lake, dextrates, magnesium stearate, maltodextrin, microcrystalline cellulose, peppermint flavoring, silicon dioxide

**Questions or comments?**  
877.506.4291

Reorder CS41220

Distributed by  
**Thompson Safety**  
Houston, TX 77073  
877.506.4291  
www.thompson-safety.com



0 73408 4 1220 1

# CHEWABLE BISMUTH TABS

UPSET STOMACH / DIARRHEA RELIEF



**THOMPSON**  
SAFETY WORKS | BUSINESS THRIVES

BISMUTH SUBSALICYLATE 262 MG

36 Tablets
 TAMPER EVIDENT

THOMPSON CHEWABLE BISMUTH TABS			
bismuth subsalicylate tablet, chewable			
Product Information			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:73408-407
<b>Route of Administration</b>	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ, BISMUTH CATION - UNII:ZS9CD118YE)	BISMUTH SUBSALICYLATE	262 mg	
Inactive Ingredients			
Ingredient Name	Strength		
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)			
<b>DEXTRATES</b> (UNII: G263MI44RU)			
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)			
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK)			
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)			
<b>D&amp;C RED NO. 27</b> (UNII: 2LRS185U6K)			
<b>ACACIA</b> (UNII: 5C5403N26O)			
<b>ASPARTAME</b> (UNII: Z0H242BBR1)			
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)			
Product Characteristics			
<b>Color</b>	pink	<b>Score</b>	no score

<b>Shape</b>	ROUND	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	RH;046
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73408-407-20	6 in 1 BOX	08/11/2021	
1		6 in 1 CELLO PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	08/11/2021	

**Labeler** - Thompson Safety LLC (080998015)

**Registrant** - Unifirst First Aid Corporation (832947092)

Revised: 9/2025

Thompson Safety LLC