

**CHEWABLE ASPIRIN- aspirin tablet, chewable  
RETAIL BUSINESS SERVICES , LLC.**

-----  
**1105B-CAR-2021-0906**

***Drug Facts***

**Active ingredient (in each tablet)**

Aspirin 81 mg (NSAID)\*

\* nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

**Uses**

- for the temporary relief of minor aches and pains or as recommended by your doctor
- ask your doctor about other uses for this product

**Warnings**

**Reye's syndrome**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

### **Ask a doctor or pharmacist before use if you are** taking a prescription drug for:

- gout
- diabetes
- arthritis

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information and warnings

**Inactive ingredients**

corn starch, FD&C yellow #6 aluminum lake, flavor, magnesium stearate, silicon dioxide, sucrose

**PRINCIPAL DISPLAY PANEL**

CAREONE™

NDC 72476-205-01

Compare to the active ingredient in Bayer® Chewable Aspirin†

Low Dose

CHEWABLE ASPIRIN

Pain Reliever (NSAID)

Orange Flavored

Aspirin Regimen\*\*

Actual Size

36 TABLETS - 81mg EACH

INK AND COATING FREE  
FOR LOT AND  
EXPIRATION STAMPING



HealthCare Distribution of Bayer® Chewable Aspirin Tablets  
S-1237 0721 FT105801CAR\_10  
NDC 72476-205-01  
DO NOT USE IF IMPRINTED SEAL  
UNDER CAP IS BROKEN OR MISSING

**Drug Facts**

**Active ingredient (in each tablet)** **Purpose**  
Aspirin 81 mg (NSAID) .....Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Uses**

- for the temporary relief of minor aches and pains or as recommended by your doctor
- ask your doctor about other uses for this product

**Warnings**

- Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
- Allergy alert:** Aspirin may cause a severe allergic reaction which may include:
- hives
  - facial swelling
  - asthma (wheezing)
  - shock
- Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
- are age 60 or older
  - have had stomach ulcers or bleeding problems
  - take a blood thinning (anticoagulant) or steroid drug

**Drug Facts (continued)**

- Take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you are taking a prescription drug for:
  - gout
  - diabetes
  - arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days

**Drug Facts (continued)**

- fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - ringing in the ears or a loss of hearing occurs
- These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information and warnings

**Inactive ingredients**

- corn starch,
- FD&C yellow #6 aluminum lake, flavor,
- magnesium stearate, silicon dioxide, sucrose

CAREone™

NDC 72476-205-01

Compare to the active ingredient  
in Bayer® Chewable Aspirin®

Low Dose

**CHEWABLE ASPIRIN**

Pain Reliever (NSAID)  
Orange Flavored   
Aspirin Regimen\*\*



Actual Size 

**36 TABLETS - 81 mg EACH**

**CHEWABLE ASPIRIN**

aspirin tablet, chewable

**Product Information**

**Product Type**

HUMAN OTC DRUG

**Item Code (Source)**

NDC:72476-205

**Route of Administration**

ORAL

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

**Inactive Ingredients**

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SUCROSE (UNII: C151H8M554)	

**Product Characteristics**

Color	orange	Score	no score
Shape	ROUND	Size	8mm
Flavor	ORANGE	Imprint Code	AP;009
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72476-205-01	1 in 1 CARTON	09/06/2021	
1		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:72476-205-03	3 in 1 CARTON	09/06/2021	
2		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/06/2021	

**Labeler** - RETAIL BUSINESS SERVICES , LLC. (967989935)

Revised: 10/2024

RETAIL BUSINESS SERVICES , LLC.