

**LOW DOSE ASPIRIN- aspirin tablet, chewable**  
**RETAIL BUSINESS SERVICES , LLC.**

-----  
**1152B-CAR-2021-0906**

***Drug Facts***

**Active ingredient (in each tablet)**

Aspirin 81 mg (NSAID)\*

\* nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

**Uses**

- temporarily relieves minor aches and pains
- ask your doctor about other uses for this product

**Warnings**

**Reye's syndrome**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

- take more or for a longer time than directed

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

### **Ask a doctor or pharmacist before use if you are** taking a prescription drug for:

- gout
- diabetes
- arthritis

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

### **Other information**

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information and warnings

### **Inactive ingredients**

corn starch, FD&C yellow #6 aluminum lake, flavor, magnesium stearate, silicon dioxide, sucrose

### **PRINCIPAL DISPLAY PANEL**

CAREONE™

NDC 72476-252-01

Compare to the active ingredient in St. Joseph® Low Dose Chewable Aspirin†

LOW DOSE ASPIRIN 81 mg

Pain Reliever (NSAID)

Orange Flavored

Adult Aspirin Regimen\*\*

Actual Size

36 CHEWABLE TABLETS

INK AND COATING FREE  
FOR LOT AND  
EXPIRATION STAMPING



DO NOT USE IF IMPRINTED SEAL  
UNDER CAP IS BROKEN OR MISSING

This product is not manufactured or distributed by  
Foundation Consumer Healthcare, LLC, distributor of  
St. Joseph® Low Dose Chewable Aspirin.

**Drug Facts**

**Active Ingredient  
(in each tablet)**

**Purpose**

Aspirin 81 mg (NSAID) .....Pain reliever  
\* nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains
- ask your doctor about other uses for this product

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergic alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

**Drug Facts (continued)**

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for:**

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - stomach bleedings: pain gets worse or lasts more than 10 days

**Drug Facts (continued)**

- fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - ringing in the ears or a loss of hearing occurs
- These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

**Other information**

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information and warnings

**Inactive ingredients**

corn starch, FD&C yellow #6 aluminum lake, flavor, magnesium stearate, silicon dioxide, sucrose

NC

CAREone™

Compare to the active ingredient in St. Joseph® Low Dose Chewable Aspirin®

**LOW DOSE ASPIRIN 81 mg**

Pain Reliever (NSAID)  
Orange Flavored  
Adult Aspirin Regimen\*\*



Actual Size



**36 CHEWABLE TABLETS**

NC

**LOW DOSE ASPIRIN**

aspirin tablet, chewable

**Product Information**

**Product Type**

HUMAN OTC DRUG

**Item Code (Source)**

NDC:72476-252

**Route of Administration**

ORAL

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

**Inactive Ingredients**

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SUCROSE (UNII: C151H8M554)	

**Product Characteristics**

Color	orange	Score	no score
Shape	ROUND	Size	8mm
Flavor	ORANGE	Imprint Code	AP;009
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72476-252-01	1 in 1 CARTON	09/06/2021	
1		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/06/2021	

**Labeler** - RETAIL BUSINESS SERVICES , LLC. (967989935)

Revised: 10/2024

RETAIL BUSINESS SERVICES , LLC.