

STERITALC 4G- talc powder
STERITALC 2G- talc powder
STERITALC 3G- talc powder
Novatech SA

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use STERITALC® safely and effectively. See full prescribing information for STERITALC®.

STERITALC® (talc), powder, for intrapleural use
Initial U.S. Approval: 2003

RECENT MAJOR CHANGES -----

Warnings and Precautions (5.4) 06/2025

INDICATIONS AND USAGE -----

INDICATIONS AND USAGE _____

STERITALC® is a sclerosing agent indicated: (1)

- To decrease the recurrence of malignant pleural effusions in symptomatic patients following maximal drainage of the pleural effusion. (1.1)
- In adults to decrease the recurrence of pneumothorax. (1.2)

DOSAGE AND ADMINISTRATION -----

DOSAGE AND ADMINISTRATION _____ (2)

- Malignant Pleural Effusion: The recommended dose is 2 to 5 grams administered intrapleurally. (2.2)
- Pneumothorax: The recommended dose is 2 grams administered intrapleurally.
- Do not exceed a total cumulative dosage of 10 grams per procedure. (2.3)
- Prepare and administer as recommended. (2.5)

DOSAGE FORMS AND STRENGTHS -----

DOSAGE FORMS AND STRENGTHS _____ (3)

- Powder: 2 grams, in a 50 mL single-dose vial (3)
- Powder: 4 grams, in a 50 mL single-dose vial (3)
- Powder: 3 grams, in a 10 mL single-dose vial (3)

CONTRAINDICATIONS -----

CONTRAINDICATIONS _____ (4)

Pregnancy. (4, 5.3, 8.1) (4)

WARNINGS AND PRECAUTIONS -----

WARNINGS AND PRECAUTIONS _____ (5)

- Pneumonitis and Acute Respiratory Distress Syndrome (ARDS): Acute Pneumonitis and ARDS, have been reported with intrapleural use of various talc products. (5.1)
- Interference with Future Procedures: Sclerosis of the pleural space may preclude or complicate subsequent ipsilateral surgery and diagnostic procedures. (5.2)
- Lead Content: STERITALC® contains lead as an impurity. May cause lead toxicity, especially in children. (5.3)
- Asbestos Content: Talc products may contain trace amounts of asbestos. Exposure to asbestos may increase the risk of mesotheliomas and other cancers. (5.4, 11)

(5)

ADVERSE REACTIONS -----

ADVERSE REACTIONS _____ (6)

Common adverse reactions are fever and pain. (6) (6)

(6)

(6)

(6)

To report suspected adverse reactions contact Boston Medical Product Inc., at 508-898-9300 and FDA at 1-800-332-1088 or www.fda.gov/medwatch. (6)

USE IN SPECIFIC POPULATIONS -----

-----**USE IN SPECIFIC POPULATIONS**----- (7)

- Females and Males of Reproductive Potential: Advise females of reproductive potential to use effective contraception. (8.3)
- Lactation: Advise women not to breastfeed. (8.2)

Revised: 12/2024

FULL PRESCRIBING INFORMATION: CONTENTS*

- 1. INDICATIONS AND USAGE**
- 2. DOSAGE AND ADMINISTRATION**
- 3. DOSAGE FORMS AND STRENGTHS**
- 4. CONTRAINDICATIONS**
- 5. WARNINGS AND PRECAUTIONS**
- 6. ADVERSE REACTIONS**
- 8. USE IN SPECIFIC POPULATIONS**
- 11. DESCRIPTION**
- 12. CLINICAL PHARMACOLOGY**
- 13. NONCLINICAL TOXICOLOGY**
- 14. CLINICAL STUDIES**
- 16. HOW SUPPLIED/STORAGE AND HANDLING**

Principal Display Label, Steritalc 3g vial label

Principal Display Label, Steritalc 4g vial label

Principal Display Label, Steritalc 2g vial label

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1. INDICATIONS AND USAGE

1. INDICATIONS AND USAGE

1.1 Malignant Pleural Effusion

STERITALC is indicated to decrease the recurrence of malignant pleural effusions in symptomatic patients following maximal drainage of the pleural effusion.

1.2 Pneumothorax

STERITALC is indicated in adults to decrease the recurrence of pneumothorax.

2. DOSAGE AND ADMINISTRATION

2.1 Important Use Information

STERITALC is for pleurodesis only - Do NOT administer STERITALC intravenously.

Administer STERITALC after adequate drainage of the pleural effusion or air.

2.2 Recommended Dose for Malignant Pleural Effusion

The recommended dose for malignant pleural effusion is 2 to 5 grams administered intrapleurally. According to the physician's discretion, and in consideration of diagnosis and patient's condition, different dosages may be applied, but a total dosage of 10 grams should not be exceeded.

2.3 Recommended Dose for Pneumothorax

The recommended dose for pneumothorax is 2 grams administered intrapleurally. According to physician's discretion and in consideration of diagnosis and patient's condition, different dosages may be applied, but a cumulative dosage of 10 grams should not be exceeded.

2.4 Preparation

Slurry for Tube Thoracostomy

STERITALC 2 grams and 4 grams dosage forms

Do not prepare the slurry in advance. Use the slurry immediately after preparation.

Prepare the talc suspension using aseptic technique in an appropriate laminar flow hood as follows:

Step 1: Fully bend or remove the flap into the direction of the arrow. The top can now be punctured to mix the slurry.

Step 2: Using a 16 gauge needle attached to a 60-mL Luer Lok syringe, draw up 50 mL of 0.9 % Sodium Chloride injection, USP. Vent the talc bottle using a needle. Slowly inject the 50 mL of 0.9% Sodium Chloride Injection, USP into the glass vial.

Step 3: Swirl the glass vial to disperse the talc powder.

Step 4: Divide the contents of the glass vial equally into two 60-mL Luer Lok syringes, each attached with a 16 gauge needle, by withdrawing 25 mL of the suspension into each syringe with continuous swirling. Add 0.9% Sodium Chloride Injection, USP to a total volume of 50 mL in each syringe. Draw 10 mL of air into each syringe to the 60 mL mark to serve as a headspace for mixing prior to administration.

For STERITALC 2 grams, each syringe should contain 1 gram of Sterile Talc Powder in 50 mL of 0.9% Sodium Chloride Injection, USP with an air headspace of 10 mL.

For STERITALC 4 grams, each syringe should contain 2 grams of Sterile Talc Powder in 50 mL of 0.9% Sodium Chloride Injection, USP with an air headspace of 10 mL.

Step 5: Label the syringes with the talc concentration, the expiration date and time, the identity of the patient intended to receive the material, and the following statements:

“SHAKE SYRINGE WELL to resuspend before administration”

“FOR PLEURODESIS ONLY - not for intravenous administration”

Step 6: If not used immediately, store prepared suspension in refrigerator. Discard the prepared suspension if not used within 12 hours.

Insufflation/Poudrage

STERITALC 2 grams and 4 grams dosage forms

Use an FDA-approved or cleared device for STERITALC insufflation/poudrage.

Step 1: Fully remove the top from the vial.

Step 2: Fill the content into the applicator for insufflation/poudrage.

Step 3: Follow the selected device manufacturer's instructions for insufflation/poudrage.

STERITALC 3 grams dosage form

STERITALC 3 grams is designed to be used for the administration of talc in combination with a compatible FDA approved or FDA cleared device intended for manual insufflation of medical grade talc into the pleural cavity during pleurodesis such as Novatech SA's NOVATECH® TALCAIR™.

Step 1: Remove the aluminum tear-off cap from the vial.

Step 2: Remove the stopper from the vial.

Step 3: Close the vial with the vial coupling.

Step 4: Firmly press the vial coupling onto the vial until you feel and hear the lid "click" onto the vial top. To do this, place the vial on a firm and level base.

Step 5: Connect the insufflation bulb to the vial coupling using the Luer lock connector.

Step 6: Ensure that all components are firmly connected with each other.

2.5 Administration

Slurry for Tube Thoracostomy

STERITALC 2 grams and 4 grams dosage forms

Prior to administration, continuously agitate the syringes to evenly redisperse the talc and avoid settlement. Immediately prior to administration, vent the 10 mL air headspace from each syringe. Administer the talc suspension through the chest tube according to standard procedures.

Step 1: Inject the slurry through the pleural drainage into the pleural cavity.

Step 2: Clamp the pleural drainage. Keep the negative pressure in the pleural cavity.

Step 3: While the slurry remains in the pleural cavity, reposition the patient regularly to achieve even distribution of the slurry.

Step 4: Aspirate the slurry through the pleural drainage.

Insufflation/Poudrage

Perform intervention by means of a tube thoracoscopy. Follow the selected device manufacturer's instructions for insufflation/poudrage.

STERITALC 2 grams and 4 grams dosage forms

Step 1: Introduce the cannula into the trocar.

Step 2: Distribute the talc evenly in the pleural cavity. To do this, carefully spray several times. After spraying a few times, change the cannula direction.

After use, some talc may remain in the vial. Discard unused portion.

STERITALC 3 grams dosage form

Keep the product upright during use. Avoid contact between the cannula tip and tissue/body fluids. Otherwise the cannula may be blocked. In case of blockage: Shorten the cannula with a scalpel to be able to continue the intervention. Cut perpendicularly to the cannula and ensure that the cutting edge is straight and free of burrs.

Step 1: Introduce the cannula into the trocar.

Step 2: Distribute the talc evenly in the pleural cavity. To do this, press the insufflation bulb carefully and at regular intervals. After spraying a few times, change the cannula direction.

After use, some talc may remain in the vial. Discard unused portion

3. DOSAGE FORMS AND STRENGTHS

White or off-white to light gray sterile powder provided in the following strengths:

- Powder, for intrapleural use: STERITALC, 2 grams in a 50 mL single-dose vial
- Powder, for intrapleural use: STERITALC, 4 grams in a 50 mL single-dose vial
- Powder, for intrapleural use: STERITALC, 3 grams in a 10 mL single-dose vial, for use with Novatech SA's NOVATECH® TALCAIR™

4. CONTRAINDICATIONS

Due to lead content, STERITALC® is contraindicated in pregnant women and can cause fetal harm and potential loss of pregnancy [*see Warnings and Precautions (5.3) and Use in Specific Populations (8.1)*].

5. WARNINGS AND PRECAUTIONS

5.1 Pneumonitis and Acute Respiratory Distress Syndrome (ARDS)

Acute Pneumonitis and ARDS, including fatal cases, have been reported with intrapleural use of various talc products. Systemic exposure of talc could be affected by the integrity of the visceral pleura and could be increased if talc is administered immediately after mechanical abrasion or biopsy of the pleura. The literature also suggests a possible correlation between talc particle size distribution and risk of toxicity.

There are published reports of two large, prospective trials conducted to evaluate the safety of STERITALC administered intrapleurally. One trial evaluated 558 patients treated with STERITALC 4g by poudrage for MPE. The second trial evaluated 418 patients with recurrent primary spontaneous pneumothorax treated with STERITALC 2 g by poudrage. No cases of ARDS or talc-related lung injury were reported.

5.2 Interference with Future Procedures

Sclerosis of the pleural space may preclude or complicate subsequent ipsilateral surgery and diagnostic procedures. Consider the possible effects of the use of STERITALC on future diagnostic and therapeutic procedures prior to administration.

5.3 Lead Content

Lead is present in STERITALC as an impurity. The main target organ for lead toxicity is the nervous system, but effects of lead exposure also include increased blood pressure, anemia, decreased sperm production, and damage to the kidneys in children and adults. Minimal risk levels of lead have not been derived for humans because clear thresholds for effects have not been identified.

Children are more sensitive to lead toxicity than adults, and no safe blood level has been determined in children. Cognitive and neurobehavioral deficits are observed in children exposed to lead. Exposure of a pregnant woman to lead may cause miscarriage, premature birth, lower birth weights and slow or impaired mental development in the child.

Administration of STERITALC at the highest recommended dose of 10 grams may deliver up to 40 mcg of lead.

5.4 Asbestos Content

Talc products may contain trace amounts of asbestos. Asbestos is a known human carcinogen. Exposure to asbestos may increase the risk of mesotheliomas and other cancers including lung, larynx, and ovarian cancer.

STERITALC is tested for asbestos and asbestiform fibers [see DESCRIPTION (11)].

6. ADVERSE REACTIONS

The following clinically significant adverse reactions are described elsewhere in the labeling.

- Pneumonitis and Acute Respiratory Distress Syndrome (ARDS) [see Warnings and Precautions (5.1)]
- Lead Content [see Warnings and Precautions (5.3)]
- Asbestos Content [see Warnings and Precautions (5.4)]

Common adverse reactions observed with intrapleural use of various talc products are fever and pain.

Other adverse reactions include dyspnea, arrhythmia, empyema, pneumonitis and acute respiratory distress syndrome (ARDS).

Procedure related adverse reactions such as bleeding, hemothorax, wound infections, atelectasis and pneumonia may occur.

8. USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

STERITALC is contraindicated for use in pregnant women because it contains lead, which can cause fetal harm and potential loss of pregnancy [see *Warnings and Precautions (5.3)*]. There are no human data on the use of STERITALC in pregnant women. In an animal reproduction study, oral administration of talc in pregnant rabbits during organogenesis revealed no evidence of teratogenicity at dose up to approximately 5

times the human dose [see Data].

Data

Animal Data

In an embryo-fetal developmental toxicity study in rabbits, talc was administered to rabbits by oral gavage daily during the period of organogenesis at doses up to 900 mg/kg (approximately 5 times the human dose on a mg/m² basis). No significant dose-related toxicity was reported except at maternally toxic doses. In multiple animal studies, intrapleurally administered talc was not absorbed systemically.

8.2 Lactation

Risk Summary

There is no information regarding the presence of talc in human milk, the effects on the breastfed infants, or the effects on milk production. STERITALC contains lead, which is known to have adverse effects on health and development in neonates, infants and children [see Warnings and Precautions (5.3)]. Because of the potential for serious adverse reactions in breast-fed infants from lead content in STERITALC, advise lactating women not to breastfeed during treatment with STERITALC and for 5 months after the final dose.

8.3 Females and Males of Reproductive Potential

Contraception

STERITALC contains lead, which can cause fetal harm when administered to a pregnant woman [see Warnings and Precautions (5.3) and Use in Specific Populations (8.1)]. Advise females of reproductive potential to use effective contraception during treatment and for 5 months following the final dose of STERITALC.

Infertility

STERITALC contains lead, which may impair fertility in males of reproductive potential [see Warnings and Precautions (5.3)].

8.4 Pediatric Use

Safety and effectiveness have not been established in pediatric patients. Lead is present in STERITALC as an impurity [see Warnings and Precautions (5.3)].

11. DESCRIPTION

STERITALC (talc) is a sclerosing agent for intrapleural administration. The molecular formula for talc is Mg₃Si₄O₁₀(OH)₂. The molecular weight is 379.3 g/mole. The chemical name of talc is hydrated magnesium silicate.

The talc powder contains ≥ 95% of hydrated magnesium silicate; associated minerals include chlorites (hydrated aluminum and magnesium silicates), magnesite (magnesium carbonate), calcite (calcium carbonate), and dolomite (calcium and magnesium carbonate). The talc elementary sheet is composed of a layer of magnesium-oxygen/hydroxyl octahedra, sandwiched between two layers of silicon-oxygen tetrahedra. Talc is insoluble in water.

STERITALC is tested for asbestos and asbestiform fibers consistent with USP standards for talc-containing products.

STERITALC is supplied as 2 grams, 3 grams, and 4 grams white or off-white to light gray, sterile powder. STERITALC contains particle size-controlled talc, graded to decrease the proportion of smaller particles.

12. CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Talc instilled into the pleural cavity is thought to result in an inflammatory reaction. This reaction promotes adherence of the visceral and parietal pleura to prevent reaccumulation of pleural air or fluid.

13. NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Studies on the carcinogenicity of talc have been performed using non-standard designs which prevent firm conclusions on its carcinogenicity. With single intraperitoneal administration to mice at 20 mg and observation for at least 6 months, or 4 weekly doses administered intraperitoneally at 25 mg/dose to rats with observation for at least 84 weeks, tumor incidence was not increased. In these studies the talc and its asbestos content were not characterized.

Genotoxicity was tested in cultures of rat pleural mesothelial cells (RPMC) for unscheduled DNA synthesis (UDS) and sister chromatid exchanges (SCEs). None of the talc samples induced enhancement of UDS or SCEs in treated cultures.

No information is available on impairment of fertility in animals by talc.

14. CLINICAL STUDIES

14.1 Malignant Pleural Effusion

The data demonstrating safety and efficacy of talc in the treatment of malignant pleural effusions are derived from the published medical literature. In these studies, greater than 1000 patients with malignant pleural effusions have been reported (with varying degrees of detail and durations of response) to have had successful pleurodesis with talc, administered either by poudrage or as a slurry via chest tube. There are published efficacy data for more than 200 patients treated with STERITALC for malignant pleural effusion, with a success rate of approximately 89% (range 73-91%).

14.2 Pneumothorax

The data demonstrating safety and efficacy of talc in the treatment of pneumothorax are derived from the published medical literature. There are published efficacy data for more than 500 patients treated with STERITALC for pneumothorax, some in conjunction with surgical procedures, with successful pleurodesis rates of 97-100%.

16. HOW SUPPLIED/STORAGE AND HANDLING

STERITALC, white or off-white to light gray, sterile powder is supplied as

- two grams of talc powder in a single-dose 50 mL Type III glass vial, closed with a stopper, and crimped with an aluminum cap; with 4 vials packaged in one carton.
- NDC 62327-222-02 (vial)

NDC 62327-222-42 (carton)

four grams of talc powder in a single-dose 50 mL Type III glass vial, closed with a stopper, and crimped with an aluminum cap; with 4 vials packaged in one carton.

NDC 62327-444-04 (vial)

NDC 62327-444-44 (carton)

- three grams of talc powder in a single-dose 10 mL Type I brown glass vial, closed with a stopper, and crimped with an aluminum cap; with 4 vials packaged in one carton.

NDC 62327-333-03 (vial)

NDC 62327-333-43 (carton)

Store at 20°C - 25°C (68°F - 77°F); excursions permitted between 15°C - 30°C (59°F - 86°F) [see USP Controlled Room Temperature]. Protect against sunlight.

17. PATIENT COUNSELLING INFORMATION

- Advise patients to notify their healthcare provider if new or worsening pulmonary symptoms develop [*see Warnings and Precautions (5.1)*].
- Inform patients that the lead content in STERITALC can be harmful to a developing fetus. Advise females of reproductive potential to use effective contraception during treatment and for 5 months after the last dose of STERITALC [*see Warnings and Precautions (5.3) and Use in Specific Populations (8.3)*].

Manufactured for: Novatech SA

Z.I. Athélia III - 1058, Voie Antiope

13705 La Ciotat CEDEX

France

By: bess pro gmbh

Gustav-Krone-Strasse 7

14167 Berlin

Germany

NO128-03

Principal Display Label, Steritalc 3g package label

NDC: 62327-333-03

STERITALC® **3g/vial**

(Talc) Powder
for intrapleural administration

Manufactured for:
Novatech SA
Z.I. Athelia III - 1058, Voie Antiope
13705 La Ciotat CEDEX
France

By:
bess pro gmbh
Gustav-Krone-Str. 7
14167 Berlin
Germany

3 grams of sterile talc per vial
Refer to the Prescribing Information for dosage and preparation. Single-dose vial. Discard unused portion.
Store at 20-25°C (68-77°F);
excursions permitted between 15°C to 30°C (59°-86°F)
(See USP controlled Room temperature).
Protect against sunlight.

Rx only

LOT:

Expiry:



Principal Display Label, Steritalc 3g vial label

NDC: 62327-333-03

STERITALC® **3g/vial**

(Talc) Powder
for intrapleural administration

Manufactured for:
Novatech SA
Z.I. Athelia III - 1058, Voie Antiope
13705 La Ciotat CEDEX
France

By:
bess pro gmbh
Gustav-Krone-Str. 7
14167 Berlin
Germany

3 grams of sterile talc per vial
Refer to the Prescribing Information for dosage and preparation. Single-dose vial. Discard unused portion.
Store at 20-25°C (68-77°F);
excursions permitted between 15°C to 30°C (59°-86°F)
(See USP controlled Room temperature).
Protect against sunlight.

Rx only

LOT:

Expiry:



Principal Display Label, Steritalc 4g package label

NDC: 62327-444-04

STERITALC® **4g/vial**

(Talc) Powder
for intrapleural administration

Manufactured for:
Novatech SA
Z.I. Athelia III - 1058, Voie Antiope
13705 La Ciotat CEDEX
France

By:
bess pro gmbh
Gustav-Krone-Str. 7
14167 Berlin
Germany

4 grams of sterile talc per vial
Refer to the Prescribing Information for dosage and preparation. Single-dose vial. Discard unused portion.
Store at 20-25°C (68-77°F);
excursions permitted between 15°C to 30°C (59°-86°F)
(See USP controlled Room temperature).
Protect against sunlight.

Rx only

LOT:

Expiry:



Principal Display Label, Steritalc 4g vial label

NDC: 62327-444-04

STERITALC® **4g/vial**

(Talc) Powder
for intrapleural administration

Manufactured for:
Novatech SA
Z.I. Athelia III - 1058, Voie Antiope
13705 La Ciotat CEDEX
France

By:
bess pro gmbh
Gustav-Krone-Str. 7
14167 Berlin
Germany

4 grams of sterile talc per vial
Refer to the Prescribing Information for dosage and
preparation. Single-dose vial. Discard unused portion.
Store at 20-25°C (68-77°F);
excursions permitted between 15°C to 30°C (59°-86°F)
(see USP controlled Room temperature).
Protect against sunlight.

Rx only

LOT:

Expiry:



1.14.2.1.6 Pg. 1

Principal Display Label, Steritalc 2g package label

NDC: 62327-222-02

STERITALC® **2g/vial**

(Talc) Powder
for intrapleural administration

Manufactured for:
Novatech SA
Z.I. Athelia III - 1058, Voie Antiope
13705 La Ciotat CEDEX
France

By:
bess pro gmbh
Gustav-Krone-Str. 7
14167 Berlin
Germany

2 grams of sterile talc per vial
Refer to the Prescribing Information for dosage and
preparation. Single-dose vial. Discard unused portion.
Store at 20-25°C (68-77°F);
excursions permitted between 15°C to 30°C (59°-86°F)
(see USP controlled Room temperature).
Protect against sunlight.

Rx only

LOT:

Expiry:



1.14.2.1.4 Pg. 1

Principal Display Label, Steritalc 2g vial label

NDC: 62327-222-02

STERITALC® **2g/vial**

(Talc) Powder
for intrapleural administration

Manufactured for:
Novatech SA
Z.I. Athelia III - 1058, Voie Antiope
13705 La Ciotat CEDEX
France


By:
bess pro gmbh
Gustav-Krone-Str. 7
14167 Berlin
Germany

2 grams of sterile talc per vial
Refer to the Prescribing Information for dosage and
preparation. Single-dose vial. Discard unused portion.
Store at 20-25°C (68-77°F);
excursions permitted between 15°C to 30°C (59°-86°F)
(see USP controlled Room temperature).
Protect against sunlight.

Rx only

LOT:

Expiry:



1.14.2.1.4 Pg. 1

Steritalc final carton label 4g N444

115x49x210 mm

NDC 62327-444-44

STERITALC® 4g/vial
(Talc) Powder
for intrapleural administration

NDC 62327-444-44

STERITALC® 4g/vial
(Talc) Powder
for intrapleural administration

4 vials
4 grams of sterile talc per vial

Refer to the Prescribing Information for dosage and preparation.
Single-dose vial. Discard unused portion.
Store at 20-25°C (68-77°F); excursions permitted between 15°C to 30°C (59°-86°F)
[see USP controlled Room temperature].
Protect against sunlight.

Rx only



NDC 62327-444-44
STERITALC® 4g/Vial
(Talc) Powder
for intrapleural administration

NDC 62327-444-44

STERITALC® 4g/vial
(Talc) Powder
for intrapleural administration

4 vials
4 grams of sterile talc per vial

Refer to the Prescribing Information for dosage and preparation.
Single-dose vial. Discard unused portion.
Store at 20-25°C (68-77°F); excursions permitted between 15°C to 30°C (59°-86°F)
[see USP controlled Room temperature].
Protect against sunlight.

Rx only



3 02327 44444 4

N444-02 - 2020/09

STERITALC® 4g/vial

NDC 62327-444-44

(Talc) Powder for intrapleural administration

By: bios pro gmbh
Helmholtzstrasse 7
44183 Datteln
Germany

Manufactured for:
Novatech SA
11000, Via Joffre
13705 Le Castellet
France



Distributed in the U.S. by:
Boston Medical Products, Inc.
1000 Main Street
Shrewsbury, MA 01545
800-433-2674
508-895-5200



115x49x210 mm

NDC 62327-333-43

STERITALC® 3g/vial
(Talc) Powder
for intrapleural administration

NDC 62327-333-43

STERITALC® 3g/vial
(Talc) Powder
for intrapleural administration

4 vials
3 grams of sterile talc per vial

Refer to the Prescribing Information for dosage and preparation.
Single-dose vial. Discard unused portion.
Store at 20-25°C (68-77°F); excursions permitted between 15°C to 30°C (59°-86°F)
[see USP controlled Room temperature].
Protect against sunlight.

Rx only



NDC 62327-333-43
STERITALC® 3g/vial
(Talc) Powder
for intrapleural administration

NDC 62327-333-43

STERITALC® 3g/vial
(Talc) Powder
for intrapleural administration

4 vials
3 grams of sterile talc per vial

Refer to the Prescribing Information for dosage and preparation.
Single-dose vial. Discard unused portion.
Store at 20-25°C (68-77°F); excursions permitted between 15°C to 30°C (59°-86°F)
[see USP controlled Room temperature].
Protect against sunlight.

Rx only



3 02327 33343 4

By: bios pro gmbh
Helmholtzstrasse 7
44139 Dortmund
Germany

Manufactured for:
Novatech SA
118, Rue de la
Liberte
13785 La Cote d'Azur
France



Distributed in the U.S. by:
Boston Medical Products, Inc.
1000 Main Street
Shrewsbury, MA 01545
800-433-2674
508-895-9300




N443-03 - 2023/12

STERITALC® 3g/vial

NDC 62327-333-43

(Talc) Powder for intrapleural administration

115x49x210 mm



STERITALC 4G

talc powder

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:62327-444
Route of Administration	INTRAPLEURAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TALC (UNII: 7SEV7J4R1U) (TALC - UNII: 7SEV7J4R1U)	TALC	4 g in 50 mL

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62327-444-44	4 in 1 BOX	08/01/2017	
1	NDC:62327-444-04	50 mL in 1 VIAL, GLASS; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA205555	08/01/2017	

STERITALC 2G

talc powder

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:62327-222
Route of Administration	INTRAPLEURAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TALC (UNII: 7SEV7J4R1U) (TALC - UNII:7SEV7J4R1U)	TALC	2 g in 50 mL

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62327-222-42	4 in 1 BOX	11/01/2017	

1	NDC:62327-222-02	50 mL in 1 VIAL, GLASS; Type 0: Not a Combination Product		
----------	------------------	---	--	--

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA205555	11/01/2017	

STERITALC 3G

talc powder

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:62327-333
Route of Administration	INTRAPLEURAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TALC (UNII: 7SEV7J4R1U) (TALC - UNII:7SEV7J4R1U)	TALC	3 g in 10 mL

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62327-333-43	4 in 1 BOX	11/01/2017	
1	NDC:62327-333-03	10 mL in 1 VIAL, GLASS; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA205555	11/01/2017	

Labeler - Novatech SA (777211640)

Registrant - Novatech SA (777211640)**Establishment**

Name	Address	ID/FEI	Business Operations
Novatech SA		777211640	label(62327-444, 62327-333, 62327-222)

Revised: 6/2025

Novatech SA