

**MULTI-SYMPATOM COLD AND FLU RELIEF DAYTIME NIGHTTIME-
acetaminophen dextromethorphan hbr phenylephrine hcl and acetaminophen
dextromethorphan hbr doxylamine succinate
RITE AID**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

**695T RITE AID 11822-1988 Multi-Symptom Cold & Flu Relief Daytime
Nighttime (Acetaminophen 325mg Dextromethorphan HBr10mg
Phenylephrine HCl 5mg Acetaminophen 325mg Dextromethorphan HBr 15mg
Doxylamine Succinate 6.25mg)**

DRUG FACTS

daytime cold & flu

Active ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

nighttime cold & flu

Active ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

daytime cold & flu

Purpose

Pain reliever/Fever reducer

Cough suppressant

Nasal decongestant

nighttime cold & flu

Purpose

Pain reliever/Fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion (**Daytime only**)
- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever
- runny nose & sneezing (**Nighttime only**)

Warnings

liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional

conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma (**nighttime only**)
- heart disease (**Daytime only**)
- high blood pressure (**Daytime only**)
- thyroid disease (**Daytime only**)
- diabetes (**Daytime only**)
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema (**Daytime only**)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema (**nighttime only**)

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

- taking the blood thinning drug warfarin

When using this product, do not use more than directed.(daytime only)

- excitability may occur, especially in children (**nighttime only**)
- marked drowsiness may occur (**nighttime only**)
- avoid alcoholic drinks (**nighttime only**)
- be careful when driving a motor vehicle or operating machinery. (**nighttime only**)
- alcohol, sedatives, & tranquilizers may increase drowsiness. (**nighttime only**)

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless (**daytime only**)
- pain, nasal congestion or cough get worse or last more than 7 days (**daytime only**)
- pain or cough gets worse or lasts more than 7 days (**nighttime only**)

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be Signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults and for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 4 doses per 24 hours

adults & children 12 yrs & over: 2 softgels with water every 4 hrs (daytime only)

adults & children 12 yrs & over: 2 softgels with water every 6 hrs (nighttime only)

children 4 to under 12 yrs:ask a doctor

children under 4 yrs:do not use

OTHER INFORMATION

Other information

- store at room temperature

daytime cold & flu

Inactive ingredients FD&C red#40, FD&C yellow#6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide

nighttime cold & flu

Inactive ingredients FD&C blue#1, D&C yellow#10, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide

Questions?call 1-877-290-4008

Print/Varnish Omit Area


Compare to the active ingredients in Vicks® DayQuil® and NyQuil® Cold & Flu LiquiCaps®*

NDC 11822-1988-0

MULTI-SYMPATOM COLD & FLU RELIEF

Varnish Omit Area

Varnish Omit Area

DAYTIME	NIGHTTIME
ACETAMINOPHEN 325 mg DEXTROMETHORPHAN HBr 10 mg PHENYLEPHRINE HCl 5 mg PAIN RELIEVER / FEVER REDUCER COUGH SUPPRESSANT NASAL DECONGESTANT NON-DROWSY Temporarily relieves aches, fever, sore throat, cough, and nasal congestion	ACETAMINOPHEN 325 mg DEXTROMETHORPHAN HBr 15 mg DOXYLAMINE SUCCINATE 6.25 mg PAIN RELIEVER / FEVER REDUCER COUGH SUPPRESSANT ANTIHISTAMINE Temporarily helps relieve aches, fever, sore throat, cough, runny nose, and sneezing
ACTUAL SIZE	ACTUAL SIZE
	
When using Daytime and Nighttime products, carefully read the labeling to ensure correct dosing.	
16 DAYTIME SOFTGELS	8 NIGHTTIME SOFTGELS

Drug Facts	nighttime cold & flu	Drug Facts	daytime cold & flu
Active ingredients (in each softgel)	Purpose	Active ingredients (in each softgel)	Purpose
Acetaminophen 325 mg	Pain reliever/Fever reducer	Acetaminophen 325 mg	Pain reliever/Fever reducer
Dextromethorphan HBr 15 mg	Cough suppressant	Dextromethorphan HBr 10 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine	Phenylephrine HCl 5 mg	Nasal decongestant
Uses temporarily relieves common cold/flu symptoms: • cough due to minor throat & bronchial irritation • sore throat • headache • minor aches & pains • fever		Uses temporarily relieves common cold/flu symptoms: • nasal congestion • cough due to minor throat & bronchial irritation • sore throat • headache • minor aches	

SAFETY SEALED: DO NOT USE IF THE INDIVIDUAL BLISTER UNIT IS OPEN OR TORN

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

SATISFACTION GUARANTEE
If you're not satisfied, we'll happily refund your money.

6957 0621
*This product is not manufactured or distributed by The Proctor & Gamble Company. Vicks®, DayQuil®, NyQuil®, and LiquiCaps® are registered trademarks of The Proctor and Gamble Company.

Drug Facts (continued)

• runny nose & sneezing

Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 4 doses in 24 hrs, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • liver disease • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema • trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers • taking the blood thinning drug warfarin

When using this product • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, & tranquilizers may increase drowsiness

Stop use and ask a doctor if • pain or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults and for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions • take only as directed • do not exceed 4 doses per 24 hours

adults & children 12 yrs & over	2 softgels with water every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information • store at room temperature

Inactive ingredients FD&C blue#1, D&C yellow#10, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide

Questions? call 1-877-290-4008

Drug Facts (continued)

• pains • fever

Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 4 doses in 24 hrs, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if • you get nervous, dizzy or sleepless • pain, nasal congestion or cough get worse or last more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults and for children even if you do not notice any signs or symptoms.

Directions • take only as directed • do not exceed 4 doses per 24 hours

Drug Facts (continued)

adults & children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information • store at room temperature

Inactive ingredients FD&C red#40, FD&C yellow#6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide

Questions? call 1-877-290-4008


PARENTS:
Learn about how medicine works
www.StopMedicineAbuse.org

DISTRIBUTED BY: RITE AID,
30 HUNTER LANE,
CAMP HILL, PA 17011
www.riteaid.com
MADE IN INDIA

0 11822 19883 7

Lot #:
Exp. Date:

Print/Varnish Omit Area



MULTI-SYMP TOM COLD AND FLU RELIEF DAYTIME NIGHTTIME

acetaminophen dextromethorphan hbr phenylephrine hcl and acetaminophen dextromethorphan hbr doxylamine succinate kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-1988
---------------------	----------------	---------------------------	----------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-1988-0	2 in 1 CARTON	09/15/2021	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:11822-1988-9	4 in 1 CARTON	09/15/2021	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1		16
Part 2		8

Part 1 of 2

MULTI-SYMPATOM COLD AND FLU RELIEF DAYTIME

acetaminophen 325 mg dextromethorphan hbr 10 mg phenylephrine hci 5 mg capsule, liquid filled

Product Information

Item Code (Source)	NDC:11822-1989
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
SORBITAN (UNII: 6O92ICV9RU)	
SHELLAC (UNII: 46N107B710)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
POVIDONE (UNII: FZ989GH94E)	
WATER (UNII: 059QF0KO0R)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL (OBLONG SHAPED)	Size	21mm
Flavor		Imprint Code	70
Contains			

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/15/2021	

Part 2 of 2

MULTI-SYMPATOM COLD AND FLU RELIEF NIGHTTIME

acetaminophen 325 mg dextromethorphan hbr 15 mg doxylamine succinate 6.25 mg capsule, liquid filled

Product Information

Item Code (Source)	NDC:11822-1987
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg

Inactive Ingredients

Ingredient Name	Strength
SORBITAN (UNII: 6O921CV9RU)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	
SHELLAC (UNII: 46N107B71O)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
GELATIN (UNII: 2G86QN327L)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	71

Contains**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/15/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/15/2021	

Labeler - RITE AID (014578892)**Registrant** - TIME CAP LABORATORIES, INC. (037052099)**Establishment**

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(11822-1988)

Revised: 12/2022

RITE AID