

CHLORZOXAZONE- chlorzoxazone tablet
Cintex Services, LLC

Chlorzoxazone Tablets, USP 250 mg

For Painful Musculoskeletal Conditions


DESCRIPTION

Each tablet contains:

Chlorzoxazone* 250 mg

* 5-Chloro-2-benzoxazolinone

Structural Formula:



Molecular Formula: C₇H₄ClNO₂

Molecular Weight: 169.57

Chlorzoxazone, USP is a white or practically white, practically odorless, crystalline powder. Chlorzoxazone is slightly soluble in water; sparingly soluble in alcohol, in isopropyl alcohol, and in methanol; soluble in solutions of alkali hydroxides and ammonia.

Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, docusate sodium, sodium benzoate, lactose monohydrate, magnesium stearate and microcrystalline cellulose.

Meets USP Dissolution Test 4

CLINICAL PHARMACOLOGY

Chlorzoxazone is a centrally-acting agent for painful musculoskeletal conditions. Data available from animal experiments as well as human study indicate that chlorzoxazone acts primarily at the level of the spinal cord and subcortical areas of the brain where it inhibits multisynaptic reflex arcs involved in producing and maintaining skeletal muscle spasm of varied etiology. The clinical result is a reduction of the skeletal muscle spasm with relief of pain and increased mobility of the involved muscles. Blood levels of chlorzoxazone can be detected in people during the first 30 minutes and peak levels may be reached, in the majority of the subjects, in about 1 to 2 hours after oral administration of chlorzoxazone. Chlorzoxazone is rapidly metabolized and is excreted in the urine, primarily in a conjugated form as the glucuronide. Less than one percent of a dose of chlorzoxazone is excreted unchanged in the urine in 24 hours.

INDICATIONS AND USAGE

Chlorzoxazone is indicated as an adjunct to rest, physical therapy, and other measures for the relief of discomfort associated with acute, painful musculoskeletal conditions. The mode of action of this drug has not been clearly identified, but may be related to its sedative properties. Chlorzoxazone does not directly relax tense skeletal muscles in man.

CONTRAINDICATIONS

Chlorzoxazone is contraindicated in patients with known intolerance to the drug.

WARNINGS

Serious (including fatal) hepatocellular toxicity has been reported rarely in patients receiving chlorzoxazone. The mechanism is unknown but appears to be idiosyncratic and unpredictable. Factors predisposing patients to this rare event are not known. Patients should be instructed to report early signs and/or symptoms of hepatotoxicity such as fever, rash, anorexia, nausea, vomiting, fatigue, right upper quadrant pain, dark urine, or jaundice. Chlorzoxazone should be discontinued immediately and a physician consulted if any of these signs or symptoms develop. Chlorzoxazone use should also be discontinued if a patient develops abnormal liver enzymes (e.g., AST, ALT, alkaline phosphatase and bilirubin).

The concomitant use of alcohol or other central nervous system depressants may have an additive effect.

Usage in Pregnancy: The safe use of chlorzoxazone has not been established with respect to the possible adverse effects upon fetal development. Therefore, it should be used in women of childbearing potential only when, in the judgment of the physician, the potential benefits outweigh the possible risks.

PRECAUTIONS

Chlorzoxazone should be used with caution in patients with known allergies or with a history of allergic reactions to drugs. If a sensitivity reaction occurs such as urticaria, redness, or itching of the skin, the drug should be stopped.

If any symptoms suggestive of liver dysfunction are observed, the drug should be discontinued.

ADVERSE REACTIONS

After extensive clinical use of chlorzoxazone-containing products, it is apparent that the product is well tolerated and seldom produces undesirable side effects. Occasional patients may develop gastrointestinal disturbances. It is possible in rare instances that chlorzoxazone may have been associated with gastrointestinal bleeding. Drowsiness, dizziness, lightheadedness, malaise, or overstimulation may be noted by an occasional patient. Rarely, allergic-type skin rashes, petechiae, or ecchymoses may develop during treatment. Angioneurotic edema or anaphylactic reactions are extremely rare. There is

no evidence that the drug will cause renal damage. Rarely, a patient may note discoloration of the urine resulting from a phenolic metabolite of chlorzoxazone. This finding is of no known clinical significance.

OVERDOSAGE

Symptoms: Initially, gastrointestinal disturbances such as nausea, vomiting, or diarrhea together with drowsiness, dizziness, lightheadedness or headache may occur. Early in the course there may be malaise or sluggishness followed by marked loss of muscle tone, making voluntary movement impossible. The deep tendon reflexes may be decreased or absent. The sensorium remains intact, and there is no peripheral loss of sensation. Respiratory depression may occur with rapid, irregular respiration and intercostal and substernal retraction. The blood pressure is lowered, but shock has not been observed.

Treatment: Gastric lavage or induction of emesis should be carried out, followed by administration of activated charcoal. Thereafter, treatment is entirely supportive. If respirations are depressed, oxygen and artificial respiration should be employed and a patent airway assured by use of an oropharyngeal airway or endotracheal tube. Hypotension may be counteracted by use of dextran, plasma, concentrated albumin or a vasopressor agent such as norepinephrine. Cholinergic drugs or analeptic drugs are of no value and should not be used.

DOSAGE AND ADMINISTRATION

Usual Adult Dosage: One tablet (250 mg) three or four times daily. Initial dosage for painful musculoskeletal conditions should be two tablets (500 mg) three or four times daily. If adequate response is not obtained with this dose, it may be increased to three tablets (750 mg) three or four times daily. As improvement occurs dosage can usually be reduced.

HOW SUPPLIED

Chlorzoxazone tablets, USP, 250 mg are supplied as white, Capsule shaped tablet, debossed "A26" on one side and "250" on the other side.

They are available as:

Bottles of 60 with child-resistant closure: NDC 24470-923-60

Bottles of 100 with child-resistant closure: NDC 24470-923-10

Dispense contents with a child-resistant closure (as required) and in a tight container as defined in the USP/NF.

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

Manufactured for:

Cintex Services, LLC

Dallas, TX 75243

855-899-4237


Rev. 03/2023

L7039/01

NDC 24470-923-60

**Chlorzoxazone
Tablets, USP**


250 mg

 Rx only
60 Tablets

USUAL ADULT DOSAGE:
One or two tablets three or four times daily.
For prescribing information,
see accompanying product literature.
Dispense in a tight container as
defined in the official compendium.
Store at 20° to 25°C (68° to 77°F)
[see USP Controlled Room Temperature]
Keep out of reach of children.
Not intended for household use.

Manufactured for:
Cintex Services, LLC
Dallas, TX 75243

Rev. 08/2021 Code: L7037/00




3 24470 92360 1 8

NDC 24470-923-10

**Chlorzoxazone
Tablets, USP**


250 mg

 Rx only
100 Tablets

USUAL ADULT DOSAGE:
One or two tablets three or four times daily.
For prescribing information,
see accompanying product literature.
Dispense in a tight container as
defined in the official compendium.
Store at 20° to 25°C (68° to 77°F)
[see USP Controlled Room Temperature]
Keep out of reach of children.
Not intended for household use.

Manufactured for:
Cintex Services, LLC
Dallas, TX 75243

Rev. 08/2021 Code: L7038/00



3 24470 92310 1 3

CHLORZOAZONE			
chlorzoxazone tablet			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:24470-923
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
CHLORZOAZONE (UNII: H0DE420U8G) (CHLORZOAZONE - UNII:H0DE420U8G)		CHLORZOAZONE	250 mg
Inactive Ingredients			
Ingredient Name			Strength
DOCUSATE SODIUM (UNII: F05Q2T2JA0)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)			
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			

SILICON DIOXIDE (UNII: ETJ7Z6XBU4)

CROSCARMELOSE SODIUM (UNII: M28OL1HH48)

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE (CAPSULE SHAPE TABLET)	Size	14mm
Flavor		Imprint Code	A26;250
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:24470-923-60	60 in 1 BOTTLE; Type 0: Not a Combination Product	09/23/2021	
2	NDC:24470-923-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/23/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215158	09/23/2021	

Labeler - Cintex Services, LLC (078304114)

Revised: 5/2023

Cintex Services, LLC