


METOPROLOL TARTRATE- metoprolol tablet
NCS HealthCare of KY, LLC dba Vanguard Labs

Metoprolol Tartrate Tablets, USP

Rx Only
Prescribing Information

DESCRIPTION

Metoprolol tartrate USP, is a selective beta₁-adrenoreceptor blocking agent, available as 25 mg, 50 mg and 100 mg tablets for oral administration. Metoprolol tartrate USP is (±)-1-(Isopropylamino)-3-[p-2-methoxyethyl]phenoxy]-2-propanol L-(+)-tartrate (2:1) salt, its structural formula is:



Metoprolol tartrate USP is a white, practically odorless, crystalline powder with a molecular weight of 684.82. It is very soluble in water; freely soluble in methylene chloride, in chloroform, and in alcohol; slightly soluble in acetone; and insoluble in ether.

Each tablet for oral administration contains 25 mg, 50 mg or 100 mg of metoprolol tartrate and the following inactive ingredients: microcrystalline cellulose, lactose monohydrate, povidone, croscarmellose sodium, colloidal silicon dioxide, magnesium stearate hypromellose, titanium dioxide and polyethylene glycol.

In addition, the 50 mg product contains D&C Red #30 Aluminum Lake and the 100 mg product contains FD&C Blue #2 Aluminum Lake and FD&C Blue #1 Aluminum Lake as coloring agents.

CLINICAL PHARMACOLOGY

Mechanism of Action:

Metoprolol tartrate is a beta₁- selective (cardioselective) adrenergic receptor blocker. This preferential effect is not absolute, however, and at higher plasma concentrations, metoprolol also inhibits beta₂- adrenoreceptors, chiefly located in the bronchial and vascular musculature.

Clinical pharmacology studies have confirmed the beta-blocking activity of metoprolol in

man, as shown by (1) reduction in heart rate and cardiac output at rest and upon exercise, (2) reduction of systolic blood pressure upon exercise, (3) inhibition of isoproterenol-induced tachycardia, and (4) reduction of reflex orthostatic tachycardia.

Hypertension

The mechanism of the antihypertensive effects of beta-blocking agents has not been fully elucidated. However, several possible mechanisms have been proposed: (1) competitive antagonism of catecholamines at peripheral (especially cardiac) adrenergic neuron sites, leading to decreased cardiac output; (2) a central effect leading to reduced sympathetic outflow to the periphery; and (3) suppression of renin activity.

Angina Pectoris

By blocking catecholamine-induced increases in heart rate, in velocity and extent of myocardial contraction, and in blood pressure, metoprolol reduces the oxygen requirements of the heart at any given level of effort, thus making it useful in the long-term management of angina pectoris.

Myocardial Infarction

The precise mechanism of action of metoprolol in patients with suspected or definite myocardial infarction is not known.

Pharmacodynamics

Relative beta₁ selectivity is demonstrated by the following: (1) In healthy subjects, metoprolol is unable to reverse the beta₂-mediated vasodilating effects of epinephrine. This contrasts with the effect of nonselective (beta₁ plus beta₂) beta blockers, which completely reverse the vasodilating effects of epinephrine. (2) In asthmatic patients, metoprolol reduces FEV₁ and FVC significantly less than a nonselective beta-blocker, propranolol, at equivalent beta₁-receptor blocking doses.

Metoprolol has no intrinsic sympathomimetic activity, and membrane-stabilizing activity is detectable only at doses much greater than required for beta-blockade.

Animal and human experiments indicate that metoprolol slows the sinus rate and decreases AV nodal conduction.

Significant beta-blocking effect (as measured by reduction of exercise heart rate) occurs within 1 hour after oral administration, and its duration is dose-related. For example, a 50% reduction of the maximum effect after single oral doses of 20, 50, and 100 mg occurred at 3.3, 5, and 6.4 hours, respectively, in normal subjects. After repeated oral dosages of 100 mg twice daily, a significant reduction in exercise systolic blood pressure was evident at 12 hours. When the drug was infused over a 10-minute period, in normal volunteers, maximum beta blockade was achieved at approximately 20 minutes. Equivalent maximal beta-blocking effect is achieved with oral and intravenous doses in the ratio of approximately 2.5:1.

There is a linear relationship between the log of plasma levels and reduction of exercise heart rate. However, antihypertensive activity does not appear to be related to plasma levels. Because of variable plasma levels attained with a given dose and lack of a consistent relationship of antihypertensive activity to dose, selection of proper dosage

requires individual titration.

In several studies of patients with acute myocardial infarction, intravenous followed by oral administration of metoprolol caused a reduction in heart rate, systolic blood pressure and cardiac output. Stroke volume, diastolic blood pressure and pulmonary artery end diastolic pressure remained unchanged.

In patients with angina pectoris, plasma concentration measured at 1 hour is linearly related to the oral dose within the range of 50 to 400 mg. Exercise heart rate and systolic blood pressure are reduced in relation to the logarithm of the oral dose of metoprolol. The increase in exercise capacity and the reduction in left ventricular ischemia are also significantly related to the logarithm of the oral dose.

Pharmacokinetics

Absorption: The estimated oral bioavailability of immediate release metoprolol is about 50% because of pre-systemic metabolism which is saturable leading to non-proportionate increase in the exposure with increased dose.

Distribution: Metoprolol is extensively distributed with a reported volume of distribution of 3.2 to 5.6 L/kg. About 10% of metoprolol in plasma is bound to serum albumin. Metoprolol is known to cross the placenta and is found in breast milk. Metoprolol is also known to cross the blood brain barrier following oral administration and CSF concentrations close to that observed in plasma have been reported. Metoprolol is not a significant P-glycoprotein substrate

Metabolism: Metoprolol is primarily metabolized by CYP2D6. Metoprolol is a racemic mixture of R- and S- enantiomers, and when administered orally, it exhibits stereoselective metabolism that is dependent on oxidation phenotype. CYP2D6 is absent (poor metabolizers) in about 8% of Caucasians and about 2% of most other populations. Poor CYP2D6 metabolizers exhibit several-fold higher plasma concentrations of metoprolol than extensive metabolizers with normal CYP2D6 activity thereby decreasing metoprolol's cardioselectivity.

Elimination: Elimination of metoprolol is mainly by biotransformation in the liver. The mean elimination half-life of metoprolol is 3 to 4 hours; in poor CYP2D6 metabolizers the half-life may be 7 to 9 hours. Approximately 95% of the dose can be recovered in urine. In most subjects (extensive metabolizers), less than 5% of an oral dose are excreted as unchanged drug in the urine. In poor metabolizers, up to 30% of oral doses, may be excreted unchanged; the rest is excreted by the kidneys as metabolites that appear to have no beta blocking activity. The renal clearance of the stereo-isomers does not exhibit stereo-selectivity in renal excretion.

Special populations

Geriatric patients: The geriatric population may show slightly higher plasma concentrations of metoprolol as a combined result of a decreased metabolism of the drug in elderly population and a decreased hepatic blood flow. However, this increase is not clinically significant or therapeutically relevant.

Renal impairment: The systemic availability and half-life of metoprolol in patients with renal failure do not differ to a clinically significant degree from those in normal subjects.

Hepatic Impairment: Since the drug is primarily eliminated by hepatic metabolism, hepatic impairment may impact the pharmacokinetics of metoprolol. The elimination half-life of metoprolol is considerably prolonged, depending on severity (up to 7.2 h).

Clinical Studies:

Hypertension

In controlled clinical studies, metoprolol has been shown to be an effective antihypertensive agent when used alone or as concomitant therapy with thiazide-type diuretics, at dosages of 100 to 450 mg daily. In controlled, comparative, clinical studies, metoprolol has been shown to be as effective an antihypertensive agent as propranolol, methyldopa, and thiazide-type diuretics, to be equally effective in supine and standing positions.

Angina Pectoris

In controlled clinical trials, metoprolol, administered two or four times daily, has been shown to be an effective antianginal agent, reducing the number of angina attacks and increasing exercise tolerance. The dosage used in these studies ranged from 100 to 400 mg daily. A controlled, comparative, clinical trial showed that metoprolol was indistinguishable from propranolol in the treatment of angina pectoris.

Myocardial Infarction

In a large (1,395 patients randomized), double-blind, placebo-controlled clinical study, metoprolol was shown to reduce 3-month mortality by 36% in patients with suspected or definite myocardial infarction.

Patients were randomized and treated as soon as possible after their arrival in the hospital, once their clinical condition had stabilized and their hemodynamic status had been carefully evaluated. Subjects were ineligible if they had hypotension, bradycardia, peripheral signs of shock, and/or more than minimal basal rales as signs of congestive heart failure. Initial treatment consisted of intravenous followed by oral administration of metoprolol or placebo, given in a coronary care or comparable unit. Oral maintenance therapy with metoprolol or placebo was then continued for 3 months. After this double-blind period, all patients were given metoprolol and followed up to 1 year.

The median delay from the onset of symptoms to the initiation of therapy was 8 hours in both the metoprolol- and placebo-treatment groups. Among patients treated with metoprolol, there were comparable reductions in 3-month mortality for those treated early (≤ 8 hours) and those in whom treatment was started later. Significant reductions in the incidence of ventricular fibrillation and in chest pain following initial intravenous therapy were also observed with metoprolol and were independent of the interval between onset of symptoms and initiation of therapy.

In this study, patients treated with metoprolol received the drug both very early (intravenously) and during a subsequent 3-month period, while placebo patients received no beta-blocker treatment for this period. The study thus was able to show a benefit from the overall metoprolol regimen but cannot separate the benefit of very early intravenous treatment from the benefit of later beta-blocker therapy. Nonetheless, because the overall regimen showed a clear beneficial effect on survival without evidence of an early adverse effect on survival, one acceptable dosage regimen is the precise regimen used in the trial. Because the specific benefit of very early treatment remains to be defined however, it is also reasonable to administer the drug orally to patients at a later time as

is recommended for certain other beta blockers.

INDICATIONS AND USAGE

Hypertension

Metoprolol tartrate tablets are indicated for the treatment of hypertension. They may be used alone or in combination with other antihypertensive agents.

Angina Pectoris

Metoprolol tartrate tablets are indicated in the long-term treatment of angina pectoris.

Myocardial Infarction

Metoprolol tartrate tablets are indicated in the treatment of hemodynamically stable patients with definite or suspected acute myocardial infarction to reduce cardiovascular mortality when used alone or in conjunction with intravenous metoprolol. Oral metoprolol therapy can be initiated after intravenous metoprolol therapy or, alternatively, oral treatment can begin within 3 to 10 days of the acute event (see DOSAGE AND ADMINISTRATION, CONTRAINDICATIONS, and WARNINGS).

CONTRAINDICATIONS

Hypertension and Angina

Metoprolol tartrate tablets are contraindicated in sinus bradycardia, heart block greater than first degree, cardiogenic shock, and overt cardiac failure (see WARNINGS).

Hypersensitivity to metoprolol and related derivatives, or to any of the excipients; hypersensitivity to other beta blockers (cross sensitivity between beta blockers can occur).

Sick-sinus syndrome.

Severe peripheral arterial circulatory disorders.

Myocardial Infarction

Metoprolol is contraindicated in patients with a heart rate < 45 beats/min; second- and third-degree heart block; significant first-degree heart block (P-R interval \geq 0.24 sec); systolic blood pressure < 100 mmHg; or moderate to severe cardiac failure (see WARNINGS).

WARNINGS

Heart Failure

Beta blockers, like metoprolol, can cause depression of myocardial contractility and may precipitate heart failure and cardiogenic shock. If signs or symptoms of heart failure develop, treat the patient according to recommended guidelines. It may be necessary to lower the dose of metoprolol or to discontinue it.

Ischemic Heart Disease

Do not abruptly discontinue metoprolol therapy in patients with coronary artery disease. Severe exacerbation of angina, myocardial infarction and ventricular arrhythmias have been reported in patients with coronary artery disease following the abrupt discontinuation of therapy with beta-blockers.

When discontinuing chronically administered metoprolol, particularly in patients with coronary artery disease, the dosage should be gradually reduced over a period of 1 to 2 weeks and the patient should be carefully monitored. If angina markedly worsens or acute coronary insufficiency develops, metoprolol administration should be reinstated promptly, at least temporarily, and other measures appropriate for the management of unstable angina should be taken. Patients should be warned against interruption or discontinuation of therapy without the physicians advice.

Because coronary artery disease is common and may be unrecognized, it may be prudent not to discontinue metoprolol therapy abruptly even in patients treated only for hypertension.

Use During Major Surgery

Chronically administered beta-blocking therapy should not be routinely withdrawn prior to major surgery; however, the impaired ability of the heart to respond to reflex adrenergic stimuli may augment the risks of general anesthesia and surgical procedures.

Bradycardia

Bradycardia, including sinus pause, heart block, and cardiac arrest have occurred with the use of metoprolol. Patients with first-degree atrioventricular block, sinus node dysfunction, or conduction disorders may be at increased risk. Monitor heart rate and rhythm in patients receiving metoprolol. If severe bradycardia develops, reduce or stop metoprolol.

Exacerbation of Bronchospastic Diseases

Patients with bronchospastic diseases, should, in general, not receive beta blockers, including metoprolol. Because of its relative beta₁ selectivity, however, metoprolol may be used in patients with bronchospastic disease who do not respond to, or cannot tolerate, other antihypertensive treatment. Because beta₁ selectivity is not absolute use, the lowest possible dose of metoprolol tartrate and consider administering metoprolol in smaller doses three times daily, instead of larger doses two times daily, to avoid the higher plasma levels associated with the longer dosing interval (see DOSAGE AND ADMINISTRATION). Bronchodilators, including beta₂ agonists, should be readily available or administered concomitantly.

Diabetes and Hypoglycemia

Beta-blockers may mask tachycardia occurring with hypoglycemia, but other manifestations such as dizziness and sweating may not be significantly affected.

Pheochromocytoma

If metoprolol is used in the setting of pheochromocytoma, it should be given in combination with an alpha-blocker, and only after the alpha-blocker has been initiated. Administration of beta-blockers alone in the setting of pheochromocytoma has been associated with a paradoxical increase in blood pressure due to the attenuation of beta-mediated vasodilatation in skeletal muscle.

Thyrotoxicosis

Metoprolol tartrate tablets may mask certain clinical signs (e.g., tachycardia) of hyperthyroidism. Avoid abrupt withdrawal of beta blockade, which might precipitate a thyroid storm.

PRECAUTIONS

Risk of Anaphylactic Reactions

While taking beta-blockers, patients with a history of severe anaphylactic reaction to a variety of allergens may be more reactive to repeated challenge, either accidental, diagnostic, or therapeutic. Such patients may be unresponsive to the usual doses of epinephrine used to treat allergic reaction.

Information for Patients

Advise patients to take metoprolol regularly and continuously, as directed, with or immediately following meals. If a dose should be missed, the patient should take only the next scheduled dose (without doubling it). Patients should not discontinue metoprolol without consulting the physician.

Advise patients to (1) to avoid operating automobiles and machinery or engaging in other tasks requiring alertness until the patient's response to therapy with metoprolol has been determined; (2) to contact the physician if any difficulty in breathing occurs; (3) to inform the physician or dentist before any type of surgery that he or she is taking metoprolol.

Drug Interactions

Catecholamine-depleting drugs: Catecholamine-depleting drugs (e.g., reserpine) may have an additive effect when given with beta-blocking agents or monoamine oxidase (MAO) inhibitors. Observe patients treated with metoprolol plus a catecholamine depletor for evidence of hypotension or marked bradycardia, which may produce vertigo, syncope, or postural hypotension.

In addition, possibly significant hypertension may theoretically occur up to 14 days following discontinuation of the concomitant administration with an irreversible MAO inhibitor.

Digitalis glycosides and beta blockers: Both digitalis glycosides and beta blockers slow atrioventricular conduction and decrease heart rate. Concomitant use can increase the risk of bradycardia. Monitor heart rate and PR interval.

Calcium channel blockers: Concomitant administration of a betaadrenergic antagonist with a calcium channel blocker may produce an additive reduction in myocardial contractility because of negative chronotropic and inotropic effects.

CYP2D6 Inhibitors: Potent inhibitors of the CYP2D6 enzyme may increase the plasma concentration of metoprolol which would mimic the pharmacokinetics of CYP2D6 poor metabolizer (see Pharmacokinetics section). Increase in plasma concentrations of metoprolol would decrease the cardioselectivity of metoprolol. Known clinically significant potent inhibitors of CYP2D6 are antidepressants such as fluvoxamine, fluoxetine, paroxetine, sertraline, bupropion, clomipramine, and desipramine; antipsychotics such as

chlorpromazine, fluphenazine, haloperidol, and thioridazine; antiarrhythmics such as quinidine or propafenone; antiretrovirals such as ritonavir; antihistamines such as diphenhydramine; antimalarials such as hydroxychloroquine or quinidine; antifungals such as terbinafine.

Hydralazine: Concomitant administration of hydralazine may inhibit presystemic metabolism of metoprolol leading to increased concentrations of metoprolol.

Alpha-adrenergic agents: Antihypertensive effect of alpha-adrenergic blockers such as guanethidine, betanidine, reserpine, alpha-methyldopa or clonidine may be potentiated by beta-blockers including metoprolol. Beta- adrenergic blockers may also potentiate the postural hypotensive effect of the first dose of prazosin, probably by preventing reflex tachycardia. On the contrary, beta adrenergic blockers may also potentiate the hypertensive response to withdrawal of clonidine in patients receiving concomitant clonidine and beta-adrenergic blocker. If a patient is treated with clonidine and metoprolol concurrently, and clonidine treatment is to be discontinued, stop metoprolol several days before clonidine is withdrawn. Rebound hypertension that can follow withdrawal of clonidine may be increased in patients receiving concurrent beta-blocker treatment.

Ergot alkaloid: Concomitant administration with beta-blockers may enhance the vasoconstrictive action of ergot alkaloids.

Dipyridamole: In general, administration of a beta-blocker should be withheld before dipyridamole testing, with careful monitoring of heart rate following the dipyridamole injection.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies in animals have been conducted to evaluate carcinogenic potential. In a 2-year study in rats at three oral dosage levels of up to 800 mg/kg per day, there was no increase in the development of spontaneously occurring benign or malignant neoplasms of any type. The only histologic changes that appeared to be drug related were an increased incidence of generally mild focal accumulation of foamy macrophages in pulmonary alveoli and a slight increase in biliary hyperplasia. In a 21-month study in Swiss albino mice at three oral dosage levels of up to 750 mg/kg per day, benign lung tumors (small adenomas) occurred more frequently in female mice receiving the highest dose than in untreated control animals. There was no increase in malignant or total (benign plus malignant) lung tumors, or in the overall incidence of tumors or malignant tumors. This 21-month study was repeated in CD-1 mice, and no statistically or biologically significant differences were observed between treated and control mice of either sex for any type of tumor.

All mutagenicity tests performed (a dominant lethal study in mice, chromosome studies in somatic cells, a Salmonella/mammalianmicrosome mutagenicity test, and a nucleus anomaly test in somatic interphase nuclei) were negative.

Reproduction toxicity studies in mice, rats and rabbits did not indicate teratogenic potential for metoprolol tartrate. Embryotoxicity and/or fetotoxicity in rats and rabbits were noted starting at doses of 50 mg/kg in rats and 25 mg/kg in rabbits, as demonstrated by increases in preimplantation loss, decreases in the number of viable fetuses per dose, and/or decreases in neonatal survival. High doses were associated with some maternal toxicity, and growth delay of the offspring in utero, which was reflected in minimally lower weights at birth. The oral NOAELs for embryo-fetal development in mice, rats, and rabbits were considered to be 25, 200, and 12.5 mg/kg. This corresponds to dose levels that are approximately 0.3, 4, and 0.5 times, respectively, when based on surface area, the maximum human oral dose (8 mg/kg/day) of metoprolol tartrate. Metoprolol tartrate has been associated with reversible adverse effects on spermatogenesis starting at oral dose levels of 3.5 mg/kg in rats (a dose that is only 0.1-times the human dose, when based on surface area), although other studies have shown no effect of metoprolol tartrate on reproductive performance in male rats.

Pregnancy Category C

Upon confirming the diagnosis of pregnancy, women should immediately inform the doctor. Metoprolol has been shown to increase postimplantation loss and decrease neonatal survival in rats at doses up to 11 times the maximum daily human dose of 450 mg, when based on surface area. Distribution studies in mice confirm exposure of the fetus when metoprolol is administered to the pregnant animal. These limited animal studies do not indicate direct or indirect harmful effects with respect to teratogenicity (see Carcinogenesis, Mutagenesis, Impairment of Fertility). There are no adequate and well controlled studies in pregnant women. The amount of data on the use of metoprolol in pregnant women is limited. The risk to the fetus/mother is unknown. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers

Metoprolol is excreted in breast milk in a very small quantity. An infant consuming 1 liter of breast milk daily would receive a dose of less than 1 mg of the drug.

Fertility

The effects of Metoprolol on the fertility of human have not been studied. Metoprolol showed effects on spermatogenesis in male rats at a therapeutic dose level, but had no effect on rates of conception at higher doses in animal fertility studies (see Carcinogenesis, Mutagenesis, Impairment of Fertility).

Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

Geriatric Use

Clinical trials of metoprolol in hypertension did not include sufficient numbers of elderly patients to determine whether patients over 65 years of age differ from younger subjects in their response to metoprolol. Other reported clinical experience in elderly

hypertensive patients has not identified any difference in response from younger patients.

In worldwide clinical trials of metoprolol in myocardial infarction, where approximately 478 patients were over 65 years of age (0 over 75 years of age), no age-related differences in safety and effectiveness were found. Other reported clinical experience in myocardial infarction has not identified differences in response between the elderly and younger patients. However, greater sensitivity of some elderly individuals taking metoprolol cannot be categorically ruled out. Therefore, in general, it is recommended that dosing proceed with caution in this population.

ADVERSE REACTIONS

Hypertension and Angina

Most adverse effects have been mild and transient.

Central Nervous System: Tiredness and dizziness have occurred in about 10 of 100 patients. Depression has been reported in about 5 of 100 patients. Mental confusion and short-term memory loss have been reported. Headache, nightmares, and insomnia have also been reported.

Cardiovascular: Shortness of breath and bradycardia have occurred in approximately 3 of 100 patients. Cold extremities; arterial insufficiency, usually of the Raynaud type; palpitations; congestive heart failure; peripheral edema; and hypotension have been reported in about 1 of 100 patients. Gangrene in patients with pre-existing severe peripheral circulatory disorders has also been reported very rarely (see CONTRAINDICATIONS, WARNINGS, and PRECAUTIONS).

Respiratory: Wheezing (bronchospasm) and dyspnea have been reported in about 1 of 100 patients (see WARNINGS). Rhinitis has also been reported.

Gastrointestinal: Diarrhea has occurred in about 5 of 100 patients. Nausea, dry mouth, gastric pain, constipation, flatulence, and heartburn have been reported in about 1 of 100 patients. Vomiting was a common occurrence. Post-marketing experience reveals very rare reports of hepatitis, jaundice and nonspecific hepatic dysfunction. Isolated cases of transaminase, alkaline phosphatase, and lactic dehydrogenase elevations have also been reported.

Hypersensitive Reactions: Pruritus or rash have occurred in about 5 of 100 patients. Very rarely, photosensitivity and worsening of psoriasis have been reported.

Miscellaneous:

Peyronie's disease has been reported in fewer than 1 of 100,000 patients. Musculoskeletal pain, blurred vision, and tinnitus have also been reported.

There have been rare reports of reversible alopecia, agranulocytosis, and dry eyes. Discontinuation of the drug should be considered if any such reaction is not otherwise explicable. There have been very rare reports of weight gain, arthritis, and retroperitoneal fibrosis (relationship to metoprolol has not been definitely established).

The oculomucocutaneous syndrome associated with the beta-blocker practolol has not been reported with metoprolol.

Myocardial Infarction

Central Nervous System: Tiredness has been reported in about 1 of 100 patients. Vertigo, sleep disturbances, hallucinations, headache, dizziness, visual disturbances, confusion, and reduced libido have also been reported, but a drug relationship is not clear.

Cardiovascular: In the randomized comparison of metoprolol and placebo described in the CLINICAL PHARMACOLOGY section, the following adverse reactions were reported:

	Metoprolol	Placebo
Hypotension (systolic BP <90 mm Hg)	27.4%	23.2%
Bradycardia (heart rate <40 beats/min)	15.9%	6.7%
Second- or third-degree heart block	4.7%	4.7%
First-degree heart block (P-R \geq 0.26 sec)	5.3%	1.9%
Heart failure	27.5%	29.6%

Respiratory: Dyspnea of pulmonary origin has been reported in fewer than 1 of 100 patients.

Gastrointestinal: Nausea and abdominal pain have been reported in fewer than 1 of 100 patients.

Dermatologic: Rash and worsened psoriasis have been reported, but a drug relationship is not clear.

Miscellaneous: Unstable diabetes and claudication have been reported, but a drug relationship is not clear.

Potential Adverse Reactions

A variety of adverse reactions not listed above has been reported with other beta-adrenergic blocking agents and should be considered potential adverse reactions to metoprolol.

Central Nervous System: Reversible mental depression progressing to catatonia; an acute reversible syndrome characterized by disorientation for time and place, short-term memory loss, emotional lability, slightly clouded sensorium, and decreased performance on neuropsychometrics.

Cardiovascular: Intensification of AV block (see CONTRAINDICATIONS).

Hematologic: Agranulocytosis, nonthrombocytopenic purpura and thrombocytopenic purpura.

Hypersensitive Reactions: Fever combined with aching and sore throat, laryngospasm and respiratory distress.

Post-marketing Experience

The following adverse reactions have been reported during post-approval use of metoprolol tartrate: confusional state, an increase in blood triglycerides and a decrease in High Density Lipoprotein (HDL). Because these reports are from a population of

uncertain size and are subject to confounding factors, it is not possible to reliably estimate their frequency.

OVERDOSAGE

Acute Toxicity

Several cases of overdosage have been reported, some leading to death.

Oral LD50's (mg/kg): mice, 1,158 - 2,460; rats, 3,090 - 4,670.

Signs and Symptoms

Potential signs and symptoms associated with overdosage with metoprolol are bradycardia, hypotension, bronchospasm, myocardial infarction, cardiac failure and death.

Management

There is no specific antidote.

In general, patients with acute or recent myocardial infarction may be more hemodynamically unstable than other patients and should be treated accordingly (see WARNINGS, Myocardial Infarction).

On the basis of the pharmacologic actions of metoprolol, the following general measures should be employed:

Elimination of the Drug: Gastric lavage should be performed.

Other clinical manifestations of overdose should be managed symptomatically based on modern methods of intensive care.

Hypotension: Administer a vasopressor, e.g., levarterenol or dopamine.

Bronchospasm: Administer a beta₂-stimulating agent and/or a theophylline derivative.

Cardiac Failure: Administer digitalis glycoside and diuretic. In shock resulting from inadequate cardiac contractility, consider administration of dobutamine, isoproterenol or glucagon.

DOSAGE AND ADMINISTRATION

Hypertension

Individualize the dosage of metoprolol tablets. Metoprolol tartrate tablets should be taken with or immediately following meals.

The usual initial dosage of metoprolol tartrate tablets is 100 mg daily in single or divided doses, whether used alone or added to a diuretic. Increase the dosage at weekly (or longer) intervals until optimum blood pressure reduction is achieved. In general, the maximum effect of any given dosage level will be apparent after one week of therapy. The effective dosage range of metoprolol tartrate tablets is 100 to 450 mg per day. Dosages above 450 mg per day have not been studied. While once daily dosing is effective and can maintain a reduction in blood pressure throughout the day, lower doses (especially 100 mg) may not maintain a full effect at the end of the 24-hour period, and larger or more frequent daily doses may be required. This can be evaluated

by measuring blood pressure near the end of the dosing interval to determine whether satisfactory control is being maintained throughout the day. Beta₁ selectivity diminishes as the dose of metoprolol is increased.

Angina Pectoris

The dosage of metoprolol tartrate tablets should be individualized. Metoprolol tartrate tablets should be taken with or immediately following meals.

The usual initial dosage of metoprolol tartrate tablets is 100 mg daily, given in two divided doses. gradually increased at weekly intervals until optimum clinical response has been obtained or there is pronounced slowing of the heart rate.

The effective dosage range of metoprolol tartrate tablets is 100 to 400 mg per day. Dosages above 400 mg per day have not been studied. If treatment is to be discontinued, gradually decrease the dosage over a period of 1 to 2 weeks (see WARNINGS).

Myocardial Infarction

Early Treatment: During the early phase of definite or suspected acute myocardial infarction, treatment with metoprolol tartrate tablets can be initiated as soon as possible after the patient's arrival in the hospital. Such treatment should be initiated in a coronary care or similar unit immediately after the patient's hemodynamic condition has stabilized.

Begin treatment in this early phase should begin with the intravenous administration of three bolus injections of 5 mg of metoprolol tartrate each; the injections should be given at approximately 2 minute intervals. During the intravenous administration of metoprolol, monitor blood pressure, heart rate, and electrocardiogram.

In patients who tolerate the full intravenous dose (15 mg), initiate metoprolol tartrate tablets, 50 mg every 6 hours, 15 minutes after the last intravenous dose and continue for 48 hours. Thereafter, the maintenance dosage is 100 mg twice daily (see *Late Treatment* below).

Start patients who appear not to tolerate the full intravenous dose on metoprolol tablets either 25 mg or 50 mg every 6 hours (depending on the degree of intolerance) 15 minutes after the last intravenous dose or as soon as their clinical condition allows. In patients with severe intolerance, discontinue metoprolol (see WARNINGS).

Late Treatment: Start patients with contraindications to treatment during the early phase of suspected or definite myocardial infarction, patients who appear not to tolerate the full early treatment, and patients in whom the physician wishes to delay therapy for any other reason on metoprolol tartrate tablets, 100 mg twice daily, as soon as their clinical condition allows. Continue therapy for at least 3 months. Although the efficacy of metoprolol beyond 3 months has not been conclusively established, data from studies with other beta-blockers suggest that treatment should be continued for 1 to 3 years.

Special populations

Pediatric patients: No pediatric studies have been performed. The safety and efficacy of metoprolol in pediatric patients have not been established.

Renal impairment: No dose adjustment of metoprolol is required in patients with renal impairment.

Hepatic impairment: Metoprolol blood levels are likely to increase substantially in

patients with hepatic impairment. Therefore, metoprolol should be initiated at low doses with cautious gradual dose titration according to clinical response.

Geriatric patients (>65 years): In general, use a low initial starting dose in elderly patients given their greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

Method of administration:

For oral treatment, the tablets should be swallowed un-chewed with a glass of water. Metoprolol tartrate tablets should always be taken in standardized relation with meals. If the physician asks the patient to take metoprolol tartrate tablets either before breakfast or with breakfast, then the patient should continue taking metoprolol tartrate tablets with the same schedule during the course of therapy.

HOW SUPPLIED

Metoprolol Tartrate Tablets, USP are available containing 25 mg, 50 mg or 100 mg of metoprolol tartrate, USP.

The 25 mg tablets are white to off white film coated, round, biconvex tablets debossed with 'L150' on one side and breakline on other side. They are available as follows:

NDC 0615-8421-39	blistercards of 30 tablets
NDC 0615-8421-05	blistercards of 15 tablets
NDC 0615-8421-30	unit-dose boxes of 30 tablets

The 50 mg tablets are pink, film coated, round, biconvex tablets debossed with 'L151' on one side and breakline on other side. They are available as follows:

NDC 0615-8422-39	blistercards of 30 tablets
NDC 0615-8422-05	blistercards of 15 tablets
NDC 0615-8422-30	unit-dose boxes of 30 tablets

The 100 mg tablets are blue, film coated, round, biconvex tablets debossed with 'L152' on one side and breakline on other side. They are available as follows:

NDC 0615-8423-39	blistercards of 30 tablets
NDC 0615-8423-30	unit-dose boxes of 30 tablets

Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]

Protect from moisture.

Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.

Call your doctor for medical advice about side effects. You may report side effects to Alembic Pharmaceuticals Limited at 1-866 210 9797 or FDA at 1-800-FDA-1088.

Alembic Pharmaceuticals Limited


Manufactured in India

Manufactured for:

Alembic Pharmaceuticals Inc.
 Bedminster, NJ 07921, USA

Revised: 03/2023

PRINCIPAL DISPLAY PANEL - 25 MG BINGO CARD



VLI NDC 0515-8421-39
Metoprolol Tartrate
 Tabs USP
 25 mg
 LOT 8421-
 EXP
 8421-AA-39 v00
 QTY
 30
 Mfg By Alembic
 (NDC 62332-112-91)
 PKG BY VANGARD
 GLASGOW, KY 42141

(Alembic NDC 62332-112-91)

Metoprolol Tartrate
 Tabs USP
 25 mg

Received: _____

31	24	16	8
30	23	15	7
29	22	14	6
28	21	13	5
27	20	12	4
26	19	11	3
25	18	10	2
	17	9	1

Start Date _____ Start Time _____


Each film-coated tablet contains Metoprolol tartrate, USP 25 mg

Store at 20° to 25°C (68° to 77°F). (See USP Controlled Room Temperature.) Protect from moisture.

Usual Dosage: See package insert for full prescribing information.

FOR INSTITUTIONAL USE ONLY

	16	8	
30	23	15	7
29	22	14	6
28	21	13	5
27	20	12	4
26	19	11	3
25	18	10	2
	17	9	1



Omnicare 24

The overall configuration of this package is a trademark of Omnicare, Inc.

PRINCIPAL DISPLAY PANEL - 25 MG UNIT DOSE

8421-AA-UDI-V01

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
25 mg
LOT 8421-
6233211291

VG NDC 0615-8421-30		VG NDC 0615-8421-30	
Metoprolol Tartrate Tablets USP 25 mg		Metoprolol Tartrate Tablets USP 25 mg	
Rx only		Rx only	
QTY 30		FOR INSTITUTIONAL USE ONLY	
Each film-coated tablet contains Metoprolol tartrate, USP 25 mg		QTY 30 LOT: EXP:	
Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.] Protect from moisture.		Manufactured By	
[Empty Box]		Alembic	
LOT:		or	
EXP:		Vanguard Labs	
[Barcode]		Glasgow, KY	
8421-AA 1Z -01		42141	
00001		Usual Dosage: See package insert for full prescribing information.	
[Barcode]		[Barcode]	
6233211291		623321129130	

PRINCIPAL DISPLAY PANEL - 50 MG BINGO CARD



VLI NDC 0615-8422-39

Metoprolol Tartrate
Tabs USP
50 mg

LOT 8422-

EXP

8422-AA-39 v00

QTY
30

Mfg By Alembic

(NDC 62332-113-91)
PKG BY VANGARD
GLASGOW, KY 42141

(Alembic NDC 62332-113-91)
Metoprolol Tartrate
Tabs USP
50 mg

Rx only

30

23

15

7

29

22

14

6

28

21

13

5

27

20

12

4

26

19

11

3

25

18

10

2



24 17 9 1
The overall configuration of this package is a trademark of Omnicare, Inc.

Received: _____

31	23	15	7
30	22	14	6
29	21	13	5
28	20	12	4
27	19	11	3
26	18	10	2
25	17	9	1
Start Date	Start Time		

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Each film-coated tablet contains
Metoprolol tartrate, USP 50 mg

Store at 20° to 25°C (68° to 77°F). (See
USP Controlled Room Temperature).
Protect from moisture.

Usual Dosage: See package insert for full
prescribing information.

FOR INSTITUTIONAL USE ONLY

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

8422-AA-B-v02
Vanguard Labs
Glasgow, KY 42141
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

Pkg by Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tablet USP 50 mg
LOT 8422-

PRINCIPAL DISPLAY PANEL - 50 MG UNIT DOSE

8422-AA-UDI-V01

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
Tab USP
50 mg
LOT 8422-
6233211391

VG NDC 0615-8422-30		VG NDC 0615-8422-30	
Metoprolol Tartrate Tablets USP 50 mg		Metoprolol Tartrate Tablets USP 50 mg	
Rx only		Rx only	
QTY 30		FOR INSTITUTIONAL USE ONLY	
Each film-coated tablet contains Metoprolol tartrate, USP 50 mg		QTY 30 LOT: EXP:	
Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature]. Protect from moisture.		Manufactured By	
[Image Placeholder]		Alembic	
LOT:		or	
EXP:		Vanguard Labs	
[Barcode]		Glasgow, KY	
8422-AA 7 -01		42141	
00001		Usual Dosage: See package insert for full prescribing information.	
[Barcode]		[Barcode]	
6233211391		623321139130	

PRINCIPAL DISPLAY PANEL - 100 MG BINGO CARD



(Alembic NDC 62332-114-91)
Metoprolol Tartrate
Tablets, USP 100 mg

Rx only

8423-AA-39 v01

Mfg By Alembic
(NDC 62332-114-91)
PKG BY VANGUARD
GLASGOW, KY 42141

30	23	15	7
29	22	14	6
28	21	13	5
27	20	12	4
26	19	11	3
25	18	10	2
24	17	9	1

Received: _____

31	24	16	8
30	23	15	7
29	22	14	6
28	21	13	5
27	20	12	4
26	19	11	3
25	18	10	2
24	17	9	1

Start Date _____ Start Time _____

Each film-coated tablet contains Metoprolol tartrate, USP 100 mg

Store at 20° to 25°C (68° to 77°F). (See USP Controlled Room Temperature.) Protect from moisture.

Usual Dosage: See package insert for full prescribing information.

FOR INSTITUTIONAL USE ONLY

Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	8423-AA-B-02 Vanguard Labs Glasgow, KY 42141 LOT 8423- 6233211491
Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491
Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491
Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491
Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491
Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491
Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491	Pkg by Vanguard, Glasgow, KY 42141 Metoprolol Tartrate Tablet, USP 100 mg LOT 8423- 6233211491

Omnicare The overall configuration of this package is a trademark of Omnicare, Inc.

PRINCIPAL DISPLAY PANEL - 100 MG UNIT DOSE

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

Pkg By Vanguard, Glasgow, KY 42141
Metoprolol Tartrate
 Tab USP
 100 mg
 LOT 8423- EXP

 6233211491

VLI NDC 0615-8423-30 Metoprolol Tartrate Tablets USP 100 mg		VLI NDC 0615-8423-30 Metoprolol Tartrate Tablets USP 100 mg FOR INSTITUTIONAL USE ONLY	
QTY 30		QTY LOT: 30 EXP:	
Rx only STORE AT 20°-25°C (68°-77°F) (SEE USP CONTROLLED ROOM TEMPERATURE) Dispense in a tight, light-resistant container. Protect from moisture.			
LOT: EXP:	 8423-AA 0 00001-01		Manufactured By Alembic for Alembic NDC 623321149130 42141 Dosage: See aetnao365. sharepoint.com/sites/omnicare-search/drugs/SitePages/Home.aspx  623321149130

METOPROLOL TARTRATE

metoprolol tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0615-8421(NDC:62332-112)
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
METOPROLOL TARTRATE (UNII: W5S57Y3A5L) (METOPROLOL - UNII:GEB06NHM23)	METOPROLOL TARTRATE	25 mg

Inactive Ingredients	
Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	

LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
POVIDONE K30 (UNII: U725QWY32X)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	

Product Characteristics

Color	white (WHITE TO OFF WHITE)	Score	2 pieces
Shape	ROUND	Size	6mm
Flavor		Imprint Code	L;150
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0615-8421-39	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	02/15/2022	
2	NDC:0615-8421-05	15 in 1 BLISTER PACK; Type 0: Not a Combination Product	02/15/2022	
3	NDC:0615-8421-30	6 in 1 BOX, UNIT-DOSE	02/10/2022	
3		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202871	07/31/2016	

METOPROLOL TARTRATE

metoprolol tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0615-8422(NDC:62332-113)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
METOPROLOL TARTRATE (UNII: W5S57Y3A5L) (METOPROLOL - UNII:GEB06NHM23)	METOPROLOL TARTRATE	50 mg

Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
POVIDONE K30 (UNII: U725QWY32X)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
D&C RED NO. 30 (UNII: 2S42T2808B)	

Product Characteristics

Color	pink	Score	2 pieces
Shape	ROUND	Size	8mm
Flavor		Imprint Code	L;151
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0615-8422-39	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	02/08/2022	
2	NDC:0615-8422-05	15 in 1 BLISTER PACK; Type 0: Not a Combination Product	02/08/2022	
3	NDC:0615-8422-30	6 in 1 BOX, UNIT-DOSE	02/09/2022	
3		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202871	07/31/2016	

METOPROLOL TARTRATE

metoprolol tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0615-8423(NDC:62332-114)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
METOPROLOL TARTRATE (UNII: W5S57Y3A5L) (METOPROLOL - UNII:GEB06NHM23)	METOPROLOL TARTRATE	100 mg

Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
POVIDONE K30 (UNII: U725QWY32X)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

Product Characteristics

Color	blue	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	L;152
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0615-8423-39	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	02/07/2022	
2	NDC:0615-8423-30	6 in 1 BOX, UNIT-DOSE	02/10/2022	11/30/2023
2		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202871	07/31/2016	

Labeler - NCS HealthCare of KY, LLC dba Vanguard Labs (050052943)

Establishment

Name	Address	ID/FEI	Business Operations
NCS HealthCare of KY, LLC dba Vanguard Labs		050052943	repack(0615-8421, 0615-8422, 0615-8423)

Revised: 9/2024

NCS HealthCare of KY, LLC dba Vanguard Labs