

OMEPRAZOLE- omeprazole magnesium tablet, delayed release
P & L Development, LLC

OMEPRAZOLE MAGNESIUM DELAYED-RELEASE TABLETS

ACTIVE INGREDIENT

(in each tablet)

Omeprazole delayed-release tablet 20 mg
(equivalent to 20.6 mg omeprazole magnesium)

Purpose

Acid reducer

Use

- treats frequent heartburn (occurs ***2 or more*** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

taking a prescription drug.

Acid reducers may interact with certain drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take everyday for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- Keep the carton. It contains important information.
- store at 20° to 25°C (68° to 77°F) and protect from moisture

Inactive ingredients

acetyl tributyl citrate, colloidal silicon dioxide, corn starch, croscarmellose sodium,

hydroxypropyl cellulose, hypromellose 2910, magnesium stearate, methacrylic acid copolymer type C, microcrystalline cellulose, polyethylene glycol 400, polyethylene glycol 3350, polyethylene glycol 8000, polyvinyl alcohol, red iron oxide, sucrose, talc, titanium dioxide, triethyl citrate

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

Compare to the active ingredient in Prilosec OTC®*

Omeprazole Delayed-Release Tablets 20 mg

ACID REDUCER

Treats FREQUENT Heartburn 24 Hr

One-14-Day Course of Treatment

14 Tablets

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole delayed-release tablet 20 mg (equivalent to 20.5 mg omeprazole magnesium).....Acid reducer</p> <p>Purpose</p> <p>Acid reducer</p> <p>Use</p> <ul style="list-style-type: none"> ■ Treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect <p>Warnings</p> <p>Allergy alert: Do not use if you are allergic to omeprazole</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating, pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain <p>These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p>	<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ swallow whole. Do not chew or crush tablets. <p>Repeated 14-Day Courses (if needed)</p> <ul style="list-style-type: none"> ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor <p>■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.</p>	<p>Drug Facts (continued)</p> <p>Other information</p> <ul style="list-style-type: none"> ■ Read the directions and warnings before use ■ Keep the carton. It contains important information. ■ Store at 20° to 25° C (68° to 77° F) and protect from moisture <p>Inactive ingredients: acetyl tributyl citrate, colloidal silicon dioxide, corn starch, croscarmellose sodium, hydroxypropyl cellulose, hypromellose 2910, magnesium stearate, methacrylic acid copolymer type C, microcrystalline cellulose, polyethylene glycol 400, polyvinyl alcohol, red iron oxide, sucrose, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? Call 1-877-753-3835 Monday-Friday 9AM-5PM EST</p> <p>This product is not affiliated with, manufactured by or produced by the makers or owners of Prilosec OTC®.</p> <p>Manufactured for: P & L Development LLC Westbury, NY 11590</p> <p>PLD-A000A Rev. 03/20 FC000000</p> <p>TAMPER EVIDENT: Do not use if caplin is missing or full seal under cap, printed with "SEALED FOR YOUR PROTECTION" is missing, open or broken.</p>	<p style="text-align: right;">NDC 59726-297-14 ¹Compare to the active ingredient in Prilosec OTC®</p> <h2 style="text-align: center;">Omeprazole</h2> <p style="text-align: center;">Delayed-release tablets 20 mg Acid reducer</p> <p style="text-align: center;"><i>Treats FREQUENT heartburn 24 HR</i></p> <p style="text-align: center;">14 Tablets <i>One-14-day course of treatment</i> <i>May take 1 to 4 days for full effect</i></p>	<p>Tips for Managing Heartburn</p> <ul style="list-style-type: none"> ■ Do not lie flat or bend over after eating ■ Do not wear tight-fitting clothing around the stomach ■ Do not eat before bedtime ■ Raise the head of your bed ■ Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables ■ Eat slowly and avoid big meals ■ If overweight, lose weight ■ Quit smoking
<p style="text-align: center;">Lot No.: Exp. Date:</p> <div style="text-align: center;">  <p>3 59726 29714 1</p> </div>				

Omeprazole Tablets

PACKAGE LABEL, PRINCIPAL DISPLAY PANEL

Compare to the active ingredient in Prilosec OTC®*

Omeprazole Delayed-Release Tablets 20 mg

ACID REDUCER

SWALLOW - DO NOT CHEW

Treats FREQUENT Heartburn 24 HR

14 Tablets

One-14-Day Course of Treatment

Coated with Wildberry Flavor

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole delayed-release tablet 20 mg (equivalent to 20.6 mg omeprazole magnesium).....Acid reducer</p> <p>Purpose Acid reducer</p> <p>Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings Allergy alert: Do not use if you are allergic to omeprazole Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p>	<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours 14-Day Course of Treatment ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ swallow whole. Do not chew, crush, or suck tablets. Repeated 14-Day Courses (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.</p>	<p>Drug Facts (continued)</p> <p>Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20° to 25°C (68° to 77°F) and protect from moisture</p> <p>Inactive ingredients acetyl tributyl citrate, colloidal silicon dioxide, corn starch, croscarmellose sodium, dextrose, FD&C Blue #2, FD&C Red #40, flavor, hydroxypropyl cellulose, hypromellose 2910, magnesium stearate, methacrylic acid copolymer type C, microcrystalline cellulose, polyethylene glycol 400, polyethylene glycol 3350, polyethylene glycol 8000, polyvinyl alcohol, polyvinyl glycol, sucrose, talc, titanium dioxide, triacetin, triethyl citrate</p> <p>Questions or comments? Call 1-877-753-3555 Monday-Friday 9AM-5PM EST</p> <p>*This product is not affiliated with, manufactured by or produced by the makers or owners of Prilosec OTC®.</p> <p>Manufactured for: P & L Development LLC Westbury, NY 11590</p> <p>P.L.D.-A000A Rev. 04/20 FC000000</p> <p>TAMPER EVIDENT: Do not use if carton is opened or foil seal under cap, printed with "SEALED FOR YOUR PROTECTION" is missing, open or broken.</p>	<p>NDC 59726-737-14 †Compare to the active ingredient in Prilosec OTC®</p> <h2>Omeprazole</h2> <p>Delayed-release tablets 20 mg Acid reducer</p> <p>Treats FREQUENT heartburn 24 HR SWALLOW-DO NOT CHEW</p> <p>14 Tablets One-14-day course of treatment May take 1 to 4 days for full effect</p> <p>Coated with Wildberry Flavor</p>	<p>Tips for Managing Heartburn</p> <ul style="list-style-type: none"> ■ Do not lie flat or bend over after eating ■ Do not wear tight-fitting clothing around the stomach ■ Do not eat before bedtime ■ Raise the head of your bed ■ Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables ■ Eat slowly and avoid big meals ■ If overweight, lose weight ■ Quit smoking
<p>Lot No: Exp. Date:</p>  <p>3 5 9 7 2 6 7 3 7 1 4 2</p>				

Omeprazole Wildberry Tablets

OMEPRAZOLE			
omeprazole magnesium tablet, delayed release			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59726-297
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	OMEPRAZOLE MAGNESIUM (UNII: 426QFE7XLK) (OMEPRAZOLE - UNII:KG60484QX9)	OMEPRAZOLE	20.6 mg

Inactive Ingredients

Ingredient Name	Strength
ACETYLTRIBUTYL CITRATE (UNII: 0ZBX0N59RZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
HYPROMELLOSE 2910 (15 MPA.S) (UNII: 36SFW2JZ0W)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
POLYETHYLENE GLYCOL 8000 (UNII: Q662QK8M3B)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	red	Score	no score
Shape	OVAL (capsule-shaped)	Size	13mm
Flavor		Imprint Code	OM;20
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59726-297-14	1 in 1 CARTON	06/02/2020	12/31/2025
1		14 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:59726-297-28	2 in 1 CARTON	06/02/2020	12/31/2025
2		14 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:59726-297-42	3 in 1 CARTON	06/02/2020	12/31/2025
3		14 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206582	06/02/2020	12/31/2025

OMEPRAZOLE

omeprazole magnesium tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59726-737
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OMEPRAZOLE MAGNESIUM (UNII: 426QFE7XLK) (OMEPRAZOLE - UNII:KG60484QX9)	OMEPRAZOLE	20.6 mg

Inactive Ingredients

Ingredient Name	Strength
ACETYLTRIBUTYL CITRATE (UNII: 0ZBX0N59RZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
HYPROMELLOSE 2910 (15 MPA.S) (UNII: 365FW2JZ0W)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
POLYETHYLENE GLYCOL 8000 (UNII: Q662QK8M3B)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
DEXTROSE (UNII: IY9XDZ35W2)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
TRIACETIN (UNII: XHX3C3X673)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	

Product Characteristics

Color	purple	Score	no score
Shape	OVAL	Size	13mm
Flavor	BERRY	Imprint Code	OM;20
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59726-737-14	1 in 1 CARTON	06/02/2020	12/31/2025
1		14 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:59726-737-28	2 in 1 CARTON	06/02/2020	12/31/2025
2		14 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:59726-737-42	3 in 1 CARTON	06/02/2020	12/31/2025
3		14 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206582	06/02/2020	12/31/2025

Labeler - P & L Development, LLC (800014821)

Revised: 4/2023

P & L Development, LLC