

PHARBETOL- acetaminophen 500mg tablet, coated
Ulai Health LLC

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- the common cold
- headache
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or a pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease.

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">• take 2 caplets every 6 hours while symptoms last• do not take more than 6 caplets in 24 hours, unless directed by a doctor• do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- store at 20-25 °C (68-77 °F)

Inactive ingredients

hypromellose, povidone, pregelatinized corn starch, sodium starch glycolate, stearic acid, talc, triacetin

Questions or comments?

(866) 562-2756 (Mon - Fri 8 AM to 4 PM EST)

PHARBEST

NDC 73057-350-11

Manufactured in the USA

Extra Strength *Compare to the active ingredient in Extra Strength Tylenol® Caplet

Contains no Aspirin

PHARBETOL

Acetaminophen 500 mg each

Pain Reliever/Fever Reducer

1000 CAPLETS

PHARBEST
NDC 73057-350-11

 **Manufactured in the USA**

Extra Strength
Contains no Aspirin

*Compare to the active ingredient in Extra Strength Tylenol® Caplet

PHARBETOL
Acetaminophen 500 mg each
Pain Reliever/Fever Reducer

1000 CAPLETS 

THIS PACKAGE FOR HOUSEHOLD WITHOUT YOUNG CHILDREN

TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each caplet)
Acetaminophen 500 mg Pain reliever/fever reducer

Uses
Temporarily relieves minor aches and pains due to: ■ the common cold ■ headache ■ backache ■ minor pain of arthritis ■ toothache ■ muscular aches ■ premenstrual and menstrual cramps ■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or a pharmacist.
■ if you are allergic to acetaminophen or any of the inactive ingredients in this product
Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if ■ Pain gets worse or lasts more than 10 days ■ Fever gets worse or lasts more than 3 days ■ Nausea or swelling is present ■ New symptoms occur These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
■ do not take more than directed (see overdose warning)
■ take 2 caplets every 6 hours while symptoms last
■ do not take more than 6 caplets in 24 hours, unless directed by a doctor
■ do not use for more than 10 days unless directed by a doctor

Children under 12 years ask a doctor

Other information
■ store at 20-25°C (68-77°F)

Inactive ingredients hypromellose, povidone, pregelatinized corn starch, sodium starch glycolate, stearic acid, talc, triacetin

Questions or comments? (866) 562-2756 (Mon-Fri 8 AM to 4 PM EST)

*This product is not manufactured or distributed by McNeil Consumer Healthcare Division of McNeil-PPC, Inc., owner of the registered trademark, Tylenol®.

Org. 06/19
3 7 3 0 5 7 1 3 5 0 1 1 5
Distributed by
ULAI HEALTH, LLC
Alexandria, VA 22312

PHARBETOL

acetaminophen 500mg tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73057-350
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	

STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE (Capsule shaped tablet)	Size	18mm
Flavor		Imprint Code	PH016
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73057-350-11	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/08/2019	
2	NDC:73057-350-08	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/08/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/08/2019	

Labeler - Ulai Health LLC (081181535)