### GOOD NEIGHBOR PHARMACY SINUS RELIEF- acetaminophen, guaifenesin, phenylephrine hcl capsule, coated Amerisource Bergen

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

### Amerisource Bergen Sinus Pressure & Pain Relief Drug Facts

### Active ingredients (in each caplet)

Acetaminophen 325 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

### Purpose

Pain reliever

Expectorant

Nasal decongestant

### Uses

- temporarily relieves:
- nasal congestion
- headache
- minor aches and pains
- sinus congestion and pressure
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

### Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

### When using this product

### do not use more than directed

### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or persistent headache. These could be signs of a serious condition.

### If pregnant or breast-feeding,

ask a health professional before use.

### Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symp

### **Directions**

- do not take more than directed (see overdose warning)
- do not take more than 10 caplets in any 24-hour period
- adults and children 12 years and older: take 2 caplets every 4 hours
- children under 12 years of age: do not use

### Other information

- each caplet contains: sodium 4 mg
- store at 20-25°C (68-77°F)

### **Inactive ingredients**

croscarmellose sodium, crospovidone, FD&C yellow #5 aluminum lake (tartrazine), FD&C yellow #6 aluminum lake, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, silicon dioxide, stearic acid, talc, titanium dioxide

### Questions or comments?

1-800-719-9260

### Package/Label Principal Display Panel

Compare to  $Mucinex^{\ensuremath{\mathbb{R}}}$  Sinus- $Max^{\ensuremath{\mathbb{R}}}$  Pressure & Pain active ingredients

See New Warnings

Sinus Relief Pressure & Pain

FOR AGES 12+

acetaminophen – pain reliever

guaifenesin – expectorant

phenylephrine HCl – nasal decongestant

Maximum Strength

Relieves Sinus Pressure & Congestion – Relieves Headache

Thins & Loosens Mucus

20 caplets

actual size



# Sinus Relief Pressure & Pain





Compare to Mucinex® Sinus-Max® Pressure & Pain active ingredients\*

NDC 46122-271-60

See New Warnings

# Sinus Relief Pressure & Pain

**acetaminophen** • pain reliever guaifenesin • expectorant phenylephrine HCl • nasal decongestant

### Maximum Strength\*\*

- Relieves Sinus Pressure & Congestion Relieves Headache Thins & Loosens Mucus

20 caplets

actual size







\*Good Neighbor Pharmacy®
Sinus Relief Pressure & Pain is
not manufactured or
distributed by Reckitt
Benckiser, distributor of
Mucinex® Sinus-Max®



Distributed By
AmerisourceBergen
1300 Morris Drive
Chesterbrook, PA 19087
Visit us at
w.goodneighborpharmacy.con

### Drug Facts

# Active ingredients (in each caplet)

### Purpose

Acetaminophen 325 mg......Pain reliever Guaifenesin 200 mg.....Expectorant Phenylephrine HCl 5 mg...Nasal decongestant

### Uses

- temporarily relieves: nasal congestion
   headache minor aches and pains
  - sinus congestion and pressure
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

### Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek
medical help right away.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

### Ask a doctor before use if you have

- liver disease heart disease diabetes
- high blood pressure
   thyroid disease

### Drug Facts (continued)

■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough that occurs with too much phleqm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

# Directions ■ do not take more than directed (see overdose warning)

- do not take more than 10 caplets in any 24-hour period
- adults and children 12 years and older: take 2 caplets every 4 hours
- children under 12 years of age: do not use

### Other information

- each caplet contains: sodium 4 mg
- store at 20-25°C (68-77°F)

Inactive ingredients croscarmellose sodium, crospovidone, FD&C yellow #5 aluminum lake (tartrazine), FD&C yellow #6 aluminum lake, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, silicon dioxide, stearic acid, talc, titanium dioxide



### Do not use if blister unit is broken or torn

\*\*Per 4-hour dose, dose every 4 hours Keep carton for complete product information.

### GOOD NEIGHBOR PHARMACY SINUS RELIEF

acetaminophen, guaifenesin, phenylephrine hcl capsule, coated

#### **Product Information**

Product TypeHUMAN OTC DRUGItem Code (Source)NDC:46122-271

Route of Administration ORAL

# Active Ingredient/Active Moiety Ingredient Name Basis of Strength ACETAMINOPHEN (UNII: 36209 ITL9D) (ACETAMINOPHEN - UNII:36209 ITL9D) ACETAMINOPHEN (UNII: 495W7451VQ) (GUAIFENES IN - UNII:495W7451VQ) GUAIFENES IN (UNII: 495W7451VQ) (GUAIFENES IN - UNII:495W7451VQ) PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - PHENYLEPHRINE HYDROCHLORIDE 5 mg

Inactive Ingredients		
Ingredient Name	Strength	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)		
CROSPOVIDONE (15 MPA.S AT 5%) (UNII: 68401960MK)		
MALTO DEXTRIN (UNII: 7CVR7L4A2D)		
MICRO CRYSTALLINE CELLULO SE (UNII: OP1R32D61U)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)		
PO VIDONE, UNSPECIFIED (UNII: FZ989GH94E)		
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)		
STEARIC ACID (UNII: 4ELV7Z65AP)		
TALC (UNII: 7SEV7J4R1U)		
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)		

<b>Product Characteristic</b>	roduct Characteristics			
Color	ORANGE	Score	no score	
Shape	CAPSULE	Size	19 mm	
Flavor		Imprint Code		

### **Contains**

Packaging					
# Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>		
1 NDC:46122-271-60	10 in 1 CARTON	11/06/2014			
1	$2\ \mbox{in}\ 1\ \mbox{BLISTER}$ PACK; Type $0$ : Not a Combination Product				

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC monograph final	part341	11/06/2014			

## Labeler - Amerisource Bergen (007914906)

Revised: 12/2019 Amerisource Bergen