CHILDRENS PAIN AND FEVER - acetaminophen suspension Aurohealth LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient(in each 5 mL)

Acetaminophen, USP 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen.

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for more than 10 days for pain unless directed by a doctor.
- for more than 3 days for fever unless directed by a doctor.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dose (see overdose warning).

Stop using and ask a doctor if

- symptoms do not improve
- new symptoms occur
- pain or fever persists or gets worse

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

^{*} or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

• each 5 mL contains: sodium 6 mg

- store at 20°-25°C (68°-77°F)
- do not use if printed seal over cap is torn or missing
- see bottom panel for lot number and expiration date

Inactive ingredients

anhydrous citric acid, butylparaben, carboxymethylcellulose sodium, flavor, glycerin, high fructose corn syrup, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

Questions or comments?

1-855-274-4122

Distributed by: Aurohealth LLC. 2572 Brunswick Pike Lawrenceville, NJ 08648

PACKAGE LABEL-PRINCIPAL DISPLAY PANEL 4 FL OZ (118 mL)

NDC 58602-503-24
PrimaryHealth
Ages 2-11 years
Children's
Pain &
Fever
SUSPENSION LIQUID
Pain Reliever & Fever Reducer
ACETAMINOPHEN
ORAL SUSPENSION
160 mg per 5 mL

- Alcohol Free
- Aspirin Free
- Ibuprofen Free
- Gluten Free

Dye-Free

Cherry Flavor

Use ONLY enclosed dosage cup DO NOT USE IF PRINTED SEAL OVER CAP IS TORN OR MISSING

4 FL OZ (118 mL)



Drug Facts Active ingredient (in each 5 mL)

Purpose

Uses temporarily: = reduces fever = relieves minor aches and pains due to:

= the common cold = flu = headache = sore throat = toothache

Warnings Liver warning: This product contains acetaminophen. Severe liver
damage may occur if your child takes = more than 5 doses in 24 hours, which
is the maximum daily amount = with other drugs containing acetaminophen.

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may
include: = skin reddening = blisters = resh [16 skin reactions course stop use Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: mskin reddening misiters mash. If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. Do not use myth any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. mfor more than 10 days for pain unless directed by a doctor. mfor more than 3 days for fever unless directed by a doctor. mfor more than 3 days for fever unless directed by a doctor. mfor before use if your child has liver disease. Ask a doctor or pharmacist before use if your child has liver disease. Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin. When using this product do not exceed recommended dose (see overdose warning). Stop using and ask a doctor if msymptoms do not improve mew symptoms occur pain or fever persists or gets worse Keep out of reach of children. Overdose ■ pain or fever persists or gets worse **Keep out of reach of children. Overdose** warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Directions athis product does not contain directions or complete warnings for adult use = do not give more than directed (see overdose warning) = shake well before using = mL = milliliter = find right dose on chart below.

If possible, use weight to dose; otherwise, use age. ■ remove the child protective cap and squeeze your child's dose into the dosing cup ■ repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours

[Weight (lb)	Age (yr)	Dose (mL)*	
[under 24	under 2 years	ask a doctor	
[24-35	2-3 years	5 mL	
[36-47	4-5 years	7.5 mL	
[48-59	6-8 years	10 mL	
	60-71	9-10 years	12.5 mL	
ſ	72-95	11 years	15 mL	

or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information meach 5 mL contains: sodium 6 mg ■ store at 20°-25°C (68°-77°F) ■ do not use if printed seal over cap is torn or missing

Inactive ingredients anhydrous citric acid, butylparaben, carboxymethylcellulose sodium, flavor, glycerin, high fructose corn syrup, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

Questions or comments? © 1 - 855 - 274 - 4122 🗟

Distributed by: Aurohealth LLC. 2572 Brunswick Pike, Lawrenceville, NJ 08648

E E E

PACKAGE LABEL-PRINCIPAL DISPLAY PANEL 4 FL OZ (118 mL)

NDC 58602-503-24

PrimaryHealth

Compare To Children's

TYLENOL® Oral Suspension

active ingredient**

Ages 2-11 years

Children's

Pain &

Fever

SUSPENSION LIQUID

Pain Reliever & Fever Reducer

ACETAMINOPHEN

ORAL SUSPENSION

160 mg per 5 mL

- Alcohol Free
- Aspirin Free
- Ibuprofen Free
- Gluten Free

Dve-Free

Cherry Flavor

Use ONLY enclosed dosage cup

4 FL OZ (118 mL)



CHILDRENS PAIN AND FEVER

acetaminophen suspension

Product Information Product Type HUMAN OTC DRUG Item Code (Source) NDC:58602-503 Route of Administration ORAL

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINO PHEN (UNII: 36209 ITL9 D) (ACETAMINO PHEN - UNII: 36209 ITL9 D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients	
Ingredient Name	Strength
ANHYDRO US CITRIC ACID (UNII: XF417D3PSL)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)	
CHERRY (UNII: BUC5I9595W)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6 UN3QB6S)	
MICRO CRYSTALLINE CELLULO SE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics			
Color	WHITE (White to off-white)	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging			
# Item Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:58602-503-24	1 in 1 CARTON	09/24/2018	
1	118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part343	09/24/2018	

Labeler - Aurohealth LLC (078728447)

Establishment			
Name	Address	ID/FEI	Business Operations
Aurohealth LLC		078728447	MANUFACTURE(58602-503)

Revised: 11/2019 Aurohealth LLC