

ELECTROLYTE SOLUTION WITH DEXTROSE- dexolyte injection, solution Covetrus

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Electrolyte solution with dextrose

INDICATIONS:

For use in conditions associated with fluid and electrolyte loss, such as dehydration, shock, vomiting and diarrhea, particularly when an immediate source of energy is also indicated.

CONTRAINDICATIONS:

Do not administer intraperitoneally to horses.

CAUTION:

This product contains no preservatives.
Use entire contents when first opened. Discard any unused solution.

DOSAGE AND ADMINISTRATION:

Warm solution to body temperature and administer slowly (10 to 30 mL per minute) by intravenous or intraperitoneal injection, using strict aseptic procedures.

Adult Cattle and Horses - 1000 to 2000 mL

Calves, Ponies and Foals - 500 to 1000 mL

Adult Sheep and Swine - 500 to 1000 mL

These are suggested dosages. The actual amount and rate of fluid administration must be judged by the veterinarian in relation to the condition being treated and the clinical response of the animal, being careful to avoid overhydration.

TAKE TIME OBSERVE LABEL DIRECTIONS

Questions? (855) 724-3461

Reorder #069174

18-804

RMS 92-1175

AH-069174-01

Rev: 0719

Lot No.

Exp. Date

Manufactured by
Nova-Tech, Inc.
Grand Island, NE 68801

Distributed by:
Covetrus North America
400 Metro Place North
Dublin, OH 43017

Volume: 16.907 fl oz (500 mL)

Assembled in USA

FOR ANIMAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

Caution:

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

COMPOSITION:

Each 500 mL of sterile aqueous solution contains:

Dextrose . H₂O.....12.50 g

Sorbitol.....12.50 g

Sodium lactate.....3.95 g

Sodium chloride.....2.40 g

Potassium chloride.....0.37 g

Magnesium chloride . 6H₂O.....0.21 g

Calcium chloride . 2H₂O.....0.19 g

Milliequivalents per liter

Cations

Sodium.....153 mEq/L

Potassium.....9 mEq/L

Calcium.....6 mEq/L

Magnesium.....4 mEq/L

Anions

Chloride.....101 mEq/L

Lactate.....71 mEq/L

Osmolarity (calc.) 602 mOsmol/L

Store between 15°-30°C (59°-86F).



Indicaciones: Para usar en condiciones asociadas con pérdidas de fluidos y electrolitos, como deshidratación, shock, vómitos y diarrea, especialmente cuando esté indicada también una fuente inmediata de energía.

Contraindicaciones: No administrar por vía intraperitoneal en caballos

Precaución: Este producto no contiene agentes conservantes. Usar todo el contenido después de abrir. Descartar cualquier solución no utilizada.

Dosis y administración: Elevar la temperatura de la solución a la del cuerpo y administrar lentamente (de 10 a 30 mL por minuto) por inyección intravenosa o intraperitoneal, usando procedimientos de asepsia estricta.

Bovinos y caballos adultos - 1000 a 2000 mL

Terminos/becerras, ponis y potrillos - 500 a 1000 mL

Ovejas y cerdos adultos - 500 a 1000 mL

Estas son dosis sugeridas. La cantidad y velocidad reales de la administración de fluidos en efecto deben ser juzgadas por el veterinario en relación a la condición que se trata y a la respuesta clínica del animal, con cuidado de evitar la sobrehidratación.

Table with 2 columns: Component and Milliequivalents per liter. Includes Dextrose, Sorbitol, Sodium lactate, Sodium chloride, Potassium chloride, Magnesium chloride, Calcium chloride, Chloride, and Lactate.

Almacénese entre 15° y 30°C (59° y 86°F). Tomarse el tiempo requerido para observar las direcciones de la etiqueta. N° para nuevo pedido 069174



NDC: 11695-6413-5

Electrolyte solution with dextrose

ES Solución electrolítica con dextrosa

For animal use only
Keep out of reach of children

Solamente para uso animal
Manténgase fuera del alcance de los niños

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Precaución: Leyes federales restringen el uso de esta droga por o bajo las órdenes de un veterinario con licencia.

Volume: 16.907 fl oz (500 mL)

Indications: For use in conditions associated with fluid and electrolyte loss, such as dehydration, shock, vomiting and diarrhea, particularly when an immediate source of energy is also indicated.

Contraindications: Do not administer intraperitoneally to horses.

Caution: This product contains no preservatives. Use entire contents when first opened. Discard any unused solution.

Dosage and administration: Warm solution to body temperature and administer slowly (10 to 30 mL per minute) by intravenous or intraperitoneal injection, using strict aseptic procedures.

Adult cattle and horses - 1000 to 2000 mL

Calves, ponies and foals - 500 to 1000 mL

Adult sheep and swine - 500 to 1000 mL

These are suggested dosages. The actual amount and rate of fluid administration must be judged by the veterinarian in relation to the condition being treated and the clinical response of the animal, being careful to avoid overhydration.

Table with 2 columns: Component and Milliequivalents per liter. Includes Dextrose, Sorbitol, Sodium lactate, Sodium chloride, Potassium chloride, Magnesium chloride, Calcium chloride, Chloride, and Lactate.

Store between 15°-30°C (59°-86°F).

Questions? (855) 724-3461

Reorder #069174

18-804

RMS 92-1175

AH-069174-01

REV: 0719

Lot No.

Distributed by: Covetrus North America 400 Metro Place North Dublin, OH 43017 covetrus.com

Manufactured by: Nova-Tech, Inc. Grand Island, NE 68801

Exp. Date



NDC: 11695-6413-1

Electrolyte solution with dextrose

ES Solución electrolítica con dextrosa

For animal use only
Keep out of reach of children

Solamente para uso animal
Manténgase fuera del alcance de los niños

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Precaución: Leyes federales restringen el uso de esta droga por o bajo las órdenes de un veterinario con licencia.

Volume: 33.814 fl oz (1000 mL)

Indications: For use in conditions associated with fluid and electrolyte loss, such as dehydration, shock, vomiting and diarrhea, particularly when an immediate source of energy is also indicated.

Contraindications: Do not administer intraperitoneally to horses.

Caution: This product contains no preservatives. Use entire contents when first opened. Discard any unused solution.

Dosage and administration: Warm solution to body temperature and administer slowly (10 to 30 mL per minute) by intravenous or intraperitoneal injection, using strict aseptic procedures.

Adult cattle and horses - 1000 to 2000 mL

Calves, ponies and foals - 500 to 1000 mL

Adult sheep and swine - 500 to 1000 mL

These are suggested dosages. The actual amount and rate of fluid administration must be judged by the veterinarian in relation to the condition being treated and the clinical response of the animal, being careful to avoid overhydration.

Table with 2 columns: Component and Milliequivalents per liter. Includes Dextrose, Sorbitol, Sodium lactate, Sodium chloride, Potassium chloride, Magnesium chloride, Calcium chloride, Chloride, and Lactate.

Store between 15°-30°C (59°-86°F).

Questions? (855) 724-3461

Reorder #069175

18-804

RMS 92-1175

AH-069175-01

REV: 0719

Lot No.

Distributed by: Covetrus North America 400 Metro Place North Dublin, OH 43017 covetrus.com

Manufactured by: Nova-Tech, Inc. Grand Island, NE 68801

Exp. Date



ELECTROLYTE SOLUTION WITH DEXTROSE

dexolyte injection, solution

Product Information

| | | | |
|--------------------------------|------------------------------|---------------------------|----------------|
| Product Type | PRESCRIPTION ANIMAL DRUG | Item Code (Source) | NDC:11695-6413 |
| Route of Administration | INTRAVENOUS, INTRAPERITONEAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|----------------------|-------------------|
| Dextrose Monohydrate (UNII: LX22YL083G) (Anhydrous Dextrose - UNII:5SLOG7R0OK) | Dextrose Monohydrate | 12.50 g in 500 mL |
| Sorbitol (UNII: 506T60A25R) (Sorbitol - UNII:506T60A25R) | Sorbitol | 12.50 in 500 mL |
| Sodium Lactate (UNII: TU7HW0W0QT) (Sodium Cation - UNII:LYR4M0NH37) | Sodium Lactate | 3.95 in 500 mL |
| Sodium Chloride (UNII: 451W47IQ8X) (Chloride Ion - UNII:Q32ZN48698) | Sodium Chloride | 2.40 in 500 mL |
| Potassium Chloride (UNII: 660YQ98I10) (Potassium Cation - UNII:295O53K152) | Potassium Chloride | 0.37 in 500 mL |
| Magnesium Chloride (UNII: 02F3473H9O) (Magnesium Cation - UNII:T6V3LHY838) | Magnesium Chloride | 0.21 in 500 mL |
| Calcium Chloride (UNII: M4I0D6VV5M) (Calcium Cation - UNII:2M83C4R6ZB) | Calcium Chloride | 0.19 in 500 mL |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|------------------------------|----------------------|--------------------|
| 1 | NDC:11695-6413-5 | 500 mL in 1 BOTTLE, PLASTIC | | |
| 2 | NDC:11695-6413-1 | 1000 mL in 1 BOTTLE, PLASTIC | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------------|--|----------------------|--------------------|
| unapproved drug other | | 06/01/2016 | |

Labeler - Covetrus (603750329)