

DAYTIME SINUS- acetaminophen, phenylephrine hydrochloride capsule, liquid filled
PuraCap Pharmaceutical LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

DayTime Sinus

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg

Phenylephrine HCl 5 mg

Purposes

Pain reliever/Fever reducer

Nasal decongestant

Uses

- temporarily relieves symptoms due to the common cold
- minor aches and pains
- headache
- fever
- nasal congestion
- sinus congestion & pressure
- temporarily relieves symptoms due to hay fever or other upper respiratory allergies
- minor aches and pains
- headache
- nasal congestion
- sinus congestion & pressure

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- you get nervous, dizzy or sleepless

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 4 doses per 24 hrs

adults and children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- store at room temperature 15°-30°C (59°-86°F)

Inactive ingredients

FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol special and white edible ink

Questions or comments?

Call toll free: 1-855-215-8180

PRINCIPAL DISPLAY PANEL

DayTime Sinus 24 SOFTGELS

NDC 51013-157-04

Compare to Vicks® QlearQuil® Daytime Sinus & Congestion active ingredients*

MARKET BASKET
"MORE FOR YOUR DOLLAR"

Multi-Symptom Non-Drowsy
DayTime Sinus
ACETAMINOPHEN - Pain reliever/Fever reducer
PHENYLEPHRINE HCl - Nasal decongestant

Compare to
Vicks® QlearQuil™
DayTime Sinus & Congestion
active ingredients*
NDC 51013-157-04

Multi-Symptom Non-Drowsy
DayTime Sinus
ACETAMINOPHEN - Pain reliever/Fever reducer
PHENYLEPHRINE HCl - Nasal decongestant
Relieves:
• Minor aches & pains
• Fever & headache
• Nasal congestion
• Sinus congestion & pressure
Alcohol & Antihistamine Free

24 SOFTGELS

MARKET BASKET
"MORE FOR YOUR DOLLAR"

Compare to
Vicks® QlearQuil™
DayTime Sinus & Congestion
active ingredients*
NDC 51013-157-04

Actual size

24 SOFTGELS

KEEP OUTER CARTON FOR COMPLETE
WARNINGS AND PRODUCT INFORMATION

LOT NO: _____
EXP DATE: _____

DISTRIBUTED BY:
DEMOLAS SUPERMARKETS INC.
TEWKSBRURY, MA 01876
Product Made in China
M809-00

STONE AT ROOM TEMPERATURE
AVOID EXCESSIVE HEAT.

**THIS PRODUCT IS PACKAGED IN
A CHILD RESISTANT AND
TAPESEALED BUBBLE PACKAGING
ONLY IF ALLEGED AND INDICATED**

Drug Facts (continued)
Ask a doctor before use if you have:
• liver disease • heart disease • high blood pressure
• kidney disease • diabetes
• trouble swallowing or enlarged prostate gland
Ask a doctor or pharmacist before use if you are taking the
blood thinning drug warfarin.
When using this product, do not use more than directed.
Stop use and ask a doctor if:
• pain or nasal congestion does not improve after 7 days
• fever gets worse or lasts more than 3 days
• redness or swelling is present
• new symptoms occur
• you get nervous, dizzy or sleepless
If you get a headache, ask a health professional
to see you.
Keep out of reach of children. In case of overdose, get
medical help or contact a Poison Control Center right away.
Always use Aspirin when it may cause severe side reactions.
Symptoms may include: • dizziness • blurred vision •
stomach pain • nausea • vomiting • rash
It may interact with other drugs you are taking. Tell your
doctor about all the drugs you are taking, including over-the-
counter drugs.
Do not use with any other drug containing acetaminophen
(paracetamol) or aspirin. If you are not sure whether
a drug contains acetaminophen, ask a doctor or pharmacist.
If you are now taking a prescription medicine or other
medication (MAOI) (certain drugs for depression, psychiatric or
anxiety disorders, or Parkinson's disease), or for 2 weeks
after stopping the MAOI drug (You do not know if you
are taking this product), ask a doctor or pharmacist
before taking this product.
Do not use with any other drug containing acetaminophen
(paracetamol) or aspirin. If you are not sure whether
a drug contains acetaminophen, ask a doctor or pharmacist.
If you are now taking a prescription medicine or other
medication (MAOI) (certain drugs for depression, psychiatric or
anxiety disorders, or Parkinson's disease), or for 2 weeks
after stopping the MAOI drug (You do not know if you
are taking this product), ask a doctor or pharmacist
before taking this product.

Warnings
Save: liver damage may occur if you take
more than 4 doses in 24 hours, which is the maximum daily
amount for this product. • with other drugs containing
acetaminophen. • 3 or more alcohol drinks every day while
using this product.
Always use Aspirin when it may cause severe side reactions.
Symptoms may include: • dizziness • blurred vision •
stomach pain • nausea • vomiting • rash
It may interact with other drugs you are taking. Tell your
doctor about all the drugs you are taking, including over-the-
counter drugs.
Do not use with any other drug containing acetaminophen
(paracetamol) or aspirin. If you are not sure whether
a drug contains acetaminophen, ask a doctor or pharmacist.
If you are now taking a prescription medicine or other
medication (MAOI) (certain drugs for depression, psychiatric or
anxiety disorders, or Parkinson's disease), or for 2 weeks
after stopping the MAOI drug (You do not know if you
are taking this product), ask a doctor or pharmacist
before taking this product.

Uses
Temporarily relieves symptoms due to the common cold
• minor aches and pains • headache • fever
• nasal congestion • sinus congestion, pressure
• upper respiratory tract pain • headache
Temporarily relieves symptoms due to hay fever or other
• nasal congestion • sinus congestion, pressure
• minor aches and pains • headache • fever
Temporarily relieves symptoms due to the common cold
• minor aches and pains • headache • fever
• nasal congestion • sinus congestion, pressure
• upper respiratory tract pain • headache
Temporarily relieves symptoms due to hay fever or other

Directions
Take only as directed • do not exceed 4 doses per 24 hrs
adults and children 12 yrs & over 2 softgels with water
every 4 hrs
children 4 to under 12 yrs ask a doctor
children under 4 yrs do not use

Other Information
store at room temperature (59°-86°F)
Keep out of reach of children. In case of overdose, get
medical help or contact a Poison Control Center right away.
Always use Aspirin when it may cause severe side reactions.
Symptoms may include: • dizziness • blurred vision •
stomach pain • nausea • vomiting • rash
It may interact with other drugs you are taking. Tell your
doctor about all the drugs you are taking, including over-the-
counter drugs.
Do not use with any other drug containing acetaminophen
(paracetamol) or aspirin. If you are not sure whether
a drug contains acetaminophen, ask a doctor or pharmacist.
If you are now taking a prescription medicine or other
medication (MAOI) (certain drugs for depression, psychiatric or
anxiety disorders, or Parkinson's disease), or for 2 weeks
after stopping the MAOI drug (You do not know if you
are taking this product), ask a doctor or pharmacist
before taking this product.

Drug Facts (continued)
Ask a doctor before use if you have:
• liver disease • heart disease • high blood pressure
• kidney disease • diabetes
• trouble swallowing or enlarged prostate gland
Ask a doctor or pharmacist before use if you are taking the
blood thinning drug warfarin.
When using this product, do not use more than directed.
Stop use and ask a doctor if:
• pain or nasal congestion does not improve after 7 days
• fever gets worse or lasts more than 3 days
• redness or swelling is present
• new symptoms occur
• you get nervous, dizzy or sleepless
If you get a headache, ask a health professional
to see you.
Keep out of reach of children. In case of overdose, get
medical help or contact a Poison Control Center right away.
Always use Aspirin when it may cause severe side reactions.
Symptoms may include: • dizziness • blurred vision •
stomach pain • nausea • vomiting • rash
It may interact with other drugs you are taking. Tell your
doctor about all the drugs you are taking, including over-the-
counter drugs.
Do not use with any other drug containing acetaminophen
(paracetamol) or aspirin. If you are not sure whether
a drug contains acetaminophen, ask a doctor or pharmacist.
If you are now taking a prescription medicine or other
medication (MAOI) (certain drugs for depression, psychiatric or
anxiety disorders, or Parkinson's disease), or for 2 weeks
after stopping the MAOI drug (You do not know if you
are taking this product), ask a doctor or pharmacist
before taking this product.

Active Ingredients (in each softgel) Purpose
Acetaminophen 325 mg Pain reliever/Fever reducer
Phenylephrine HCl 5 mg Nasal decongestant

Glue

No Coating Area

No Coating Area

DAYTIME SINUS

acetaminophen, phenylephrine hydrochloride capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:510 13-157
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	orange (clear)	Score	no score
Shape	capsule (oblong)	Size	19mm
Flavor		Imprint Code	PC12
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:510 13-157-04	2 in 1 CARTON	07/26/2016	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/26/2016	

Labeler - PuraCap Pharmaceutical LLC (962106329)

Establishment

Name	Address	ID/FEI	Business Operations
Humanwell PuraCap Pharmaceutical (Wuhan) Co., Ltd.		421293287	manufacture(510 13-157) , analysis(510 13-157)

Revised: 1/2020

PuraCap Pharmaceutical LLC