

PEROXYGUARD 105- hydrogen peroxide spray
AgroChem Inc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).

Peroxyguard 105

PEROXYGUARD 105 SPRAY & FOAM

Teat Spray

WARNING

CAUSES SERIOUS EYE IRRITATION

Use Peroxyguard effectively to reduce the spread of mastitis causing organisms. Recommended for pre milking applications.

FOR FARM AND INDUSTRIAL USE ONLY.

KEEP OUT OF REACH OF CHILDREN.

KEEP OUT OF DIRECT SUNLIGHT.

AVOID FREEZING

Warning

Causes serious eye irritation.

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant

Storage: Store locked up.

INGREDIENTS:

ACTIVE INGREDIENTS:

Hydrogen Peroxide 1.0% (Stabilized)

Lactic Acid Salt - 0.5%

Glycerine - 5.0%

Directions:

General Use

Ready to use. Do not dilute. Do not mix product with water, any other teat dips, or other product.

Use a teat dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Teat Sprays can also be foamed or sprayed using recommended equipment.

Pre Dipping Use

Perform fore-stripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels

Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product.

Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product.

Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe.

If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking. Dipping should be resumed at least twice daily one week before calving.

PEROXYGUARD 105 TEAT SPRAY & FOAM

TEAT SPRAY

WARNING

**CAUSES SERIOUS EYE IRRITATION
FOR FARM AND INDUSTRIAL USE ONLY.
KEEP OUT OF REACH OF CHILDREN.
KEEP OUT OF DIRECT SUNLIGHT.
AVOID FREEZING**



PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

INGREDIENTS:

ACTIVE INGREDIENTS:

Hydrogen Peroxide 1.0% (Stabilized)

Lactic Acid Salt - 0.5%


Glycerine - 5.0%

Available with a glycerine emollient of 5% or 10%. PeroxyGuard is available as a dip, a spray & foam, or in a concentrate.

NET CONTENTS: 265 GALLON



38145



Use Peroxyguard effectively to reduce the spread of mastitis causing organisms. Recommended for pre milking applications.

Directions:

General Use

Ready to use. Do not dilute. Do not mix product with water, any other teat dips, or other product.

Use a teat dip cup to apply. Clean dip cup just prior to use. If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip. If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product. If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container. Teat Sprays can also be foamed or sprayed using recommended equipment.

Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis. Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product. Allow product to stand for 30 seconds. Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product. Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.

Dipping should be resumed at least twice daily one week before calving.



AgroChem

26 Freedom Way • Saratoga Springs, NY 12866 • (518)226-4850

PEROXYGUARD 105 TEAT SPRAY & FOAM

TEAT SPRAY

WARNING

CAUSES SERIOUS EYE IRRITATION
FOR FARM AND INDUSTRIAL USE ONLY.
KEEP OUT OF REACH OF CHILDREN.
KEEP OUT OF DIRECT SUNLIGHT.
AVOID FREEZING



PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

INGREDIENTS:

ACTIVE INGREDIENTS:

Hydrogen Peroxide 1.0% (Stabilized)

Lactic Acid Salt - 0.5%

Glycerine - 5.0%

Available with a glycerine emollient of 5% or 10%. PeroxyGuard is available as a dip, a spray & foam, or in a concentrate.

NET CONTENTS: 265 GALLON



38145

Use Peroxyguard effectively to reduce the spread of mastitis causing organisms. Recommended for pre milking applications.

Directions:

General Use

Ready to use. Do not dilute. Do not mix product with water, any other teat dips, or other product.

Use a teat dip cup to apply. Clean dip cup just prior to use. If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip. If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product. If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container. Teat Sprays can also be foamed or sprayed using recommended equipment.

Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis. Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product. Allow product to stand for 30 seconds. Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product. Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.

Dipping should be resumed at least twice daily one week before calving.



AgroChem

26 Freedom Way • Saratoga Springs, NY 12866 • (518)226-4850

PEROXYGUARD 105

hydrogen peroxide spray

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:17307-3813
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROGEN PEROXIDE (UNII: BBX060AN9V) (HYDROGEN PEROXIDE - UNII:BBX060AN9V)	HYDROGEN PEROXIDE	0.01 kg in 1 kg
LACTIC ACID (UNII: 33X04XA5AT) (LACTIC ACID - UNII:33X04XA5AT)	LACTIC ACID	0.005 kg in 1 kg

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	0.89 kg in 1 kg
GLYCERIN (UNII: PDC6A3C0OX)	0.05 kg in 1 kg

BEHENAMIDO PROPYLTRIMONIUM METHOSULFATE (UNII: 7MY9WEU70B)	0.02 kg in 1 kg
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	0.0009 kg in 1 kg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17307-3813-1	19 kg in 1 CONTAINER		
2	NDC:17307-3813-2	57 kg in 1 DRUM		
3	NDC:17307-3813-3	208 kg in 1 DRUM		
4	NDC:17307-3813-4	1040 kg in 1 CONTAINER		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/01/2013	

Labeler - AgroChem Inc (805427700)

Establishment

Name	Address	ID/FEI	Business Operations
AgroChem Inc		805427700	api manufacture, manufacture

Revised: 2/2018

AgroChem Inc