INFANT AND CHILDREN NOVEX ACETAMINOPHEN- acetaminophen suspension Herbion Pakistan (Pvt.) Ltd

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active ingredient

(in each 5 mL)

Acetaminophen 160 mg

Purpose:

fever reducer / Pain reliever

Uses:

temporarily: \blacksquare reduces fever \blacksquare relieves minor aches and pains due to: \blacksquare the common cold \blacksquare flu \blacksquare headache \blacksquare sore throat \blacksquare toothache

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if your child takes ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Overdose warning:

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions:

- this product does not contain directions or complete warnings for adult use.
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and use the dosing vial / measuring cup provided
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*	
Under 24	Under 2 years	Ask a doctor	
24-35	2-3 years	5 mL	
36-47	4-5 years	7.5 mL	
48-59	6-8 years	10 mL	
60-71	9-10 years	12.5 mL	
72-95	11 years	15 mL	

^{*}or as directed by a doctor

Attention: use only enclosed dosing devices provided, specifically designed for use with this product.

Other Information

- each 5 mL (1 tsp) contains: sodium 1.2 mg
- store between 20-25⁰C (68-77⁰F)
- do not use if bottle cap seal is broken

Inactive Ingredients:

Sucrose, Sorbitol, Glycerin, Flavor, Xanthan Gum, Sodium Benzoate, Potassium Sorbate, FD&C RED NO. 40, Purified Water

Questions or Comments?

Call 1-888-(OK-HERBION) 654-3724, Monday through Friday 9am - 5pm EST





Drug Facts

Active ingredient (in each 5 mL)

Purpose

Uses temporarily: • reduces fever • relieves minor temporarily: • reduces fever • relieves minor cold aches and pains due to: a the common cold a flu a headache a sore throat a toothache

Warnings
Liver warning: This product contains
acetaminophen. Severe liver damage may occur it
your child takes = more than 5 doses in 24 hours,
which is the maximum daily amount. • with other
drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

• skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

a with any other drug containing acetaminophen
tyrescription or nonprescription). If you are not
sure whether a drug contains acetaminophen, ask
a doctor or pharmacist.

a if your child is altergic to acetaminophen or any
of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if pain gets worse or lasts more than 5 days a fever gets worse or lasts more than 3 days a new symptoms occur a redness or swelling is present These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Posson Control Center right away. (1-800-222-1222) Quick medical attendos is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Drug Pacts (continued)

Directions

In this product does not contain directions or complete warnings for adult use,

do not give more than directed (see overdose warning)

shake well before using

mit. = milliter, tsp = teaspoonful

indiright dose on chart below. If possible, use weight to dose, otherwise, use age,

remove the child protective cap and use the dosing vial / measuring cup provided

repeat dose every 4 hours while symptoms tast

do not give more than 5 times in 24 hours

wedge (to)	MORE CITY	Dose (titr).	
under 24	under 2 years	ask a doctor	
24-35	2-3 years	5 mL	
36-47	4-5 years	7.5 mt.	
48-59	6-8 years	10 mL	
60-71	9-10 years	12.5 mL	
72-95	11 years	15 mL	

for as directed by a doctor

Attention: use only enclosed dosing devices provided, specifically designed for use with this product.

Other information:

• each 5est, contains; sodium 1,2 mg

• store between 20-25°C (68-77°F)

• do not use if bottle cap seal is broken

Inactive ingredients: Sucrose, Sorbitot, Glycerin, Flavor, Xanthan Gum, Sodium Benzoate, Potassium Sorbate, FDSC RED NO. 40, Purified Water

Questions or comments? call 1-888-(OK-HERBION) 654-3724, Monday through Friday, 9am - 5pm EST

Herbion

wile. NY 11801 5 (OK-HERBION) 664-3724

Lot No. Exp.

Barcode

INFANT AND CHILDREN NOVEX ACETAMINOPHEN

acetaminophen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:44237-040
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name Basis of Strength Strength				
ACETAMINO PHEN (UNII: 36209 ITL9D) (ACETAMINO PHEN - UNII: 36209 ITL9D)	ACETAMINOPHEN	160 mg in 5 mL		

Inactive Ingredients			
Ingredient Name	Strength		
SUCROSE (UNII: C151H8M554)			
SORBITOL (UNII: 506T60A25R)			
GLYCERIN (UNII: PDC6A3C0OX)			
XANTHAN GUM (UNII: TTV12P4NEE)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)			
WATER (UNII: 059QF0KO0R)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			

Product Characteristics				
Color	pink	Score		
Shape		Size		
Flavor	STRAWBERRY	Imprint Code		
Contains				

Packaging			
# Item Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:44237-040-	1 in 1 CARTON	03/09/2020	
1	120 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	03/09/2020	

Labeler - Herbion Pakistan (Pvt.) Ltd (645482936)

Registrant - Herbion Pakistan (Pvt.) Ltd (645482936)

Establishment

Name	Address	ID/FEI	Business Operations
Herbion Pakistan (Pvt.) Ltd		645482936	manufacture(44237-040)

Revised: 3/2020 Herbion Pakistan (Pvt.) Ltd