

**MCKESSON IBUPROFEN 200 MG- ibuprofen tablet, coated**  
**McKesson Medical-Surgical**

-----  
**McKesson Ibuprofen 200 mg**

***Drug Facts***

***Active ingredient (in each tablet)***

Ibuprofen 200 mg (NSAID\*).

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

Temporarily relieves minor aches and pains associated with

■ headache ■ toothache ■ backache ■ menstrual cramps

■ common cold ■ muscular aches ■ minor arthritis pain

Temporarily reduces fever.

***Warnings***

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

■ hives ■ skin reddening ■ asthma (wheezing) ■ facial swelling ■ rash ■ shock ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

■ are age 60 or older

■ have had stomach ulcers or bleeding problems

■ take a blood thinning (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ have 3 or more alcoholic drinks every day while using this product

■ take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better
- you have symptoms of heart problems or stroke • chest pain • trouble breathing • weakness in one part or side of body • slurred speech • leg swelling
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present in the painful area
- any new or unexpected symptoms occur

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- **do not use more than directed**
- **the smallest effective dose should be used**
- do not take longer than 10 days, unless directed by a doctor (see Warnings)

**Adults and children: (12 years and older)**

Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used.

Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

**Children under 12 years:**

Ask a doctor.

**Other information**

- read all product information before using
- store at 68-77°F (20-25°C)
- avoid excessive heat 104°F (above 40°C)
- tamper evident sealed packets
- do not use any opened or torn packets

### Inactive ingredients

colloidal silicon dioxide, corn starch, iron oxide red, polyethylene glycol, povidone (K-30), sodium starch glycolate, stearic acid, talc, titanium dioxide

**Questions or comments?** 1-800-777-4908

NDC 68599-8051-4

This Package is for Households without Young Children

McKesson

Ibuprofen

200 mg

Pain Reliever | Fever Reducer | (NSAID)

2 Per Pack | 200 Packs Per Box

MFR# 543-24805

Drug Facts		Drug Facts (continued)	
<b>Active ingredient (in each tablet)</b> Ibuprofen 200 mg (NSAID) <sup>1</sup>	<b>Purpose</b> Pain reliever/fever reducer	<b>When using this product</b> ● Use with food or milk to stomach upset occurs	<b>Stop use and ask a doctor if</b> ● you have had or are at risk for stomach upset occurs ● heart failure ● severe blood ● have bloody or black stools ● have stomach pain that does not get better ● you have symptoms of heart problems or stroke ● chest pain ● trouble breathing ● weakness in one part or one of body ● blurred vision ● ring around ● see spots or floaters in your ● have gone slow or fast for more than 2 days ● are not or getting worse in the genital area ● any new or unexplained symptoms occur
<b>Uses</b> Temporarily relieves minor aches and pains associated with: ● headache ● cold/flu ● sore throat ● menstrual cramps ● common cold ● muscle aches ● joint aches/pain Temporarily reduces fever	<b>Warnings</b> ● Allergy alerts: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ● hives ● difficulty breathing ● facial swelling ● rash ● dizziness ● asthma (wheezing) If an allergic reaction occurs, stop use and seek medical help right away. <b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause some stomach bleeding. The chance is higher if you: ● are age 60 or older ● have had stomach ulcers or bleeding problems ● take blood thinning (anticoagulant) or steroid drugs ● take other drugs, including prescription or over-the-counter (OTC) drugs (aspirin, ibuprofen, naproxen, or others) ● have had drug-induced peptic ulcers or gastrointestinal (GI) bleeding ● take more or for a longer time than directed Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. <b>Do not use</b> ● if you have ever had an allergic reaction to any other pain reliever/fever reducer ● right before or after heart surgery	<b>Directions</b> ● Do not use more than directed. ● If the medicine effective dose should be used (see directions). ● Take 1 tablet every 6 to 8 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor. <b>Children under 12 years:</b> Ask a doctor	<b>Other information</b> ● Read all of the information before using. ● Store at 68°F to 77°F (20° to 25°C). ● Do not store above 86°F (30°C). ● Do not use after the expiration date. ● Do not use any opened or torn packets. <b>Inactive ingredients</b> colloidal silicon dioxide, corn starch, iron oxide red, polyethylene glycol, povidone (K-30), sodium starch glycolate, stearic acid, talc, titanium dioxide
<b>Questions or comments?</b> Call 1-800-777-4908			

Distributed by McKesson Medical-Surgical Inc.  
Richmond, VA 23233  
PWN C0823  
Made in India or USA

MFR # 543-24805  
6 12479 26110 1

NDC 68599-8051-4

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

**McKesson**

**Ibuprofen**  
**200 mg**

PAIN RELIEVER | FEVER REDUCER | (NSAID)

2 PER PACK | 200 PACKS PER BOX | MFR # 543-24805

NDC 68599-8051-4

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

**McKesson**

**Ibuprofen**  
**200 mg**

PAIN RELIEVER | FEVER REDUCER | (NSAID)

2 PER PACK | 200 PACKS PER BOX | MFR # 543-24805

NDC 68599-8051-4

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

**McKesson**

**Ibuprofen**  
**200 mg**

PAIN RELIEVER | FEVER REDUCER | (NSAID)

2 PER PACK | 200 PACKS PER BOX | MFR # 543-24805

## MCKESSON IBUPROFEN 200 MG

ibuprofen tablet, coated

Product Information			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68599-8051
<b>Route of Administration</b>	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
Ibuprofen	200 mg		

<b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg
---	-----------	--------

### Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	

### Product Characteristics

<b>Color</b>	red (Reddish Brown)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	G;2
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68599-8051-4	200 in 1 BOX	09/01/2021	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	09/01/2021	

**Labeler** - McKesson Medical-Surgical (023904428)

**Registrant** - Unifirst First Aid Corporation (832947092)